



Calcareous Nannofossils from the Shiranish Formation at Bekhaer anticline, Dhouk area, Northern Iraq

Omar A. Al-Badrani, Mahfoud A. Al-Hadeedy, Ibrahim Y. Al-Shareefi

Department of Geology, College of Science, Mosul University, Mosul, Iraq

DOI: <http://dx.doi.org/10.25130/tjps.27.2022.035>

ARTICLE INFO.

Article history:

-Received: 8 / 1 / 2022

-Accepted: 14 / 4 / 2022

-Available online: / / 2022

Keywords: Nannofossils; Shiranish Formation; Nannobiostratigraphy; Bekhare.

Corresponding Author:

Name: Ibrahim Y. Al-Shareefi

E-mail:

ibrahimshareefi@uomosul.edu.iq

Tel:

ABSTRACT

Calcareous nannofossils are examined up on the Shiranish Formation from the Bekhaer anticline in Northern Iraq. The study area includes about 55 meters to find calcareous nannofossils assemblages, from which (36) species have identified.

Three biozones are postulated based on the above studied assemblages, in order of biozones: *Tranolithus phacelosus* biozone; *Reinhardtites levis* biozone; *Arkhangelskiella cymbiformis* biozone. From a regional perspective, these biozones were connected with other calcareous nannofossils biozones, leading to the conclusion that the section is Late Campanian-Maastrichtian in age.

1. Introduction

Henson in (1940) was firstly described the Shiranish Formation in an unpublished report, near the type section that is located near the village of Shiranish Islam near Zakho in northern Iraq [1]. It is represented by a section exposed on the surface with a thickness approximately 227.8 m. It was divided into two rock units, the upper unit is 99 m thick, which is consisted of blue marl, and the lower unit is 128.8 m thick, which is consisted of dark-blue and black marly limestone. Henson (1940) in [1] explained that the age of the formation is the Maastrichtian at the top of the section, while the lower part of it represents the late Campanian age.

After the marine regression that occurred at the end of the Cenomanian - Campanian cycle, a wide transgression occurred in most parts of Iraq and continued until the end of the Maastrichtian age. The reason for this was attributed to the diminishing or ending of the Sub-Hercynian tectonic movement within the inner parts of the Eugeosyncline [2]. This transgression led to the flooding of the whole of Kurdistan region, northern Iraq, as well as other neighboring regions such as southern Turkey and eastern parts of Syria, in addition to western Iran. This inundation with water led to the result of diverse sediments and their distribution over most of the area of the sedimentary basin, forming a sedimentary form

that differs greatly from the previous stages. [3] mentioned that Laramide Orogeny is responsible for these events and that as a result of it, three environments have developed in this cycle, the deep sedimentary basin environment and the neritic environment on the shelf, and these two environments overlap and add to them the deeper environment affected by flesh deposits.

This formation has been studied by many researchers with different specializations, where [4, 5] studied it with other formations based on fossils of Ostracoda, foraminifera, and some other groups. The formation was studied by relying on foraminifera fossils, such as [6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19]. The formation based on calcareous nannofossils has also been studied by [20, 21, 22].

The study section is located within the Bekhaer anticline, in the Dohuk governorate, northern Iraq, (Fig.1). It is represented by an outcrop section exposed to the surface with a thickness of 55 meters that was studied through 29 rock samples elicited from this section (Fig. 2). The lithology of the section from the bottom is limestone with intercalation marls separating it from the hard limestone detritus rocks below it, which represented by the Bekhme Formation. An unconformity is clear between them. Then the lithology of the formation starts with rock

sample no. 15, which is represented by marl with intercalation chalky limestone until it reaches the shale and sandstone beds that are representative of the Kolosh Formation. The contact between both formations is unconformable.

The aim of the current study is to identify the calcareous nannofossils assemblages that are

distributed within this formation, and then determine the biozones that represent the formation based on these fossils genera/species and compare them with previous studies from different regions. Also, the research aims to conclude the age of the formation accurately.




Fig.1: Paleogeographic map of Iraq; it shows the location of the study section.
From [23]

2. Materials and methods

We have collected twenty nine samples of stratigraphic successions from the marls and the marly limestone, to studying the calcareous nannofossils taxon using transmitted- light microscope, the specimens are extracted using the method-H [24].

It is an extraction method for microfossils that can be properly examined, when it is extracted from the rocks, each paleontologist tends to have favorite methods for these procedures. The sample preparation is decanting and smears slides that

provide method for producing slides of calcareous nannofossils by small amount of the disaggregated sample is placed in distilled water and a drop of cello size added to act as a dispersant. The cover slip is left to dry on a warm hotplate. To make permanent mounts allow the slide and residue to dry at a low temperature away from possible sources of contamination. Place a drop of mounting medium (e.g. Canada balsam) on a clean cover slip and drop this over the residue. Allow to dry before examining with transmitted light.




Fig. 2: Lithostratigraphic column of the Shiranish Formation from Bekhaer anticline, Northern Iraq.

3. Results

A Detailed systematic study of calcareous nannofossils has been carried out for the Shiranish Formation, Thirty six genera/species has determined as following (Figs. 3, 4 and 5):

1 - Heterococcoliths

Family Chiastozygaceae [25]

Genus Bukyrlithus [26]

Bukyrlithus ambiguus [26] (Fig.3a)

Genus Chiastozygus [27]

Chiastozygus platyrhethus [28] (Fig.3b)

Genus Placozygus [29]

Placozygus fibuliformis ([30]) [29] (Fig.3c)

Genus Reinhardtites [31]

Reinhardtites anthophorus ([32]) [31] (Fig.3d)

Reinhardtites levis Prins and Sissingh, in [33] (Fig.3e)

Genus Tranolithus [34]

Tranolithus phacelosus [35] (Fig.3f)

Family Eiffellithaceae [36]

Genus Eiffellithus [36]

Eiffellithus gorkae [36] (Fig.3g)

Family Stephanolithiaceae [37]

Genus Cylindralithus [38]

Cylindralithus nudus [39] (Fig.3h)

Family Axopodorhabdaceae [40]

Genus Cribrosphaerella Deflandre in [41]

Cribrosphaerella ehrenbergii (Arkhangelsky) Deflandre in [41] (Fig.3i)

Genus Nephrolithus [42]

Nephrolithus sp. (Fig.3j)

Family Prediscosphaeraceae [43]

Genus Prediscosphaera [44]

Prediscosphaera grandis [45] (Fig.3k)

Prediscosphaera sp. (Fig.3l)

Family Cretarhabdaceae [46]

Genus Retecapsa [26]

Retecapsa angustiforata [26] (Fig.4a)

Retecapsa crenulata [38] Grün in: [47] (Fig.4b)

Family Watznaueriaceae [25]

Genus Watznauia [30]

Watznaueria barnesae ([48]) [31] (Fig.4c)

Watznaueria biporta [39] (Fig.4d)

Family Arkhangelskiellaceae [39]

Genus Arkhangelskiella [44]

Arkhangelskiella cymbiformis [44] (Fig.4e)

Arkhangelskiella specillata [44] (Fig.4f)

Genus Broinsonia [39]

Broinsonia parca ([49]) [39] (Fig.4g)

Broinsonia signata ([50]) [51] (Fig.4h)

2 - Holococcoliths

Family Calyptosphaeraceae [52]

Genus *Calculites* [33]

Calculites ovalis ([49]) Prins and Sissingh in: [33] (Fig.4i)

Calculites obscurus ([32]) Prins and Sissingh in: [33] (Fig.4j)

3 -Nannoliths

Family Microrhabdulaceae [53]

Genus *Lithraphidites* [53]

Lithraphidites praequadratus [54] (Fig.4k)

Lithraphidites quadratus [38] (Fig.4l)

Genus *Microrhabdulus* [32]

Microrhabdulus decoratus [32] (Fig.5a)

Microrhabdulus undosus [55] (Fig.5b)

Family Polycyclolithaceae [56]

Genus *Eprolithus* [34]

Eprolithus floralis ([57]) [34] (Fig.5c)

Eprolithus sp. (Fig.5d)

Genus *Lithastrinus* [58]

Lithastrinus sp. (Fig.5e)

Genus *Micula* [44]

Micula staurophora ([59]) [49] (Fig.5f)

Micula swastica [60] (Fig.5j)

Genus *Radiolithus* [34]

Radiolithus planus [34] (Fig.5h)

Genus *Uniplanarius* [61]

Uniplanarius trifidus (Stradner In [58]) Prins and Perch-Nielsen in [62] (Fig.5i)

Uniplanarius sp. (Fig.5j)

Genus *Ceratolithoides* [38]

Ceratolithoides aculeus ([58]) Prins and Sissingh in [33] (Fig.5k)

Ceratolithoides sesquipedalis [63] (Fig.5l)




Fig.3: Cross-polarized micrograph of significant calcareous nannofossil taxon from the Shiranish Formation. (a) *Bukyrlithus ambiguus*; (b) *Chiastozygus platyrhethus*; (c) *Placozygus fibuliformis*; (d) *Reinhardtius anthophorus*; (e) *Reinhardtius levis*; (f) *Tranolithus phacelosus*; (g) *Eiffellithus gorkae*; (h) *Cylindralithus nudus*; (i) *Cribrosphaerella ehrenbergii*; (j) *Nephrolithus* sp.; (k) *Prediscosphaera grandis*; (l) *Prediscosphaera* sp.




Fig. 4: Cross-polarized micrograph of significant calcareous nannofossil taxon from the Shiranish Formation. (a) *Retecapsa angustiforata*; (b) *Retecapsa crenulata*; (c) *Watznaueria barnesae*; (d) *Watznaueria biporta*; (e) *Arkhangelskiella cymbiformis*; (f) *Arkhangelskiella specillata*; (g) *Broinsonia parca*; (h) *Broinsonia signata*; (i) *Calculites ovalis*; (j) *Calculites obscurus*; (k) *Lithraphidites praequadratus*; (l) *Lithraphidites quadratus*.




Fig. 5: Cross-polarized micrograph of significant calcareous nannofossil taxon from the Shiranish Formation. (a) *Microrhabdulus decoratus*; (b) *Microrhabdulus undosus*; (c) *Eprolithus floralis*; (d) *Eprolithus* sp.; (e) *Lithastrinus* sp.; (f) *Micula staurophora*; (g) *Micula swastica*; (h) *Radiolithus planus*; (i) *Uniplanarius trifidus*; (j) *Uniplanarius* sp.; (k) *Ceratolithoides aculeus*; (l) *Ceratolithoides sesquipedalis*.

4. Discussions

Stratigraphic distribution of the recorded genera/species reveals three biozones (Fig. 6) and correlated with local and regional previous studies (Figs. 7 and 8), These biozones are :

1 - *Tranolithus phacelosus* [35] Interval Biozone (CC 23)

Identification: It is Interval biozone for *Tranolithus phacelosus* [35], recorded by the Last Appearance Datum (LAD) of the species *Eiffilithus eximus* Reinhardt (1965), and extends to the Last Appearance Datum (LAD) of the species *Tranolithus phacelosus* [35], and the thickness of this biozone is about (18 meters), within the lower part of the Shiranish Formation

Correlation: This zone is compared with CC23 (*Tranolithus phacelosus*) zone which studied by the [33] that aged of the late Campanian to early Maastrichtian, and compared with UC16, UC17 zone that is studied by [40], that aged Campanian to early Maastrichtian age. On local scale its compared with *Tranolithus phacelosus* zone that is recorded by the [22, 65, 66, 67], that is also extends un age from Campanian to early Maastrichtian. Its also recorded as index of zone CC23 from the lower part of the Tanjero Formation, Sulaimani area by [68]. Accordingly, the suggested age for this zone extends late Campanian to early Maastrichtian based on geological time scale by [69].

2- *Reinhardtites levis* [33] Interval Biozone (CC24)

Identification: It is Interval biozone of the species *Reinhardtites levis* [33], recorded by Last Appearance Datum (LAD) of the species *Tranolithus phacelosus* [35], to the Last appearance Datum(LAD) of *Reinhardtites levis* [33]. The thickness of this biozone is about (12 meters). Within the middle part of the studied section of the Shiranish Formation

Correlation: This zone is compared with CC24 (*Reinhardtites levis*) zone that studied by the [33] which is indicating an age extends from the late Campanian to early Maastrichtian, and compared with the UC18 zone that is studied by [40] that aged Maastrichtian age; and compared with *Reinhardtites levis* zone that studied by the [22, 65, 66], that aged of the early Maastrichtian. Its also recorded by [68] from the lower part of Tanjero Formation in sulaimnai area. Based on this correlation, this interval biozone assigned to be as early Maastrichtian in age [69].

3- *Arkhangelskiella cymbiformis* [44] Interval Biozone (CC25)

Identification: It is Interval biozone of the species *Arkhangelskiella cymbiformis* [44], recorded by Last Appearance Datum (LAD) *Reinhardtites levis* [33] to the First Appearance Datum (FAD) *Nephrolithus frequens* [42]. The thickness of this biozone IS about (25 meters) and recorded from the upper part of Shiranish formation in the studied section

Correlation: This biozone is compared with (CC25) (*Arkhangelskiella cymbiformis*) zone by [33] , which is subdivided into three subzones namely (CC25a, CC25b, CC25c), that is based on the the first appearance of the species *Arkhangelskiella cymbiformis* and the first appearance of species *Lithraphidites quadratus* at Maastrichtian age, and compared with the UC19 zone that is studied by [40] that aged Late Maastrichtian age. IN Kurdistan region this zone also compared with *Arkhangelskiella cymbiformis* zone by the [65] which is assigned to be Maastrichtian in age ; therefore we suggested the Maastrichtian [69].

Its worthy to mention neither late Maastrichtian (CC26) nor Danian calcareous nannofossils (CP.1) biozones are recognized in the studied section of Beakher anticline, Dhok area, Kurdistan region, (N. Iraq)

5. Conclusions

The study of the Shiranish Formation at Bekhair anticline, Dhok area, Kurdistan region, Northern Iraq suggested that:

1 - Presence of three calcareous nannofossils, these are from oldest to youngest;

a. *Tranolithus phacelosus* Interval zone (CC23)

b. *Reinhardtites levis* Interval zone (CC24)

c. *Arkhangelskiella cymbiformis* Interval zone (CC25).

2 - The recorded three biozone are point to Campanian to late Maastrichtian age, but with remarkable disappearance of the latest Maastrichtian calcareous nannofossils *Nephrolithus frequens* (CC26), as well as the Danian calcareous nannofossils zone (CP1).

3 - The disappearances mostly point to a gap and unconformity between the Shiranish Formation and overlying Kolosh Formation (Paleocene), which is mostly, indicates the terminations of the Tectonic Megasequence AP9 by the influence of the tectonic activity at that time.




Fig.6: Calcareous nannofossils range chart for the Shiranish Formation from the studied area.

Age (m.y.)		Period	
65 –		Paleogene	
66 –		Paleocene	
67 –		Danian	
68 –		Maastrichtian	
69 –		Late	
70 –		Cretaceous	
71 –		Campanian	
72 –		E	
73 –		L	
74 –		Campanian	
75 –		E	
76 –		L	
Gradestien et al., 2012		CP 1	
Nephrolithus frequens CC26		Arhangelskiella cymbiformis CC25	
Reinhardtius levis CC24		Tranolithus phacelosus CC23	
Uniplanarius trifidus CC22		Campanian	
Sisssingh, 1977		CP 1	
Nephrolithus frequens CC26		Arhangelskiella cymbiformis CC25	
Reinhardtius levis CC24		Tranolithus phacelosus CC23	
Uniplanarius trifidus CC22		Campanian	
Bown, 1998		UC 20	
UC 19		UC 18	
UC 17		UC 16	
Present work		CP 1	
Nephrolithus frequens CC26		Arhangelskiella cymbiformis CC25	
Reinhardtius levis CC24		Tranolithus phacelosus CC23	
Uniplanarius trifidus CC22		Campanian	

Fig .7: Regional correlation chart for the Cretaceous calcareous nannofossils biozones

References

- [1] **Bellen, R.C. Van, Dunnington, H.V., Wetzel, R. and Morton, D.M. (1959).** Lexique stratigraphique international, V.III: Asie, Fascicule 10 a, Iraq. 333 p.
- [2] **Al-Sayab, A. S., Al-Omari, F. S., Al-Ansari, N., Al-Sheikh, Z., Al-Rawi, D., Ali, J. (1982).** Geology of Iraq. Dar Alkotob for printing and publishing. Pp. 280.
- [3] **Chatton, M. and Hart, E. (1961).** Review of the Cenomanian to Maastrichtian stratigraphy in Iraq. Iraq. Pet. Co., Limited, Palaeontology Dep. Kirkuk, 75 p.
- [4] **Al-Shareefi, I.Y. (2004).** Biostratigraphy of Ostracode and Sedimentological study of some upper Cretaceous formation from selected wells north west and middle Iraq. Unpub. Ph.d thesis, Univ. Mosul,:1 - 142.
- [5] **Al-Shareefi, I.Y., Khalaf, S.K. and Al-Eisa, M.A. (2014).** Ostracode Biostratigraphy of Shiranish, Hartha and Mushorah Formations from Selected Boreholes Northwest and Central Iraq. Iraqi National Journal of Earth Sciences, Vol.14, No.1, p. 19 – 38.
- [6] **Al-Omari, F. S. (1970).** Upper Cretaceous and Lower Cenozoic foraminifera of three oil wells in northwestern Iraq. Univ. Missouri at Rolla, Unpublished Ph. D. Thesis. 280 p.
- [7] **Kassab, I.I. (1973).** Planktonic foraminifera of the Shiranish Formation type locality (Northern Iraq). Jour. Geol. Soc. Iraq, Vol. 6 (Al-Naqib Vol.), PP.100-109.
- [8] **Kassab, I.I. (1975).** Biostratigraphy study of the subsurface Upper Cretaceous-Lower Tertiary of Well Injana No.5, Northeast Iraq. Jour. Geol. Soc. Iraq, Spec. Iss, PP.181-199.
- [9] **Al-Safawee, N. M. (1980).** Foraminiferal study of the Mesozoic – Cenozoic transition of Sasan well no.1, Northwestern Iraq. Unpublished M.Sc. Thesis, Univ. of Mosul, 146 p.
- [10] **Abawi, T. S., Abdel-Kireem, M. R. and Yousef, J. M. (1982).** Planktonic Foraminiferal stratigraphy of Shiranish Formation, Sulaimaniah – Dokan region, northeastern Iraq. Rev. Esp. Micropal., Vol. 14, No. 1, PP. 153 – 164.
- [11] **Abdel-Kireem, M. R. (1983).** A study of the paleoecology and bathymetry of the Foraminiferal assemblages of the Shiranish Formation (Upper Cretaceous) northeastern Iraq. Paleogeol. Paleocli. Paleoeco., Vol. 43, PP. 169 – 180.
- [12] **Hammoudi, R. A. (2000).** Planktonic foraminiferal biostratigraphy of the Shiranish Formation (Upper Cretaceous) in Jambur Well No. 13, Northern Iraq. Raf. Jour. Sci., Vol. 11, No. 4, PP. 50 – 58.
- [13] **Bamerni, A. A. Q. (2010).** Biostratigraphy of the Shiranish Formation in the subsurface section of Khanke well no.1, Duhok area, North of Iraq. Unpub. M.Sc. thesis, Univ. Mosul,:1 -111.
- [14] **Al-Juboury, F. N. H. (2011).** Planktonic Foraminiferal Biostratigraphy, Depositional Environment and Sequence Stratigraphy of Upper Cretaceous - Middle Eocene Sequences at selected Wells in Khabbaz Oil Field Northeastern Iraq. Unpub. Ph.d thesis, Univ. Mosul,:1 -198.
- [15] **Ameen, F. A., and Gharaib, H. (2014).** Biostratigraphy of the Tethyan cretaceous successions from northwestern Zagros fold-thrust belt, Kurdistan region, NE Iraq. Arabian Journal of Geosciences, 7(7), 2689-2710.
- [16] **Jaff, R. B. N., Williams, M., Wilkinson, I. P., Lawa, F., Lee, S., Zalasiewicz, J. (2014).** A refined foraminiferal biostratigraphy for the Late Campanian-Early Maastrichtian succession of northeast Iraq. Geo Arabia, 19 (1), 161-180.
- [17] **Jaff, R. B. N., Wilkinson, I. P., Lee, S., Zalasiewicz, J., Lawa, F., Williams, M. (2015).** Biostratigraphy and palaeoceanography of the early Turonian-early Maastrichtian planktic foraminifera of northeast Iraq. Journal of Micropalaeontology, 34 (2), 105-138.
- [18] **Jaff, R. B. N. & Lawa, F. A. (2019).** Palaeoenvironmental signature of the Late Campanian-Early Maastrichtian benthonic foraminiferal assemblages of Kurdistan, Northeast Iraq. Journal of African Earth Sciences, 151, 255–273.
- [19] **Jaff, R. B.N. and Lawa, F. A.A. (2020).** Biostratigraphy and Systematic Palaeontology of Late Cretaceous Heterohellicidae Foraminifera from Kurdistan Region Northeastern Iraq. Iraqi National Journal of Earth Sciences, Vol. 20, No. 2, pp. 33-63.
- [20] **Al-Badrani, O. A. & Mahmood, R. M. (2014).** Study of nannoliths from upper Cretaceous Shiranish Formation, Higran area, Northern Iraq. Arab. Jour. Geosci. 7:5571–5578.
- [21] **Al-Shareefi, I., Al-Badrani O., and Yousif A. (2019).** Micropaleontological Study of the contact between Kometan and Shiranish Formations from Dokan area, Northern Iraq. In: Mabrouk Boughdiri *et al.* (eds). Paleobiodiversity and Tectono-Sedimentary Records in the Mediterranean Tethys and Related Eastern Areas. 117-119.
- [22] **Al-Badrani O.A., and Al-Shareefi, I.Y. (2020).** Calcareous Nannofossils Biostratigraphy of the Shiranish Formation from Ajeel-10 well, North Iraq. Iraqi Bulletin of Geology and Mining Vol.16, No.2, 2020 p 21 – 31.
- [23] **Jassim, S. Z. and Goff, J. C. (2006).** Geology of Iraq, Published by Dolin, Prague and Moravian Museum, Brno, pp. 179-181.
- [24] **Armstrong, H. and Brasier, M. (2005).** Microfossils Black well publishing. 296p.
- [25] **Rood, A. P., Hay, W. W. & Barnard, T. (1973).** Electron microscope studies of Lower and Middle Jurassic coccoliths. Eclogae Geologicae Helvetiae. 66: 365-382. gs V O
- [26] **Black, M. (1971).** Coccoliths of the Speeton Clay and Sutterby Marl. Proceedings of the Yorkshire

Geological Society, vol. 38, part 3, no. 18, pp. 381-424.

[27] **Gartner, S. (1968).** Coccoliths and related calcareous nannofossils from Upper Cretaceous deposits of Texas and Arkansas. University Kansas Paleontology Contribution-Protista, Article 1, pp. 1-56.

[28] **Hill, M.E. (1976).** Lower Cretaceous calcareous nannofossils from Texas and Oklahoma. *Palaeontologica B*, 156, 103-79.

[29] **Hoffmann, N. (1970).** Coccolithineen aus der weissen Schreiekreide (Unter Maastricht) von Jasmund auf Rügen. *Geologie*, 19, 846-79.

[30] **Perch-Nielsen, K. (1968).** Der feinbau und die klassifikation der coccolithen aus dem Maastrichtien von Danemark. *Kong. Danske videnskab. Selskab., Biol. Skr.*, vol. 16, pp. 1-96.

[31] **Sissingh, W. (1977).** Biostratigraphy of Cretaceous calcareous nannoplankton. *Geol. Mjnb.*, vol. 56, pp. 37-65.

[32] **Stover, E. (1966).** Cretaceous coccoliths and associated nannofossils from France and Netherlands. *Micropaleontology*, vol. 12, no. 2, pp. 133-167.

[33] **Reinhardt, P. (1966).** Zur taxonomie und biostratigraphie des fossilen nannoplanktons aus dem malm, der kreide und dem tertiär Mitteleuropas. *Freiberger Forschungshefte, Paläontologie, Series C*, 196, pp. 513-524.

[34] **Reinhardt, P. (1965).** Neue familien für fossile kalkflagellaten (Coccolithophoridae, Coccolithineae). *Monatsber. Deutsch. Akad. Wiss. Berlin*, vol. 7, pp. 30-40.

[35] **Black, M. (1968).** Taxonomic problems in the study of coccoliths. *Palaeontology*, 11 (5), 793-813.

[36] **Bramlette, M. N., and Martini, E. (1964).** The great change in calcareous nannoplankton fossils between the Maastrichtian and Danian. *Micropaleontology*, vol. 10, no. 3, pp. 291-322, pls. 1-7.

[37] **Bukry, D. (1969).** Upper Cretaceous coccoliths from Texas and Europe. *The University Kansas Paleontology Contribution, Article 51, (Protista 2)*, pp. 1-79.

[38] **Bown, P.R. and Young, J.R. (1997).** Mesozoic calcareous nannoplankton classification. *Journal of Nannoplankton Research*, vol. 19, no. 1, pp. 21-36.

[39] **Piveteau, J. (1952).** *Traité de Paléontologie*, vol. I, pp. 107-15. Masson, Paris.

[40] **Górka, H. (1957).** Les coccolithophoridés du Maastrichtien supérieur de Pologne. *Acta Palaeont. Polonica*, vol. 2, no. 2-3, pp. 235-284.

[41] **Rood, A.P., Hay, W.W., & Barnard, T. (1971).** Electron Microscope Studies of Oxford Clay Coccoliths. *Eclogae geologicae Helveticae*, 64, 245-272.

[42] **Vekshina, V. N. (1959).** Coccolithophoridae of the Maastrichtian deposits of the West Siberian lowland. *Trudy Sibir. Nauch-Issled. Inst. Geol. Geofiz. Min. Syrja*, vol. 2, pp. 56-81.

[43] **Perch-Nielsen, K. (1979).** Calcareous nannofossils from the Cretaceous between the North Sea and the Mediterranean. *Aspekte der Kreide Europas, IUGS Series A*, vol. 6, pp. 223-272.

[44] **Thierstein, H. R. (1973).** Lower Cretaceous calcareous nannoplankton biostratigraphy. *Abh. Geol. B. -A.* vol. 29, p. 52.

[45] **Grün, W., & Allemann, F. (1975).** The Lower Cretaceous of Caravaca (Spain). Berriasian Calcareous Nannoplankton of the Miravetes Section (Subbetic Zone, Prov. of Murcia). *Eclogae geologicae Helveticae*, 68, 147-211.

[46] **Reinhardt, P. (1964).** Einige kalkflagellaten - gattungen (Coccolithophoridae, Coccolithineae) aus dem Mesozoikum Deutschlands. *Monatsber. Deutsch. Akad. Wiss. Berlin*, vol. 6, pp. 749-759.

[47] **Black, M. and Barnes, B. (1959).** The structure of Coccoliths from the English Chalk. *Geological Magazine*, 96: 321-328.

[48] **Stradner, H. (1963).** New contributions to Mesozoic stratigraphy by means of nannofossils. *Proceedings of the Sixth World Petroleum Congress, Section 1 Paper 4*, pp. 167-183.

[49] **Noël, D. (1969).** Arkhangelskiella (coccolithes cretaces) et formes affines du Bassin de Paris. *Rev. Micropaleontol.*, 11 (4) 191-204.

[50] **Noël, D. (1970).** Coccolithes Cretaces. I. *ij Craie Campanienne, du Bassin de Paris. CNRS, Paris*. 129 pp

[51] **Boudreaux, J. E. & Hay, W. W. (1969).** Calcareous nannoplankton and biostratigraphy of the Late Pliocene-Pleistocene-Recent sediments of the Submarex cores. *Rev. Esp. Micropaleontol.*, 1, 249-92.

[52] **Deflandre, G. (1959).** Sur les nannofossiles calcaires et leur systematique. *Rev. Micropaleontology*, vol. 2, no. 3, pp. 127-152.

[53] **Deflandre, G. (1963).** Sur les Microrhabdulidés, famille nouvelle de nannofossiles calcaires. *C.R. Acad. Sci., Paris*, no. 256, pp. 3484-3486.

[54] **Roth, P. H. (1978).** Calcareous nannoplankton biostratigraphy and oceanography of the Northwestern Atlantic ocean. In *Benson, W. E., Sheridan, R. E., et. al., Initial reports of the Deep sea drilling project. Washington (U.S. Government Printing office)*, vol. 44, pp. 731-759.

[55] **Perch-Nielsen, K. (1973).** Neue coccolithen aus dem Maastrichtien von Dänemark, Madagaskar und Ägypten. *Bull. geol. Soc Denmark*. Vol. 22, pp. 306-333.

[56] **Varol, O. and Girgis, M. (1994).** New taxa and taxonomy of some Jurassic to Cretaceous calcareous nannofossils. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, 192: 221-253.

[57] **Stradner, H. (1962).** Über neue und wenig bekannte Nannofossilien aus Kreide und Alttertiär. *Sonderabdruck aus den Verhandlungen der Geologischen Bundesanstalt*, vol. 2, pp. 363-377.

[58] **Stradner, H. and Papp, A. (1961).** Tertiäre Discoasteriden aus Österreich und deren stratigraphische Bedeutung mit Hinweisen auf

Mexico, Rumanien und Italien. Jahrbuch der Geologischen Bundesanstalt (Wien), vol.7, pp. 1-159.

[59] Gardet, M. (1955). Contribution à l'étude des coccolithes des terrains néogènes de l'Algérie. Publications du Service de la Carte Géologique de l'Algérie (Nouvelle Série), vol. 5, pp. 477-550.

[60] Stradner, H. and Stein metz, J. (1984). Cretaceous calcareous nannofossils from the Angola Basin, Deep Sea Drilling Project Site 530. Initial Reports of the Deep Sea Drilling Project, vol. 75, pp. 565-649.

[61] Hattner, J.G. and Wise, J.R. (1980). Upper Cretaceous calcareous nannofossil biostratigraphy of south Carolina. South Carolina. Geol., Vol.24, No.2, p. 44 – 117.

[62] Manivit, H., Perch-Nielsen, K., Prins, B. and Verbeek, J.W. (1977). Mid Cretaceous calcareous nannofossil biostratigraphy. Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen, B80(3), pp. 169-181.

[63] Burnett, J.A. (1997). New species and new combinations of Cretaceous nannofossils, and a note on the origin of Petrarhabdus (Deflandre) Wind and Wise. Journal of Nannoplankton Research, vol.19, no. 2, pp. 133-146.

[64] Burnett, J.A. (1998). Upper Cretaceous. In: Bown, P.R. (Editor), Calcareous Nannofossil

Biostratigraphy. British Micropalaeontological Society Publications Series. Chapman and Hall, London, pp. 132-199.

[65] Al-Maamari, A.M. and Al-Badrani, O.A. (2019). Calcareous Nannofossils Biostratigraphy of Shiranish Formation (K-306) well, Northern Iraq. Iraqi National Journal of Earth Sciences Vol.19, No.2, p. 1 – 10.

[66] Al-Badrani, O. A. (2012). Nannobiostratigraphy of the Lower part of Shiranish Formation, Sinjar Anticline, NW Iraq. Iraqi National Journal of Earth Sciences, Vol. 12, No. 1, pp. 1-16.

[67] Al-Badrani, O.A. and Al-Assaf, E.N. (2011). Nannobiostratigraphy of Shiranish Formation in Balad Well No. 8, Northern Baghdad, Iraq. Iraqi National Journal of Earth Sciences, Vol. 11, No. 2, pp. 65 – 80.

[68] Al-Sulivani, R. F. and Al-Badrani, O.A. (2021). Calcareous Nannofossils Biostratigraphy of Tanjero Formation at Azmer Anticline, Sulimaniya, Northern Iraq. Iraqi Journal of Science, Vol.62, No. 5, pp: 1605-1621.

[69] Gradstein, F. M., Ogg, J. G., Schmitz, M. D. and Ogg, G., M. (2012). The geologic time scale. The Boulevard, Langford lane, Kidlington, Oxford OX5 1GB, UK, vol. 2, pp. 437-1144.

متحجرات النانو الكلسية من تكوين شرانش في طية بيخير المحدبة، منطقة دهوك، شمال العراق

عمر احمد البدراني ، محفوظ عبدالله.الحديدي ، إبراهيم يونس الشريفي

قسم علوم الارض ، كلية العلوم ، جامعة الموصل ، الموصل، العراق

الملخص

تم دراسة متحجرات النانو الجيرية في تكوين شرانش من طية بيخير المحدبة في منطقة دهوك في شمال العراق. بلغ سمك المقطع المدروس حوالي 55 مترًا ، حيث تم ملاحظة تجمعات لمتحجرات النانو الجيرية وامكن تحديد 36 نوعًا.

تم تحديد ثلاثة انطقة حياتية بناءً على التجمعات المدروسة أعلاه وهي مرتبة من الاقدم الى الاحدث كالتالي:

Tranolithus phacelosus biozone, *Rienhardtites lives* biozone, *Arkhangelskilla cymbiformis* biozone.

من منظور المقارنة الإقليمية ، تم ربط هذه الانطقة الحياتية مع انطقة حياتية أخرى لمتحجرات النانو الجيرية ، مما أدى إلى استنتاج أن المقطع المدروس هو بعمر الكامبانيان – الماسترختيان المتأخر.