


Chemical profiling and antimicrobial potential of crude extract and essential oil from *Conyza bonariensis* leaves

Simeon P. Nsindagi , Cyprian B. Mpinda  and Fulgence N. Mpenda 

Department of Molecular Biology and Biotechnology, University of Dar es Salaam, Dar es salaam, P.O Box 35179, Tanzania



Received 16-10-2023

Revised 19-1-2024

Accepted 20-1-2024

Corresponding Author

Fulgence N. Mpenda

mpenda.fulgence@udsm.ac.tz

DOI <https://doi.org/10.47419/bjbabs.v4i03.244>

Pages: 136-150

Distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are properly cited

Copyright: © 2023 the Authors

ABSTRACT

Background and objectives: Microbial infections have made significant contributions to the global health burden, particularly in developing countries with lagging economies. Challenges arising from the evolution of antimicrobial-resistant (AMR) bacteria have led to the search for novel antimicrobial agents, especially those derived from medicinal plants. Therefore, this study aimed to assess the antimicrobial activity of ethyl acetate crude extracts and essential oils from *C. bonariensis* collected from the Mbeya region of Tanzania.

Methods: Initial experimental activities involved the preparation of crude extracts and essential oil, which were then, tested for antimicrobial activity against selected test organisms (*Staphylococcus aureus* (ATCC29213), *Bacillus subtilis* (ATCC6051), *Escherichia coli* (ATCC8736), *Salmonella typhi* (ATCC6539), and *Candida albicans* (DSM1665)) by using the disc diffusion method. On the other hand, the chemical composition profiling of ethyl acetate and essential oils was performed by gas chromatography-mass spectrometry (GC-MS).

Results: The crude extracts and essential oil of *C. bonariensis* leaves from Tanzania demonstrated to have antibacterial and antifungal activity. The minimum inhibitory concentration (MIC) of ethyl acetate leaf crude extract was 25 mg/mL > MIC > 12.5 mg/mL for tested microorganisms, with the exception of *C. albicans*, where the MIC was 50 mg/mL > MIC > 25 mg/mL. Similarly, for all tested microorganisms, the MIC of leaves essential oils was 1:1 > MIC > 1:2, except *S. typhi*, where the MIC was 2:1 > MIC > 1:1. On the other hand, 2,4-di-tert-butyl phenol in leaves crude extracts had the highest percentage composition of peak area (41%); whereas, in essential oil, 2,4-Di-tert-butyl phenol had the highest percentage composition (38.8%).

Conclusions: The present findings have demonstrated antimicrobial potential of ethyl acetate crude extract and essential oil of *C. bonariensis* leaves from Tanzania, and validate the wide use of the plant by local communities, particularly the local community of Mbeya region in Tanzania.

OPEN ACCESS

Keywords *Conyza bonariensis*, Antimicrobial Potential, Essential Oil

INTRODUCTION

Microbial infections have a significant contribution to the health burden worldwide, particularly in developing countries due to lagging economies.¹ The microbial infection burden is further exacerbated by the emergence of antimicrobial resistant microbes.^{1, 2} It has been estimated that about 700,000 people die each year as a result of antimicrobial resistance (AMR) around the world.³ In Sub-Saharan Africa (SSA), it was found that 73.4% of deaths caused by microbial infections were linked to antibiotic resistance of infectious agents.³ Specifically, in Tanzania, about 25% to 50% of the mortality caused by microbial infections was associated with antimicrobial resistance. Furthermore, antibiotic resistance threatens not just the country's mortality rate but also its economic stability, particularly in developing nations like Tanzania.⁴ Moreover, within clinical settings in Tanzania, the prevalence of multidrug-resistant bacteria ranges from 25% to 50%.^{1, 4, 5} These challenges caused by the evolution of antimicrobial resistant bacteria entail searching for novel antimicrobial agents, particularly from medicinal plants.

Medicinal plants are plants with therapeutic properties and have beneficial pharmacological effects on the human or animal body.⁶ Medicinal plants are a good source of bioactive substances such as flavonoids, tannins, curcumin, and resveratrol, which have antibacterial, anti-diabetic, anti-ulcer, and antioxidant properties.⁷ For example, *Plantago major* from Spain.⁸ Has been reported to have antimicrobial activities. *Aloe vera* which is from China, has been reported to have anti-diabetic effects⁹ and *Momordica charantia* which is from Turkey has antimicrobial and antioxidant activities.¹⁰ To assess the antibacterial activity of medicinal plants leaves crude extracts and essential oil have been found to be more appropriate.

Essential oils are aromatic, volatile liquids obtained from plant material through steam distillation and named after the plant from which they are derived.¹¹ Essential oils extracted from various medicinal plants have been shown to contain high levels of bioactive metabolites that have antimicrobial activity.¹² Essential oils have been widely used in the treatment of infectious diseases like skin diseases, respiratory diseases, and intestinal disorders. For example, essential oil from *Abies balsamea* is used for the treatment of skin infections.¹³ Also, essential oils from *Lavandula angustifolia* and *Mentha piperita* medicinal plants have been used in treatment of respiratory disorders.¹⁴

One empirical investigation was conducted in Tanzania to assess the antibacterial efficiency of *C. bonariensis*, and the results were promising.¹⁵ The inhibitory concentration of ethyl crude extracts of *C. bonariensis* leaves collected in Tanzania was around 1.6, 1.6, 6.3 and 1.7 mg/mL against *E. coli*, *S. aureus*, *S. typhi*, and *C. albicans*, respectively.¹⁵ However, essential oil from *C. bonariensis* was not tested for antibacterial activity. Also, it is

widely understood that the composition of secondary metabolites in medicinal plants can be influenced by variety of circumstances, including temperature, water, and soil properties.¹⁶ This suggests a wide-ranging surveillance study to assess the antimicrobial activity of crude extracts and essential oils of *C. bonariensis* collected from other places including Mbeya region of Tanzania. The aim of the present study was to determine the antimicrobial activity of crude extracts and essential oil from *C. bonariensis* leaves and to profile chemical composition of *C. bonariensis* leaves crude extracts and essential oil.

MATERIALS AND METHODS

Collection of plant materials

Plant samples were collected from Mbeya region of Tanzania. Purposive random sampling was used to select healthy leaves between November and December 2021 for extraction of leaves crude extracts and essential oil from *C. bonariensis*. For each subsequent sampling, a sample voucher was included for identification and was provided with voucher number FMM4139 after identification by a plant taxonomist at Botany Department, University of Dar Es Salaam.

Tested microorganisms

Five test microorganisms used were Gram-positive bacteria *Staphylococcus aureus* (ATCC29213), *Bacillus subtilis* (ATCC6051) Gram-negative bacteria *E. coli* (ATCC8736), *Salmonella typhi* (ATCC 6539) and the yeast *Candida albicans* (DSM1665), all obtained from the microbial strains library of the Department of Molecular Biology and Biotechnology (MBB), University of Dar Es Salaam. Bacteria and fungi were grown in nutrient agar and potato dextrose agar and incubated at 37 °C and 30°C for 24 hours and 48 hours respectively. Bacterial and fungal colonies were suspended in normal saline, which were then compared with 0.5 McFarland standards (prepared by adding 0.05 mL of 1.175% BaCl₂ to 9.95 mL of 1% H₂SO₄). Suspensions of tested organisms were kept in the test tubes at 4 °C for a maximum of overnight.¹⁷

Leaves crude extracts and essential oil preparation

Leaves crude extracts preparation

Leaves were kept in the shade for 21 days to dry. The dried leaves were ground into a fine powder using a grinder (Panasonic AC MX-AC300-H 550-Watt Mixer Grinder). The ground powder was soaked in 99.55% ethyl acetate and left for 72 hours so as to extract the bioactive constituents. The mixture was filtered using Whatman No.1 filter paper (pore size 11 µm) and the filtrate was then concentrated on the rotary evaporator (Model No RE-501) at 40°C and 100 mbar pressure. The obtained crude extract was stored at 4°C for further use.^{6, 16}

Essential oil preparation

Fresh leaves were cleaned in tap water and placed in a round-bottomed flask. An electric burner was used to set up a steam distillation setup, which was followed by heating the fresh leaves to 100 °C.^{16, 18} The essential oil was gathered in the collector after 45 minutes of boiling and followed by the addition of an equivalent volume of dichloromethane to the harvest. The lower layer was removed and concentrated using a rotary evaporator at 40 °C and 100 mbar pressure. The oil was collected and stored in dark glass tubes at 4 °C for later use.^{12, 16}

Screening for antimicrobial activity and Minimum inhibitory concentration

Screening for antimicrobial activities

Sterile 5 mm discs, which were prepared from Whatman No. 1 filter paper (pore size of 11 µm), were soaked with 20 µL of various concentrations (200 mg/mL, 100 mg/mL, 50 mg/mL, and 25 mg/mL) of *C. bonariensis* leaves crude extract. For *C. bonariensis* essential oils 20 µL of 1:1 with Dimethyl sulfoxide (DMSO) was used to soak discs. Discs were placed at 4°C in the refrigerator for 30 minutes to allow proper soaking before transferring to Petri dishes. Well-soaked discs were placed on Petri dishes, which were initially inoculated with test organism suspension by using sterile cotton swabs. Then, Petri dishes were incubated at 37°C and 30°C for 24 hours and 48 hours for bacteria and yeast, respectively. Ten percent of dimethyl sulfoxide (DMSO) was used as negative control; whereas, Chloramphenicol (0.25mg/ml) and Fulconazole (0.25mg/ml) were used as positive control for Bacteria and Fungi, respectively. The experiment was performed in triplicate then, zones of growth inhibition were measured after 24 hours of incubation for bacteria, and after 48 hours of incubation for fungi.¹⁹

Minimum inhibitory concentration

For test organisms that were inhibited by extracts and/or essential oils, the minimum inhibitory concentration (MIC) was determined. The MIC was carried out using the disc diffusion method, as described before.¹⁹ Each of the twofold serial dilutions of crude extracts and essential oil was tested for growth inhibition of test organisms. The discs were prepared, the test organisms were inoculated into Petri dishes, and the incubation period was as indicated in the section on screening for antimicrobial activities

Chemical analysis of Crude extract and essential oils

Chemical analysis was performed by Gas Chromatography-Mass Spectrometry recorded in GCMS-2010 Shimadzu instrument operating in Electron Ionization (EI) mode (MS) at 70 eV, and Flame Ionization Detector (FID) for GC. A Restek-5MS column (30m x 0.25mm x 0.25 µm) was used. The oven temperature program was 90°C to 280 °C and held at 90 °C for two minutes. The temperature was increased to 280 °C for 10 minutes (hold time) at a rate of 15 °C per minute. The injection temperature was 250°C with a split injection mode. The flow rate of carrier gas helium was 1.21 ml min⁻¹. The ion source temperature and interface temperature in MS were 230 °C and 300 °C, respectively. The identification of chemicals in the extracts was done by the scan method, which involves the use of Mass Spectral Library &

Search Software (NIST 11). The quantification of phytochemicals in the extracts was done using the Peak Integration method (area normalization), whereby the ion allowance was 20%, the target ion and other five quantization ions were used in the quantitative analysis. 10 μ L of the sample was dissolved in dichloromethane to make 1mL and injected in GC da Silva et al., 2021, The results were reported as percentage compositions derived from peak area of all scanned compounds in the extracts.^{20, 21}

DATA ANALYSIS

InR software, descriptive and inferential statistics were used to examine the data (R core team). For descriptive statistics, bar graphs were used to summarize the statistics; for inferential statistics, two-way analysis of variance (ANOVA) was employed to test whether there was a difference between the means of the zone of inhibition index of test among tested microorganisms. Following ANOVA, Tukey HSD was performed for pair wise comparison, with a significant difference set at $P < 0.05$.

RESULTS

Antimicrobial activity of *C. bonariensis* leaves crude extract and essential oil

The antimicrobial activity of essential oil and crude extract from *Conyza bonariensis* leaves was investigated using the disc diffusion method, as explained in subsection 2.2. Zones of inhibition against test microorganisms are depicted in Figure 1-2.

Activity indices, which were derived from zones of inhibition values, are presented in Table 1. The activity index for essential oil ranged from 0.22 to 0.47 against *S. typhi* and *B. subtilis*, respectively, and the crude extract activity index ranged from 0.35 to 0.4 against *S. aureus* and *B. subtilis*, respectively. Generally, crude extract had slightly elevated ($P > 0.05$) antimicrobial activity as compared to essential oils (Table 1).

Table 1. Antimicrobial activity of extracts from *Conyza bonariensis* leaves presented as activity index and standard error of mean

Test organism	Positive control	Essential oil	Leaves crude extract
Mean values of activity indices $\pm$ standard deviation			
<i>Escherichia coli</i>	1.00 $\pm$ 0.00 ^{b,1}	0.31 $\pm$ 0.04 ^{a,2}	0.38 $\pm$ 0.03 ^{b,2}
<i>Candida albicans</i>	1.00 $\pm$ 0.00 ^{b,1}	0.44 $\pm$ 0.04 ^{b,2}	0.37 $\pm$ 0.03 ^{b,2}
<i>Staphylococcus aureus</i>	1.00 $\pm$ 0.00 ^{b,1}	0.38 $\pm$ 0.06 ^{a,2}	0.35 $\pm$ 0.03 ^{b,2}
<i>Salmonella typhi</i>	1.00 $\pm$ 0.00 ^{b,1}	0.22 $\pm$ 0.31 ^{a,1}	0.38 $\pm$ 0.03 ^{b,1}
<i>Bacillus subtilis</i>	1.00 $\pm$ 0.00 ^{b,1}	0.47 $\pm$ 0.06 ^{b,2}	0.40 $\pm$ 0.01 ^{b,2}

Column that bear different superscript later are significant different ($P < 0.05$), and row that bear different superscript number are significant different ($P < 0.05$); positive control was chloramphenicol and fluconazole for bacteria and fungus, respectively. Concentrations




Figure 1 Representative images of zone of inhibition for crude extracts and essential oils from *C. bonariensis* leaves. A and A1 are front and back views of inhibition zone for *C. bonariensis* essential oil against *S. typhi*; B and B1 are front and back views of inhibition zone for *C. bonariensis* leaves crude extract against *E. coli*.




Figure 2 A bar graph representing zone of inhibition (mm) of ethyl acetate crude extracts and essential oils from *C. bonariensis* leaf against test organisms; error bars represent standard deviation, Positive control were chloramphenicol and fluconazole (0.25 mg/mL) for bacteria and fungi, respectively.

of positive control and leaves crude extract were 0.25 mg/mL and 100mg/mL, respectively. Essential oil concentration was 1:1 μ L dilution with 10% DMSO

The minimum inhibitory concentration of *C. bonariensis* leaves crude extract and essential oil

The MIC of crude extract and essential oil was evaluated for each of the tested microorganisms as demonstrated in Table 2. For leavecrude extract, the highest MIC was 25 mg/mL >MIC> 12.5 mg/mL, for four tested microorganisms, and the lowest MIC was 50 mg/mL >MIC> 25 mg/mL for *C. albicans*. Furthermore, for essential oil higher MIC was 1:1>MIC> 1:2 (means that essential oil was diluted with equal volume (20 μ L) of 10% DMSO) for all microorganisms and the lowest MIC was 2:1>MIC> 1:1 (means 20 μ L of essential oil was diluted with 10 μ L volume of 10% DMSO) for *S. typhi*.

Table 2. Minimum inhibitory concentration of crude extracts in mg/mL and essential oil v/v (μ L) extracted from *Conyza bonariensis*.

Extracts	Minimum inhibitory concentration of the crude extracts on selected microorganisms				
	<i>S. aureus</i>	<i>B. subtilis</i>	<i>E. coli</i>	<i>S. typhi</i>	<i>C. albicans</i>
Leaves crude	25>MIC>12.5	25>MIC>12.5	25>MIC>12.5	25>MIC>12.5	50>MIC>25
Essential oil	1:1>MIC> 1:2	1:1>MIC> 1:2	1:1>MIC> 1:2	2:1>MIC> 1:1	1:1>MIC> 1:2

For essential oils, the ratio 1:1 means that essential oil was diluted with equal volume (20 μ L) of 10% dimethyl sulfoxide (DMSO), and therefore the essential oil was diluted to half of original concentration. Whereas, for ratio 1:2, means that the essential oil was diluted to one third of original concentration.

Chemical composition of *C. bonariensis* leaves crude extract

Phytochemical analysis of the crude extract from *C. bonariensis* leaves detected different compounds based on peak area, retention time, and peak area and peak height. As illustrated in Figure 3 below, alkene and phenol predominated in crude extract from *C. bonariensis* leaves, especially 2,4-Di-tert-butylphenol (41.6%), Phytol (11.9%), 1-Nonadecene (8.7%), (-)-Spathulenol (6.6%), 1-Heptadecenol (6.4%), n-Tetracosanol-1 (6.4%), and Caryophyllene oxide (3.3%), as illustrated in figure 3 below. It is very interesting that most of the compounds detected in the present study have been demonstrated to have various health benefits as shown in Table 3.

Table 3. Biologically active chemical compounds of leaves crude extract from *Conyza bonariensis*

Name of compound	% Composition (Peak area)	R. Time	Chemical formula	Target Peak	P. Area	P. Height	Activity
2,4-Di-tert-butylphenol	41.6231	10.439	C ₁₄ H ₂₂ O	191.15	86349	49914	Antioxidant and antifungal ²²
(-)-Spathulenol	6.57777	11.244	C ₁₅ H ₂₄ O	119.15	17506	7888	Antibacterial Activity ²²
Caryophyllene oxide	3.26053	11.306	C ₁₅ H ₂₄ O	79.05	6422	3910	Antimicrobial ²³
1-Heptadecene	6.4443	13.537	C ₁₇ H ₃₄	57.1	16961	7728	Antifungal ²⁴
Phytol, acetate	15.01263	14.013	C ₂₂ H ₄₂ O ₂	68.1	33252	18003	Antioxidant ²⁵ Anti-inflammatory ²²
1-Nonadecene	8.72339	15.533	C ₁₉ H ₃₈	57.1	19951	10461	Antifungal ²⁶
Phytol	11.94556	16.66	C ₂₀ H ₄₀ O	71.1	31524	14325	Ant nociceptive ²⁷ and Antioxidant ²⁸ , anti-inflammatory and ²⁹ anti-allergic
n-Tetracosanol-1	6.41266	17.355	C ₂₄ H ₅₀ O	97.1	14111	7690	Antibacterial ³⁰




Figure 3: Chromatogram of *Conyza bonariensis* leaves crude extract

were 2,4-di-tertbutylphenol (38.8%), (-)-spathulenol (25.5%), 1-heptadecene (6.4%), n-tetracosanol-1 (6.4%), and caryophyllene oxide (8.7%) dominating, and 1-Nonadecene (6.4%), 1-Heptadecene (5.4%), Hexadecanoic acid, butyl ester (5.3%), Linoleic acid ethyl ester (5.6%) and Oleic Acid (4.2%) were relatively small . Also, most of the compounds detected from essential oil in the present study have been demonstrated to have various health benefits as demonstrated in Table4.

Table 4. Biologically active chemical compounds of essential oil extracted from *Conyza bonariensis* Leaves

Name of compound	% composition (P.area)	R. Time	Chemical formula	Target Peak	P. Area	P. Height	Activity
2,4-Di-tert-butylpheno	38.8342	10.44	C ₁₄ H ₂₂ O	191.15	58675	33411	Antioxidant and antifungal ^{31, 26}
(-)-Spathulenol	25.51055	11.243	C ₁₅ H ₂₄ O	119.15	49539	21948	Antibacterial Activity ³²
Caryophyllene oxide	8.73482	11.308	C ₁₅ H ₂₄ O	79.1	12611	7515	Antimicrobial activities ³³
1-Heptadecene	5.40129	13.537	C ₁₇ H ₃₄	57.1	11883	4647	Antifungal ³⁴
1-Nonadecene	6.40902	15.533	C ₁₉ H ₃₈	57.1	11729	5514	Antioxidant and antimicrobial ^{35, 36}
Hexadecanoic acid, butyl ester	5.33853	17.3	C ₁₇ H ₃₂ O ₂	56.1	9466	4593	Antioxidant ³⁷
Linoleic acid ethyl ester	5.64189	18.733	C ₂₀ H ₃₆ O	67.1	10399	4854	Anti-inflammatory ³⁸
Oleic Acid	4.12971	18.771	C ₁₈ H ₃₄ O ₂	55.1	8537	3553	Antioxidant activity ³⁸

DISCUSSION

Antimicrobial substances are plentiful in medicinal plants and several research have been undertaken to investigate the antimicrobial activity of plant parts such as seeds, flowers, stems, and roots. ^{39, 40} To assess antibacterial activity from medicinal plants, leaves crude extracts and essential oil have been found to be more appropriate because they possess a higher content of secondary metabolites than other form. ^{12, 18}

C. bonariensis is an annual or short-lived perennial plant. The members of genus *Conyza* are most difficult weeds to control. ⁴¹ *Conyza bonariensis* crude extracts and essential oils have been reported to have chemical constituents such as saponins, tannins, flavonoids, steroids, glycosides, diterpenoids and triterpenoids (Gutierrez et al., 2020), which have antioxidant, antimicrobial activities, anti-inflammatory activities. In Tanzania, *C. bonariensis* has traditionally been used to cure ailments like bacterial infections, fungal infections, and wound healing among various populations. ¹⁵ The Safwa tribe in Tanzania's Mbeya region traditionally uses *C. bonariensis* leaves to cure microbiological illnesses, notably fungal infections.

There is limited empirical evidence on *C. bonariensis* found in Tanzania, for example before the present findings, one study reported on *C. bonariensis* obtained from one geographical region of Tanzania (Arusha and Kilimanjaro regions, Northern part of Tanzania). ¹⁵ Furthermore, essential oil from *C. bonariensis* found in Tanzania was not yet evaluated, and it is widely understood that the composition of secondary metabolites in medicinal plants can be influenced by variety of circumstances, including temperature, water, and soil properties. ¹⁶ Suggesting crude extracts and essential oils of *C. bonariensis* from different geographic regions may have different antimicrobial performance.

In present study, crude extracts of *Conyza bonariensis* leaves had high antimicrobial activities against some of selected microorganisms. The highest MICs was found to be 25mg/mL>MIC>12.5mg/mL against all tested bacteria (*E.coli*, *S.aureus* and *S.typhi*) and lowest MIC was 50mg/mL>MIC>25mg/mL against *C. albicans*. On other hand essential oil extracted from *C. bonariensis* leaves also had antimicrobial activities against tested microorganisms, whereby the MIC ranged from 1:1>MIC> 1:2 for four tested microorganisms with the exception of *S. typhi* where the MIC range was 2:1>MIC> 1:1. Variability in antimicrobial activities to different microorganisms may be explained differences in biochemical composition of cell walls and other inherent factors of a particular microorganisms.^{42, 43} On the other hand, overall antimicrobial activity exhibited by crude extracts was slightly higher ($P>0.05$) than essential oil, these findings are challenging to compare since the two formulations (crude extracts and essential oil) have different units of measurement.

Previous studies on ethyl acetate crude extracts of *C. bonariensis* from Yemen and Tanzania reported higher antimicrobial activities compared to the findings of the present study.^{44, 15} For example, in the study that was conducted in Tanzania the MIC for *E.coli*, *S.aureus*, and *S.typhi* were 1.56 mg/mL, 6.25 mg/mL and 0.78 mg/mL, respectively. This observation may be due to a difference in geographical location from which *C. bonariensis* grows.³⁹ Also, the influence of slight genetic variability among tested microorganisms cannot be ignored.⁴⁵

In addition, in previous studies, leaves crude extracts and essential oils from medicinal plants were also reported to have antimicrobial activities.⁴² Study of antimicrobial activities of leaves crude extracts from various plants, such as *Cichorium endivia* leaves from Egypt using methanol.⁴⁶ *C. bonariensis* leaves from Arusha Tanzania using ethyl acetate.¹⁵ Crude extracts from five medicinal plants found in Saudi Arabia using ethanol extract.²¹ show high antimicrobial activities compared to the current study. In contrast, *Psidium guajava* leaves from the USA using (hexane, methanol, ethanol, and water) had shown low activity compared to the present findings.⁴⁷ The variation in antibacterial activity could be due to the crude extracts being from various plant species and the solvent utilized being different as well as variances in tested microorganisms and geographical conditions.^{48, 49, 50}

Furthermore, chemical analysis by gas chromatography mass spectrometry (GC-MS) showed that there were antimicrobial compounds in both leaves crude extracts and essential oil of *C. bonariensis*, such as 2,4-Di-tert-butylphenol, (-)-Spathulenol, Caryophyllene oxide, and 1-Heptadecene.^{51, 52} In leave crude extracts, Phytol, acetate, 1-Nonadecene, Phytol, n-Tetracosanol-1 were found. In contrast Hexadecanoic acid, butyl ester, linoleic acid ethyl ester, and Oleic Acid were observed in essential oil.^{52, 27, 31} The difference in antimicrobial activities of crude extracts and essential oil may be due to differences in chemical composition as shown in Table 3 and 4.

CONCLUSIONS

Based on in vitro antimicrobial activities and compounds screened by GC-MS, this study revealed that leaves extract and essential oils from *C. bonariensis* have antimicrobial properties. This study paves the way forward in the discovery of novel broad-spectrum drugs that can help to solve antimicrobial challenges, and chemical analysis provides directions for further in vivo research. Further studies should be done to isolate pure compounds from active fungi metabolites responsible for the shown activities.

DATA AVAILABILITY

All data presented in this article are available with the authors.

FUNDING

No external fund was received

DECLARATION OF COMPETING INTEREST

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

ACKNOWLEDGEMENT

Mr Maurice Mbago a plant taxonomist. The special appreciation to University of Dar es Salaam, the Department of Molecular Biology and Biotechnology to allow the experimental activities of the present study to be conducted.

REFERENCES

1. Blomberg B, Manji KP, Urassa WK, Tamim BS, Mwakagile DS, Jureen R. Antimicrobial resistance predicts death in Tanzanian children with bloodstream infections: a prospective cohort study. *BMC Infectious Diseases*. 2007;7(43). <https://doi.org/10.1186/1471-2334-7-43>.
2. Dadgostar P. Antimicrobial Resistance: Implications and Costs. *Infect Drug Resist*. 2019;12(3903):10–10. <https://doi.org/10.2147/IDR.S234610>.
3. Organization WH. Antimicrobial resistance; 2023. Available from: <https://www.who.int/news-room/fact-sheets/detail/antimicrobial-resistance>.
4. Manyahi J, Matee MI, Majigo M, Moyo S, Mshana SE, Lyamuya EF. Predominance of multi-drug resistant bacterial pathogens causing surgical site infections in Muhimbili national hospital. *BMC Res Notes*. 2014;7(500). <https://doi.org/10.1186/1756-0500-7-500>.

5. Moremi HCN, Stephen E, Mshana. Antimicrobial resistance pattern: a report of microbiological cultures at a tertiary hospital in Tanzania. 2016;<https://doi.org/10.1186/s12879-016-2082-1>.
6. Bisi-Johnson MA, Obi CL, Samuel BB, Eloff JN, Okoh AI. Antibacterial activity of crude extracts of some South African medicinal plants against multidrug resistant etiological agents of diarrhoea. BMC Complement Altern Med. 2017;17(321). <https://doi.org/10.1186/s12906-017-1802-4>.
7. Kianfé BY, Kühlborn J, Tchuenguem RT, Tcheegnitegni BT, Ponou BK, Groß J. Antimicrobial secondary metabolites from the medicinal plant *Crinum glaucum* A. Chev.(Amaryllidaceae). S Afr J Bot. 2020;(133):161–167. <https://doi.org/10.1016/j.sajb.2020.07.026>.
8. Pensantes-Sangay SJ, Calla-Poma RD, Requena-Mendizabal MF, Alvino-Vales MI, Millones-Gómez PA. Chemical Composition and Antibacterial Effect of Plantago Major Extract on Periodontal Pathogens 20. Pesqui Bras Odontopediatria Clín Integr. 2020;Available from: <https://doi.org/10.1590/pboci.2020.100>. <https://doi.org/10.1590/pboci.2020.100>.
9. Urmeela T, Mahomoodally F. Aloe vera extracts: From traditional uses to modern medicine. 2015;p. 211–226. Available from: https://www.researchgate.net/publication/297766937_Aloe_vera_extracts_From_traditional_uses_to_modern_medicine.
10. Masithoh DA, Kusdarwati R, Handijatno D. Antibacterial activity of bitter gourd (*Momordica charantia* L.) leaf extract against *Aeromonas hydrophila*. IOP Conf Ser: Earth Environ Sci. 2019;236:12096–12096. <https://doi.org/10.1088/1755-1315/236/1/012096>.
11. Yang C, U H, Feng DH, Y. Antibacterial activity and mode of action of the *Artemisia capillaris* essential oil and its constituents against respiratory tract infection-causing pathogens. Molecular Medicine Reports. 2015;11. <https://doi.org/10.3892/mmr.2014.3103>.
12. Hawrelak J, Cattley T, Myers S. Essential Oils in the Treatment of Intestinal Dysbiosis: A Preliminary in vitro Study. Alternative Medicine Review : A Journal of Clinical Therapeutic. 2009;14(380):4–4. Available from: <https://pubmed.ncbi.nlm.nih.gov/20030464/>.
13. Orchard A, Vuuren SV. Commercial Essential Oils as Potential Antimicrobials to Treat Skin Diseases. Evid Based Complement. Alternat Med. 2017;p. 2017–2017.
14. Horváth G, S KAC. Essential oils in the treatment of respiratory tract diseases highlighting their role in bacterial infections and their anti-inflammatory action: Flavour Fragr J. 2015;30(5). <https://doi.org/10.1002/ffj.3252>.
15. Ghwanga AAK, Chacha M. Vitro Antimicrobial Activity of *Conyza bonariensis* and *Tribulus terrestris* Growing in Tanzania Open. 2019;5(1:4):7–7. [10.21767/2472-0151.100042](https://doi.org/10.21767/2472-0151.100042).
16. Abers M, Schroeder S, Goelz L, Sulser A, Rose ST, Puchalski T, et al. Antimicrobial activity of the volatile substances from essential oils BMC Complementary Medicine and Therapies. 2021;21:124–124. <https://doi.org/10.1186/s12906-021-03285-3>.
17. Aryal S. McFarland Standards- Principle, Preparation, Uses, Limitations Microbe

- Notes. 2021; Available from: <https://microbenotes.com/mcfarland-standards/>.
18. Reddy DN, Akhtar MS, Swamy MK, Sinniah UR. Essential Oils Extracted from Medicinal Plants and Their Applications. In: editors Natural Bio-active Compounds. 2019;1:237–83. <http://dx.doi.org/10.1007/978-981-13-7154-7>.
 19. Khaki P, Sharma A, Bhalla P. Comparison of Two Disc Diffusion Methods with Minimum Inhibitory Concentration for Antimicrobial Susceptibility Testing of *Neisseria Gonorrhoeae* Isolates. *Ann Med Health Sci Res*. 2014;4(3):453–459. <https://doi.org/10.4103/2141-9248.133477>.
 20. Silva ACAD, Matias EFF, Rocha JE, Araújo AC, Freitas TS, Campina FF. Gas chromatography coupled to mass spectrometry (GC-MS) characterization and evaluation of antibacterial bioactivities of the essential oils from *Piper arboreum* Aubl *Piper aduncum* L. e *Piper gaudichaudianum* Kunth. 2020;76(1-2)(35–42.):35–42. <https://doi.org/10.1515/znc-2020-0045>.
 21. Viteri R, Peñarreta JP. Antioxidant activity and GC-MS profile of *Conyza bonariensis* L. leaves extract and fractions. vol. 73 of 3; 2020. <https://doi.org/10.15446/rfnam.v73n3.81452>.
 22. Chy AMNU, Kamal M, Barlow ATM, Faruque JW, Yang MO, X. Evaluation of antinociceptive and anti-inflammatory activities of the methanol extract of *Holigarna caustica* (Dennst.) Oken leaves. *J Ethnopharmacol*. 2019;236:401–412. Available from: https://www.researchgate.net/publication/277230263_Antimicrobial_Effect_of_15_Medicinal_Plant_Species_and_their_Dependency_on_Climatic_Conditions_of_Growth_in_Different_Geographical_and_Ecological_Areas_of_Fars_Province.
 23. Klug MJ, Markovetz AJ. Degradation of hydrocarbons by members of the genus *Candida* 3. Oxidative intermediates from 1-hexadecene and 1-heptadecene by *Candida lipolytica*. *J Bacteriol*. 1968;96(4):1115–1138. <https://doi.org/10.1128/jb.96.4.1115-1123.1968>.
 24. Santos CC, De MP, Salvadori MS, Mota VG, Costa LM, De Almeida AAC, et al. Antinociceptive and Antioxidant Activities of Phytol In Vivo and In Vitro Models. *Neurosci J*. 2013;949452. <https://doi.org/10.1155/2013/949452>.
 25. Khan IH, Javaid A. DNA cleavage of the fungal pathogen and production of antifungal compounds are the possible mechanisms of action of biocontrol agent *Penicillium italicum* against *Macrophomina phaseolina*. *Mycologia*. 2022;114(1):24–34. <https://doi.org/10.1080/00275514.2021.1990627>.
 26. Gossweiler AG, Martinez-Mier EA. Chapter 6: Vitamins and Oral Health. *Monogr Oral Sci*. 2020;28:59–67. <https://doi.org/10.1159/000455372>.
 27. Islam MT, De Alencar MVOB, Machado KDC, Machado KDC, Melo-Cavalcante AADC, De Sousa DP. Phytol in a pharma-medico- stance. *Chem Biol Interact*. 2015;p. 60–73. <https://doi.org/10.1016/j.cbi.2015.07.010>.
 28. Kohli V, Sharma D, Sandur SK, Suryavanshi S, Sainis KB. Immune responses to novel allergens and modulation of inflammation by vitamin K3 analogue: a ROS dependent mechanism. *Int Immunopharmacol*. 2011;11(2):233–276. <https://doi.org/10.1016/j.intimp.2010.11.027>.
 29. Mandal MD, Mandal S. Honey: its medicinal property and antibacterial activity.

- Asian Pac J Trop Biomed. 2011;1(2):60016–60022. [https://doi.org/10.1016/S2221-1691\(11\)60016-6](https://doi.org/10.1016/S2221-1691(11)60016-6).
30. Lv JS, Zhang LL, Zhou CXZ, F J. Chemical composition, antioxidant and antimicrobial activity of the extracts of the flowers of the Chinese plant *Chimonanthus praecox*. Nat Prod Res. 2012;26(14):1363–1370. <https://doi.org/10.1080/14786419.2011.602828>.
31. Varsha KK, Devendra L, Shilpa G, Priya S, Pandey, Nampoothiri KM. 4- Di-tert-butyl phenol as the antifungal, antioxidant bioactive purified from a newly isolated *Lactococcus* sp. Int J Food Microbiol. 2015;15(211):44–50. <https://doi.org/10.1016/j.ijfoodmicro.2015.06.025>.
32. Boeck P, Leal PC, Yunes RA, Filho VC, López S, Sortino M. Antifungal activity and studies on mode of action of novel xanthoxylone-derived chalcones. Arch Pharm (Weinheim). 2005;338(2-3):87–95. <https://doi.org/10.1002/ardp.200400929>.
33. Veres K, Csupor B, Lázár A, Hohmann J. Antifungal Activity and Composition of Essential Oils of *Conyza canadensis* Herbs and Roots. Sci World J. 2012;48(46):646–646. <https://doi.org/10.1100/2012/489646>.
34. Skanda S, Vijayakumar BS. Antioxidant and Anti-inflammatory Metabolites of a Soil-Derived Fungus *Aspergillus arcoviridensis* SSSIHL-01. Curr Microbiol. 2021;78(4):1317–1340. <https://doi.org/10.1007/s00284-021-02401-3>.
35. Tonisi S, Okaiyeto K, Hoppe H, Mabinya LV, Nwodo UU, Okoh AI. Chemical constituents, antioxidant and cytotoxicity properties of *Leonotis leonurus* used in the folklore management of neurological disorders in the Eastern Cape. South Africa 3 Biotech. 2020;10(3):141–141. <https://doi.org/10.1007/s13205-020-2126-5>.
36. Eleazu CO. Characterization of the natural products in cocoyam (*Colocasia esculenta*) using GC-MS. Pharm Biol. 2016;54(12):2880–2885. <https://doi.org/10.1080/13880209.2016.1190383>.
37. Kolar MJ, Konduri S, Chang T, Wang H, Mcnerlin C, Ohlsson L. Linoleic acid esters of hydroxy linoleic acids are anti-inflammatory lipids found in plants and mammals. J Biol Chem. 2019;294(27):10698–707. <https://doi.org/10.1074/jbc.ra118.006956>.
38. Perona JS, Arcemis C, Ruiz-Gutierrez V, Catalá A. Effect of dietary high-oleic-acid oils that are rich in antioxidants on microsomal lipid peroxidation in rats. J Agric Food. 2005;53(3):730–735. <https://doi.org/10.1021/jf048374p>.
39. Abdollahi A, Fasihi-Ramandi M, Kouhpayeh S, Najafipour S, Meshkibaf M, Naghdi M. Antimicrobial Effect of 15 Medicinal Plant Species and their Dependency on Climatic Conditions of Growth in Different Geographical and Ecological Areas of Fars Province. ZJRMS. 2012;14:31–90. Available from: https://www.researchgate.net/publication/277230263_Antimicrobial_Effect_of_15_Medicinal_Plant_Species_and_their_Dependency_on_Climatic_Conditions_of_Growth_in_Different_Geographical_and_Ecological_Areas_of_Fars_Province.
40. Mpinda C, Meyer D, Lyantagaye S. In vitro antimicrobial activity of extracts of some plant species used in the management of HIV/AIDS in Tanzania. Tanzania Journal of Science. 2018;44(1):179–90. Available from: <https://www.ajol.info/index.php/tjs/article/view/171562>.

41. Wang A, Wu H, Zhu X, Lin J. Species Identification of *Conyza bonariensis* Assisted by Chloroplast Genome Sequencing. *Front Genet.* 2018;9(374). <https://doi.org/10.3389/fgene.2018.00374>.
42. Berman J, K DJ. Drug resistance and tolerance in fungi 2020. *Nat Rev Microbiol.* 2020;18(6):319–331. <https://doi.org/10.1038/s41579-019-0322-2>.
43. Fuerst JA. Microorganisms-A Journal and a Unifying Concept for the Science of Microbiology. *Microorganisms.* 2014;2(4):140–146. <https://doi.org/10.3390/microorganisms2040140>.
44. Thabit RAS, Cheng XR, Tang X, Sun J, Shi YH, Le GW. Antioxidant and antibacterial activities of extracts from *Conyza bonariensis* growing in Yemen. *Pak J Pharm Sci.* 2015;28(1):129–163. 25553691. Available from: <https://pubmed.ncbi.nlm.nih.gov/25553691/>.
45. Gonelimali FD, Lin J, Miao W, Xuan J, Charles F, Chen M. Antimicrobial Properties and Mechanism of Action of Some Plant Extracts Against Food Pathogens and Spoilage Microorganisms. *Frontiers in Microbiology.* 2018;9(1639). <https://doi.org/10.3389/fmicb.2018.01639>.
46. Girma Y, Jiru TM. Evaluation of Antimicrobial Activity of *Conyza bonariensis* Leaf Extracts against Clinically Isolated Fungi Causing Superficial Infection. *Journal of Chemistry.* 2021;<https://doi.org/10.1155/2021/6367449>.
47. Bipul B, Kimberly R, Fredrick MCL, Dwayne D, Anand Y. Antimicrobial Activities of Leaf Extracts of Guava (*Psidium guajava* L.) on Two Gram-Negative and Gram-Positive Bacteria. *Int J Microbiol.* 2013;p. 746165–746165. <https://doi.org/10.1155/2013/746165>.
48. Karleigh B, Renee B, Cynthia D, Sean O, Andrew N, W R. Evaluation of different solvents to extract antibacterial compounds from jalapeño peppers. *PMC.* 2017;5(3). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5448369/>. <https://doi.org/10.1002/fsn3.423>.
49. Wang H, Kolar MJ, Chang T, Rizo J, Konduri S, Mcnerlin C, et al. Stereochemistry of Linoleic Acid Esters of Hydroxy Linoleic Acids. *NIH.* 2019;21(19):8080–8084. <https://doi.org/10.1021/acs.orglett.9b03054>.
50. Jarmo K, Holopainen VV, Rajendra P, Ghimire. Climate Change Effects on Secondary Compounds of Forest Trees in the Northern Hemisphere | *Plant Science. Frontiers* . 2018;9(1445). Available from: <https://www.frontiersin.org/articles/10.3389/fpls.2018.01445/full>. <https://doi.org/10.3389/fpls.2018.01445>.
51. Viszwapriya D, Prithika U, Deebika S, Balamurugan K, Pandian SK. In vitro and in vivo antibiofilm potential of 2,4-Di-tert-butylphenol from seaweed surface associated bacterium *Bacillus subtilis* against group A streptococcus. *Microbiol Res.* 2016;191:19–31. <https://doi.org/10.1016/j.micres.2016.05.010>.
52. YangD, Michel L, Chaumont JP, Millet-Clerc J. Use of caryophyllene oxide as an antifungal agent in an in vitro experimental model of onychomycosis. *Mycopathologia.* 1999;148(2):79–82. <https://doi.org/10.1023/a:1007178924408>.