



Biostratigraphy of Eocene Sediments from Naopurdan Group, Chwarta Area, Kurdistan Region, NE Iraq: Paleogeographic Implication

Imad M. Ghafor ¹ , Hemn F. Muhammad ²

^{1,2}Department of Geology, College of Science, University of Sulaimani, Sulaymaniyah, Iraq.

Article information

Received: 31- Aug -2022

Accepted: 23- Oct -2022

Available online: 31- Dec- 2022

Keywords:

Microfossils

Biostratigraphy

Eocene

Naopurdan Group

NE Iraq

Correspondence:

Name: Imad M. Ghafor

imad.gafor@univsul.edu.iq

ABSTRACT

The main purpose of the present work is to describe the microfossils and biostratigraphy of marine Eocene deposits, which are described and studied in detail from carbonates of the Naopurdan Group, Chwarta area, Kurdistan Region, Northeastern Iraq. In order to define the main index fossils and to establish the biostratigraphy and age constraints of the exposed unit in the study area. A total of thirty-eight samples, spaced ca 0.5 to 1 m apart, have been collected from the field, approximately 60 thin sections are examined and the microfossils have been identified using the polarized microscopes. Three main biozones are recorded in the studied section, these zones are: (*Nummulites globulus* - *Nummulites planulatus* zone, at the bottom of the section and assigned to the Early Eocene (Ypressian) age, *Nummulites mamillatus* - *Nummulites alsharhani* zone, collected from the middle part of the section is assigned to the Middle Eocene (Lutetian) age, and *Nummulites fabianii* - *Assilina exponens* zone collected from the upper part of the section is assigned to the Early Bartonian age. So, the studied section of the Naopurdan Group is suggested to be Ypressian - Early Bartonian age. These three biozones are correlated with the other zones inside and outside Iraq.

DOI: [10.33899/earth.2022.135618.1031](https://doi.org/10.33899/earth.2022.135618.1031), ©Authors, 2022, College of Science, University of Mosul.

This is an open-access article under the CC BY 4.0 license (<http://creativecommons.org/licenses/by/4.0/>).

الطباقية الحياتية لترسبات الايوسين ضمن مجموعة ناوبردان في منطقة جوارتا، اقليم كردستان، شمال شرق العراق: تضمين الجغرافيا القديمة

عماد محمود غفور¹ , هيمن فريق محمد²

^{1,2} قسم علوم الأرض، كلية العلوم، جامعة السليمانية، السليمانية، العراق

المخلص	معلومات الارشفة
الهدف الرئيسي من العمل الحالي هو وصف المتحجرات الدقيقة والطباقية الحياتية لرواسب الايوسين، التي تم وصفها ودراستها بالتفصيل من كربونات مجموعة ناوبردان البحرية، في منطقة جوارتا، اقليم كردستان، شمال شرق العراق، من أجل تحديد المتحجرات الدالة والرئيسية لاعطاء الطباقية الحياتية وتحديد العمر لمنطقة الدراسة. تم جمع ثمانية وثلاثين عينة من الحقل بمسافة بين 0.5 إلى 1 متر، وفحص ما يقرب من 60 من الشرائح الرقيقة وتم التعرف على المتحجرات الدقيقة باستخدام المجاهر المستقطبة. بالاعتماد على المتحجرات الدقيقة وخاصة المنخربات الكبيرة الحجم، تم تسجيل ثلاث أنطقة حياتية رئيسية في منطقة الدراسة، وهذه الانطقة هي:	تاريخ الاستلام: 31-أغسطس-2022
1- <i>Nummulites globulus</i> - <i>Nummulites planulatus</i> zone	تاريخ القبول: 23-أكتوبر-2022
نطاق (Ypressian) في الجزء السفلي من المقطع ويعود للعصر الأيوسيني المبكر	تاريخ النشر الالكتروني: 31-ديسمبر-2022
2- <i>Nummulites mamillatus</i> – <i>Nummulites alsharhani</i> zone.	الكلمات المفتاحية :
(Lutetian) تم جمعها من الجزء الأوسط للمقطع ويعود لعصر الأيوسين الأوسط	المتحجرات القيقة
3- <i>Nummulites fabianii</i> - <i>Assilina exponens</i> zone.	الطباقية الحياتية
و نطاق (Early Bartonian) التي تم جمعها من الجزء الأعلى للمقطع ويعود للبارتوني المبكر	الايوسين
لذلك يُقترح بأن يكون منطقة الدراسة لمجموعة ناوبردان يعود الى عمر بيريشين الى بارتونين المبكر (Ypressian - Early Bartonian age)	مجموعة ناو بردان
تم مضاهاة هذه الأنطقة مع الأنطقة الحياتية الاخرى داخل وخارج العراق.	شمال شرق العراق
	المراسلة:
	الاسم: عماد محمود غفور
	imad.gafor@univsul.edu.iq

DOI: [10.3389/earth.2022.135618.1031](https://doi.org/10.3389/earth.2022.135618.1031), ©Authors, 2022, College of Science, University of Mosul.

This is an open-access article under the CC BY 4.0 license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

One selected section at Chwarta area (Lat. 35°43'28.31" N and Long. 45°35'27.98" E) is studied for foraminiferal analysis. It is located at the northwestern corner of Chwarta town, which lies between Lat. 35°43'28.31" N and Long. 45°35'27.98" E in the Imbricate/Thrust Zone (Fig. 1). The area consists of dark, white and milky grey fossiliferous limestone with 38m thick and its lower boundary is with Walash Group and its upper boundary is with Red Bed Series. The description of the lithological constituents and fieldwork investigation is inferred as shown in figure (2). Bolton (1958d) was the first who described the Naopurdan Shaly Group at Naopurdan village in Rawanduz River Valley (type locality) and he determined the age of this group as Paleocene–Oligocene or possible Miocene. Naopurdan Group is subdivided into two subgroups (Naopurdan-Type Subgroup and Sidekan-Type Subgroup) (Buday, 1980). Eocene –Oligocene age of the group was recorded by Polnikov and Nikolajev (1962) in Jassim and Goff (2006). Ghafor (1988), studied the Early Eocene strata from Sinjar area, NW Iraq. According to Jassim and Goff (2006), the Naopurdan Group was deposited in the Paleogene. The Eocene rocks were described from Northeastern Iraq by Ghafor and Karim (1999). Ghafor (2000) has described the Early to Late Danian rocks in Shaqlawa area, NE Iraq, using spores and pollen. The fossils of Eocene rocks in North and Northeastern Iraq were described by Sharbazheri, et al. (2009) and Sharbazheri, et al. (2011),

who gave Early Eocene age to the Eocene rocks by using planktic and benthic foraminiferas. Al Nuaimy, et al. (2020) separated the Eocene sediments from Northeastern Iraq, based on the planktic and benthic foraminifera. Ghafor and Qadir (2009) have described three groups of larger foraminifera from the Red Bed Series, which more likely belong to Walsh–Naopurdan group. early Eocene (Ypressian) age was recorded for the group by Al Qayim, et al. (2014). Mirza, et al. (2016) studied the Walsh–Naopurdan group from Northeastern Iraq and give Middle Eocene in age. Ali, et al. (2017) recorded Eocene to Oligocene age of the Walsh and Naopurdan group in Northeastern Iraq. Al Fattah, et al. (2018), Al Fattah, et al. (2020a) and Al Fattah, et al. (2020b), studied the Early Eocene rocks in Sinjar, Shaqlawa and Dhock areas. Kharajiani, (2018) assigned the middle Eocene age for the Betwata section, NE Iraq, based on the Calcareous nannofossils. Daoud (2019), studied the Nummulitic limestone of Naopurdan Group, and he recorded Late Paleocene-Early Eocene age. Finally, Ahmad, et al. (2022) studied the condensed section within the Naopurdan Group, and they confirm the Lutetian age of succession. This work aims to reformulate the study of microfossils and the biostratigraphy of the study section.




Fig. 1. Location, tectonic and geological map of the study area, northeast Iraq. A. tectonic map. (Al-Kadhimi, et al. 1996). B. Geologic and location map.




Fig.2. Lithostratigraphic column of Late Paleocene-Early-Middle Eocene rock units from the studied section.

Materials and Methods

The idea of the present work is reconstructed after four reconnaissance trips field works and mapping during one year for the outcrops of Chuwarta area, NE Iraq (Fig. 3). Thirty-eight samples are collected from the Nummulitic limestone layers, and then microfossils are identified from the seventy-thin sections by the polarized microscope. The studies by Loeblich and Tappan (1988), and Bolli (1966) are adopted for planktic foraminiferal identification and the studies of Seyrafian and Mojikhalifeh 2005; Sirel, et al. 2013; Gedik 2014; Serra-Kiel, et al. 2016; Roozpeykar and Moghaddam 2016; Ferrandez-Canadell) and Bover-Arnal (2017) are adopted for identification of benthic foraminifera.



Fig. 3. General view of studied section (Chwarta), view toward the northwest.

Geological setting

According to Buday (1980), and Buday and Jassim (1987), the studied section is within the Imbricate / Thrust Zone. The area represents partial area where the continental sides of the Iranian and Arabian plates collide during the Eocene (Numan, 1997; Surdashi, 1997) or Upper Cretaceous (Karim, 2004). Heron and Lees (1943) clarified that the name Naopurdan is a part of the Lower Division of the Nappe Zone and is underlain by Cretaceous rocks and Red Bed Series. Al-Mehaidi (1975), Karim (2004), Numan (1997), and Surdashy (1997) show that the area is of Upper Cretaceous and Eocene age and is a part of the continental sides of the Iranian and Arabian plates.

Results

Biostratigraphy

The studied sequence yields enrichment in large benthic foraminiferal assemblages (59 species belonging to 14 genera of large benthic foraminifera and 2 species of planktic Foraminifera belonging to 2 genera) (Plates 1, 2, 3 and 4). Benthic foraminifera has been used for biostratigraphic analysis for the studied section. The stratigraphic distribution of the Early - Middle Eocene larger foraminifera recorded from the Naopurdan Group permits the recognition of three biozones (Fig. 4) and by correlation with different studies; the recognized biozones are described below, starting from older to younger.




Fig. 4. Biostratigraphy of the Naopurdan Group at the Chuwarta area of the studied section.

1- *Nummulites globulus* - *Nummulites planulatus* Assemblages Zone (Ypresian). SBZ 7-12

Definition: It is characterized by the association of the taxa *Nummulites mamillatus*, *Nummulites globulus* and *Nummulites planulatus*).

Boundaries: The lower boundary of this zone is marked by the First Appearance Datum (FAD) of *Nummulites globulus* and *Nummulites planulatus*., whereas its upper boundary is marked by the Last Appearance Datum (LAD) of *Nummulites mamillatus*.

Thickness: 9m represented by the samples (30–38).

Remarks: The most diagnostic species include: *Nummulites globulus*, *Nummulites planulatus*, *Nummulites exilis*, *Nummulites djokdjokartae*, *Nummulites mamillinus*, *Nummulites fraasi*, *Nummulites solitaries*, *Nummulites minutus*, *Nummulites* sp., *Alveolina oblonga*, *Alveolina laxa*, *Alveolina globula*, *Alveolina regularis*, *Alveolina globosa*, *Alveolina pastillicata*, *Alveolina pissiformis*, *Alveolina munieri*, *Alveolina* sp., *Assilina spira*, *Assilina exponens*, *Assilina dandotica*, *Assilina sublamina*, *Assilina granulosa*, *Assilina laminosa*, *Assilina elliptica*, *Assilina subovata*, *Assilina frumentiformis*, *Assilina solida*, *Assilina fornasinii*, *Assilina laxa*, *Assilina* sp., *Sphaerogypsina* sp., *Quinqueloculina* sp., coral, pelecypods, gastropods, echinodermata, bryozoan and algae.

Correlation and Age Determination: This zone is equivalent to *mossoluensis-carbarica-trempina-oblonga-dainelli-violae* zone of Hottinger (1960) of Ypressian age, and correlated to the SBZ 7-12 Zone of Serra Kiel, et al.(1998), to the BFZK1A - BFZK2- BFZK 3A- BFZK 3B zone of Ahmad (2010) (Fig. 5), who considered it of Ypressian age.

Age: Early Eocene (Ypressian).

2- *Nummulites mamillinus* - *Nummulites alsharhani* Interval zone (Lutetian) SBZ 13-16

Definition: This zone represents the stratigraphic range of the index species *Nummulites mamillinus* which precedes the appearance of *Nummulites alsharhani*.

Boundaries: The lower boundary of this zone is marked by the Last Appearance Datum of *Nummulites mamillatus*, whereas its upper boundary is marked by the Last Appearance Datum (LAD) of *Nummulites alsharhani*.

Thickness: 19m represented by the samples (11–29).

Remarks: This zone is recorded from the middle part of the studied section, and the most diagnostic species include: *Nummulites alsharhani*, *Nummulites globulus*, *Nummulites perforates*, *Nummulites planulatus*, *Nummulites djokdjokartae*, *Nummulites discorbinus*, *Assilina granulosa*, *Assilina laminosa*, *Assilina spira*, *Assilina exponens*, *Assilina dandotica*, *Alveolina oblonga*, *Alveolina elliptica*, *Discocyclina archiaci*, *Discocyclina varians*, *Discocyclina ranikotensis*, *Discocyclina sella*, *Discocyclina dispansa*, *Discocyclina* sp., *Lepidocyclina*, *Sphaerogypsina* sp., *miliolina*, *Asterocyclina matanzensis*, *Quinqueloculina* sp., *Morozovella formosa*, *Subbotina linaperta*, coral, bryozoans and calcareous algae.

Correlation and Age Determination: This zone is apparently equivalent to *stips-munieri-proerecta* zone which is described by (Hottinger 1960), of Lutetian age, and correlated to the SBZ13-to lower part of SBZ17 zone of Serra-Kiel, et al. (1998), to the BFZK 4- the BFZK5- the BFZK 6 zone of Ahmad (2010). Locally in Iraq, this zone is correlated with *Hantkenina alabamensis-Acaranina bulbrooki* zone of Karim and Al-Kubaysi (2015), to the *Acaranina bulbrooki-Catapsydras dissimilis-Morozovella lehneri* zone of Ghafor and Al-Qayim (2021) and finally it is correlated to the *Nummulites gizehensis- Nummulites mouculatus* zone of Al-Qayim and Ghafor (2022). (Fig. 5), who considered it of Lutetian age.

Age: Middle Eocene (Lutetian)

System	Series	Stage	Hottinger 1960	Serra Kiel et al., 1998	Ahmad, 2010	Al-Kubaysi, 2014	Karim & Al-Kubaysi, 2015	Ghafor & Al-Qayim, 2021	Al-Qayim & Ghafor, 2022	This Study	
Paleogene	Eocene	Bartonian	<i>elongata</i>	SBZ 18 SBZ 17	Not Recognized Regressive Facies	Miliolids- Penerpolid Assemblage Zone		<i>Glg. semi-involuta</i> <i>H. alabamensis</i>	<i>Assilina spira- Lokharatia huntii</i>	<i>Nummulites fobianii- Assilina exponens</i> Assemblage Zone	
		Lutetian		SBZ 16	BFZK 6	<i>Nummulites gizehensis</i> Zone	<i>Acaranina bulbrooki</i>	<i>Nummulites boyharensis</i>	<i>M. lehneri</i>	<i>Nummulites gizehensis- Nummulites mocolatus</i>	<i>Nummulites mamillatus- Nummulites alsharhani</i> Interval Zone
			<i>proerecta</i>	SBZ 15	BFZK 5 BFZK 4			<i>Alveolina lockhartia</i>	<i>A. bulbrooki C. dissimilis</i>		
			<i>munierie</i>	SBZ 14			No Larger Benthic Foraminifera				
			<i>stipes</i>	SBZ 13							
			Ypresian	<i>violae</i>	SBZ 12		BFZK 3B				
		<i>dainelli</i>		SBZ 11							
		<i>oblonga</i>		SBZ 10							
		<i>trepmina</i>		SBZ 9	BFZK 3A						
		<i>carbarica</i>		SBZ 8	BFZK 2						
		<i>mossoulensis</i>		SBZ 7	BFZK 1A						

Legend:	SBZ Shallow Benthic Zone	<i>Glg.</i> <i>Globigerinatheca</i>	<i>H.</i> <i>Hantkenina</i>	<i>M.</i> <i>Morozovella</i>	<i>A.</i> <i>Acaranina</i>	<i>C.</i> <i>Catapsydrax</i>		Not Studied
---------	-----------------------------	--	--------------------------------	---------------------------------	-------------------------------	---------------------------------	---	-------------

Fig. 5. Comparisons of zonal schemes of the studied section with those proposed by some authors.

3- *Nummulites fabianii* - *Assilina exponens* Assemblages Zone (Early Bartonian) SBZ 17-18

Definition: This zone represents the stratigraphic range and association of the taxa *Nummulites fabianii*, *Nummulites catari* and *Assilina exponens*.

Boundaries: it is represented by the First Appearance Datum (FAD) of *Nummulites fabianii* and *Assilina exponens*), and the Last Appearance Datum (LAD) of *Alveolina ellipsoidalis*. Thickness: 10m represented by the samples (1–10).

Remarks: The zone is recorded from the upper part of the section, and the most diagnostic species include: *Nummulites ptukhiani*, *Nummulites fabianii*, *Nummulites alsharhani*, *Assilina exponens*, *Alveolina ellipsoidalis*, *Discocyclus* sp., *Lepidocyclus* sp., *Heterostegina* sp., *Sphaerogypsina* sp., *Quinqueloculina* sp., *Pyrgo* sp., *Miliolina*, coral, bryozoans, *Lithophyllum* sp., and gastropods shell.

Correlation and Age Determination: This zone is apparently equivalent to elongate zone which is described by Hottinger (1960), of early Bartonian age, and correlated to the SBZ 17-18 zone of Serra-Kiel, et al. (1998), Locally in Iraq, this zone is correlated with Miliolids-peneroplid zone of Al-Kubaysi (2014), to the *Globigerina semiinvoluta*- *Hantkenina alabamensis* of Ghafor and Al-Qayim (2021), and finally it is correlated to the *Assilina spira- Lokharatina huntii* zone of Al-Qayim and Ghafor (2022), who considered it of early Bartonian age.

Age: Middle Eocene (Early Bartonian)

Discussion

The difference in age-dating of the studied area is explained by that the stratigraphic succession of the Naopurdan Group at Chwarta has difference in age; whereas at Chwarta, the Nummulitic limestone of the Naopurdan Group is considered as the lower part of the Naopurdan Group, which overlays the Red Bed Series of the Paleocene. The depositional sediments of the Iraqi red-bed basin were oriented NW–SE and bordered by the sea to the SW and a continental block to the NE (Bellen, et al. 1959). The palaeogeography of the continental block that acted as the source for most of the red-bed components is composed of three main units: (1) the Zagros Fold Belt; (2) the narrow Zagros Thrust Zone; and (3) the Sanandaj–Sirjan Zone that forms the western area of the Iranian plate.

The general larger benthic foraminifera in the section implies shallowing of the depositional environment. According to the previous studies Upper Paleocene–Middle Eocene was determined for Walash-Naopurdan Group sediments. Al-Qayim, et al. (2014) recorded the Early Eocene age of the group by direct dependence on the assemblages of foraminifera. Middle Eocene (Lutetian) age was recorded for Walash-Naopurdan Group by Al-Banna and Al-Mutwali (2008), and Kharajany (2018). Daoud (2019) studied the Nummulitic Limestone of the group and determined the Late Paleocene - Early Eocene age. Finally Ahmad, et al. (2022), recorded the middle Eocene/Lutetian age of the group. But the interpretation of the present work at Chwarta section which is studied for the first time in detail with the viewpoint of microfossils and biostratigraphy is interpreted as that the studied area is rich in large benthic foraminifera with few planktic foraminiferas in addition to microfossils of non-foraminifera and the age of Naopurdan Group is shown to be close to Ypressian-Early Bartonian, as follows:-

a- *Nummulites globulus* -*Nummulites planulatus* assemblage Zone. Ypressian age:

(Ghazi, et al. 2010; Racy, 2016; Shreif, et al. 2019; Hadi, et al. 2021; Okur and Kutluk 2020) recognized these benthic foraminifera from the Ypressian age *Alveolina laxa*, *Alveolina globula*, *Alveolina regularis*, *Alveolina globosa*, *Alveolina pastillicata*, *Alveolina pissiformis*, *Alveolina rakoveci gueroli*, *Alveolina sirelii*, *Nummulites fraasi*, *Nummulites solitaires*, *Nummulites minutus*, *Lockhartia tipperi Davies*, *Lockhartia conditi*.

b- *Nummulites mamillatus*- *Nummulites alsharhani* Interval Zone. Lutetian age:

According to Olempska (1973) and Boukhary et. al. (2002), these benthic foraminifera (*Nummulites alsharhani*, *Discocyclina varians*, *Discocyclina ranikotensis*), are the index fossils of the Lutetian age.

c- *Nummulites fabianii* - *Assilina exponens* assemblage Zone. Bartonian age:

Bartonian age is defined based on these index benthic foraminifera (*Nummulites ptukhiani*, *N. perforatus*, *Nummulites catari*), (Kövecsi, et al. 2015; Saraswati, et al. 2017).

Conclusion

The examination of the early Eocene section from the Naopurdan Group at the Imbricate/Thrust Zone from the Kurdistan region of northeast Iraq reveals important conclusions which can be summarized as follows:

a. The sequence generally consists of shallow marine carbonates depending on the large number of benthic foraminifera.

b- Three Larger benthic foraminiferal biozones are recognized that unified from old to young: *Nummulites globulus*-*Nummulites planulatus* Assemblage Zone; *Nummulites mamillatus*-*Nummulites alsharhani* Interval Zone and *Nummulites fabianii*-*Assilina exponens* Assemblage Zone. From this it is concluded that the age of the section extends from Early to Middle Eocene (Ypressian to Early Bartonian).

c- Biostratigraphic correlations with other biozones from other parts inside and outside of Iraq.

Acknowledgments

The authors acknowledge the support of the Departments of Geology in Sulaimani University. We are grateful to the staff of the Department especially Professor Dr. Kamal Haji Karim, Pshtiwan Ahmad, Zana Mahmood and Asaad Ibrahim for field assistance. We thank a lot all the reviewers for their excellent comments.

References

- Ahmad, M.P., Kharajiany, S.O.A., and Al-Khafaf., A.O.S., 2022. New contribution to biostratigraphy of aforaminiferal-condensed section within the Naopurdan Group, Chwarta, Sulaimani, Kurdistan Region/Iraq. Carbonates and Evaporites, 37(3), pp. 1-10, <https://doi.org/10.1007/s13146-022-00797-y>
- Ahmad, S., 2010. Paleogene Larger Benthic Foraminiferal Stratigraphy and Facies distribution: Implications for Tectonostratigraphic Evolution of the Kohat Basin, Potwar Basin and the Trans Indus Ranges (TIR) northwest Pakistan. Ph. D. Thesis, University of Edinburgh.
- Al-Banna, N., and Al-Mutwali, M., 2008. Microfacies and age determination of the sedimentary sequence withinWalash volcano-sedimentary group, Mawat Complex, northeast Iraq. Tikrit J Pure Science 13, pp. 88–95.
- Al Fattah, A., N., Al-Juboury, A.I., and Ghafor, I.M., 2018. Rock Magnetic Properties during the Paleocene-Eocene Thermal Maximum (PETM): Records from P/E boundary Sections (Sinjar, Shaqlawa) in Iraq. Iraqi National Journal of Earth Sciences, 18,(1), pp. 55-74. <http://dx.doi.org/10.33899/earth.2021.170031>.
- Al Fattah, A., N., Al-Juboury, A.I., and Ghafor, I.M., 2020a. Paleocene-Eocene Thermal Maximum Record of NorthernIraq: Multidisciplinary Indicators and an Environmental Scenario. ordan Journal of Earth and Environmental Sciences, 1 (2), pp. 126-145.
- Al Fattah, A.N., Al-Juboury, A.I., and Ghafor, I.M., 2020b. Significance of Foraminifera during the Paleocene-Eocene Thermal Maximum (RETM) in the Aaaliji and Kolosh Formation, North and Northeast Iraq. Iraqi Bulletin of Geology and Mining, 16(2), pp. 33-50.
- Ali, S.A., Sleabi, R.S., Talabani, M.J., and Jones B.G., 2017. Provenance of the Walash-Naopurdan back-arcearc clastic sequences in the Iraqi Zagros Suture Zone, Journal of African Earth Sciences, 125, pp. 73-87.
- Al-Kadhimi, J.A. M., Sissakian, V.K., Fattah, A.S., and Deikran, D.B., 1996. Tectonic map of Iraq, scale 1: 1000000, 2nd editl, Geosurv, Baghdad, Iraq, pp. 1-3.
- Al-Mehaidi, H.M., 1975. Tertiary Nappe in Mawat Range, N.E Iraq, Jour. Geol. Soc. Iraq, 8, pp. 31-44.
- Al Nuaimy, Q.A., Sharbazheri, K.M., and Ghafor, I.M., 2020. Cretaceous / Paleogene Boundary Analysis by Planktic Foraminiferal Biozonation in the Western Zagros Fold-Thrust Belt (Smaqli valley), Sulaimani Governorate, NE-Iraq, Kirkuk University Journal Scientific Studies (KUJSS), 15(3), pp. 45-81. <http://dx.doi.org/10.32894/kujss>.
- Al-Kubaysi, K.N., 2014. The biostratigraphy, microfacies and depositional environment of the Damam Formation In Borehole No.1, South Samawa area,Southern Desert, Iraq, Iraqi Bulletin of Geology and Mining,10(1), pp. 1-20.
- Al-Qayim, B., Ghafor, I.M, and Jaff, R., 2014. Contribution to the stratigraphy of the Walash Group, Sulaimani area, Kurdistan, Iraq. Arab. J. Geosci. <http://dx.doi.10.1007/s12517012-0809-x>.

- Al-Qayim, B., and Ghafor I.M., 2022. Biostratigraphy and Paleoenvironments of Benthic Foraminifera from Lower Part of the Damlouk Member, Western Desert, Iraq. Iraqi Journal of Science, (IJS), 63 (11), pp. 4799-4817.
- Bolton, C.M.G., 1958d. The geology of the Ranya area. Manuscript report, (271), GEOSURV, Baghdad.
- Boukhary, M., Abdelghany, O., and Bahr, S., 2002. Nummulites alsharhani n.sp. (Late Lutetian) from Jabal Hafit and Al Faiyah: Western side of the Northern Oman Mountains, United Arab Emirates, Revue Paléobiol., Genève, 21 (2), pp. 575-585.
- Bolli H.M., 1966. Zonation of Cretaceous to Pliocene marine sediments based on planktonic foraminifera. Bol. Inform. Assoc. Venezolana Geol. Min. Ret. En 9(1), pp. 3-32.
- Buday, T., 1980. The Regional Geology of Iraq. Stratigraphy and Palaeogeography. GEOSURV, Baghdad, Iraq, 445 P.
- Buday, T and Jassim, S.Z., 1987. The Regional geology of Iraq: Tectonism Magmatism, and Metamorphism. M.J Abbas and Jassim, S. Z (Eds), GEOSURV, Baghdad, Iraq, 445 P.
- Daoud, H. S., 2019. Microfaunal Assemblages and microfacies of Nummulitic Limestone of Naopurdan Group from Chwarta Area, Sulaimaniya Governorate, Northern Iraq, JZS 21 (1): (Part-A). DOI: <https://dx.doi.org/10.17656/jzs.10745>.
- Ferrandez-Canadell, C., and Bover-Arnal, T., 2017. Late Chattian larger foraminifera from the Prebetic Domain (Se Spain): New Data on Shallow Benthic Zone 23. Palaios, 32(1), pp. 83–109. DOI: <https://doi.10.2110/palo.2016.010>.
- Ghazi, S., ALI, A., Hsnief, T., Sharif, S., and Arif, S.J., 2010. Larger Benthic Foraminifera Assemblages from the Early Eocene Chor Galifformation, Saltrange, Pakistan, Geol. Bull. Punjab Univ. 45, pp. 83-91.
- Gedik, F., 2014. Benthic Foraminiferal Fauna of Malatya Oligo- Miocene Basin, (Eastern Taurids, Eastern Turkey). Bulletin of the Mineral Research and Exploration , 149(149): pp. 93–136.
- Ghafor, I.M., 1988. Planktonic foraminifera and biostratigraphy of the Aaliji Formation and the nature of its contact with the Shiranish Formation in Well Tel-Hajar No. 1. Sinjar area, Northwestern Iraq. Unpubl. Thesis, Geol. Dept. Univ. Salahaddin, Iraq, 206 P.
- Ghafor, I.M., and Kareem, K.H., 1999. Biostratigraphy of Upper part of the Kolosh Formation from Sartaq-Bamo Northeastern Iraq. J. Dohok Univ., Spec. Issue 2(4), pp. 493 510.
- Ghafor, I.M., 2000. A spores and pollen of Upper Cretaceous Lower Tertiary in Higran area, Kurdistan, Iraq. Zanco(Journal of Pure and applied Science), 12(2), pp. 47-62.
- Ghafor, I.M., and Qadir, M., 2009. Larger Foraminifera (Alveolinidae, Soritidae and Nummulitidae) from the Former Qulqula Conglomerate Formation, Kurdistan Region, Northeastern Iraq, Iraqi Journal of Earth Sciences, 9(1), pp. 35-54. DOI: <https://doi.10.33899/earth.2009.40591>.
- Ghafor, I.M., and Al-Qayim, B., 2021. Planktic Foraminiferal Biostratigraphy of the Upper Part of the Damlouk Member, Ratga Formation, Western Desert, Iraq, Iraqi National Journal of Earth Sciences, 21 (2), pp. 49-62. DOI: <https://doi.10.33899/earth.2021.170385>.
- Hadi, M., Consorti, L., Vahidinia, M., Parandavar, M., and Zoraghi, M., 2021. Ypresian Alveolina and calcareous nannofossils from the south Sabzevar area (Central Iran): biostratigraphic, taxonomic and paleobiogeographic implications. Micropaleontology, 67(1), pp. 31-52. <http://doi.org/10.47894/mpal.67.1.04>.
- Heron, A.M. and Lees, C.M., 1943. The zone of Nappe in Iraq Kurdistan. Min. of Development Report, SOM Library, No.132, Baghdad.
- Hottinger, L., 1960. Recherches sur les Alvéolines du Paléocène et de l'Eocène. Schweizerische Palaeontologische Abh & lungen 75-76, 243 P.
- Jassim, S.Z., and Goff, J.C., 2006. Geology of Iraq. Dolin, Prague and Moravian Museum, Brno, 341 P.
- Karim, K.H., 2004. Basin analysis of Tanjero Formation in Sualiamniyah area, NE. Iraq, Unpublished, Ph. D. Thesis, University of Sulaimani, Iraq.

- Karim, K.H., 2005. Some sedimentary and structural evidences of a possible Graben in Mawat-Chuwarta area, NE Iraq. *Iraqi. Journal Earth Science*, 5(2), pp. 9-18.
- Karim, S. A., and Al-Kubaysi, K.N., 2015. New occurrence of Ratha Formation (Eocene) in some subsurface sections in Nasirya, Southern desert, Iraq, *Iraqi Bulletin of Geology and Mining*, 11(1), pp. 27-43.
- Kharajiany, S.O., 2018. Calcareous nannofossils from the Eocene sequence of the Naopurdan Group, Betwat locality, Sulaimaniyah, Kurdistan Region, Iraq. *Iraqi Bulletin of Geology and Mining*, 14 (2), pp. 17-30.
- Kövecsi, SZ.A., Less, GY., Silye, L. and Filipescu, S.1., 2015. New data on the middle Eocene(Bartonian) Nummulites perforatus “banks” from the Transylvanian Basin (Romania), The 10th Romanian Symposium of Paleontology, October 16-17, 2015.
- Loeblich, J.R. and Tappan, H., 1988. Foraminiferal genera and their classification, Van Nostrand Reinhold. 305 P.
- Mirza, T., Mohialdeen, I., Hakarri, S., and Fatah, C., 2016. Geochemical assessment of Naopurdan limestone for cement making—Chwarta area, Kurdistan Region, NE Iraq. *J Zankoy Sulaimani*. <https://doi.org/10.17656/jzs.10484>.
- Numan, N. M. S., 1997. Aplate tectonic scenario for the phanerozoic succession in Iraq. *Iraqi Geological Journal*, 30(2), pp. 85-110.
- Olempska, E., 1973. The Genus Discocyclina (Foraminifera) from the Eocene of the Tatra Mts, Poland, *Acta paleontologica Polonica*, 18:(1).
- Okur, K., and Kutluk, H., 2020. Benthic foraminiferal assemblages from the Safranbolu Formation (Cuisian, Eocene), Northwest Anatolia, Turkey. *Journal of Palaeogeography*, 9 (6), pp. 1-21, <https://doi.org/10.1186/s42501-020-0054-2>.
- Racy, A., 2016. New nummulite (Foraminifera) species from the Eocene of Northern Oman, University of Cambridge on June 16, <http://doi.jm.lyellcollection.org>.
- Roozpeykar, A. and Moghaddam, I.M., 2016. Benthic foraminifera as biostratigraphical and paleoecological indicators: an example from Oligo- Miocene deposits in the SW of Zagros basin, Iran. *Geosci Front* 7(1), pp. 125-140. <https://www.sciencedirect.com/science/article/pii/S1674987115000444>.
- Saraswati, P.K., Anwar, D., and Lahiri, A., 2017. Bartonian reticulate Nummulites of Kutch, *Geodinamica Acta*, 29(2), pp. 14–23, <http://dx.doi.org/10.1080/09853111.2017.1300847>.
- Serra-Kiel, J., Gallardo-Garcia A., Razin P., Robinet, J., Roger, J., and Grelaud Robin, C.C., 2016. Middle Eocene-Early Miocene larger foraminifera from Dhofar (Oman) and Socotra Island (Yemen). *Arab J Geosci*, 9:(5). <https://doi.org/10.1007/s12517-015-243-3>.
- Serra-Kiel, J., Hottinger, L., Caus, E., Drobne, K., Fernandez, C., Jauhri, A.K., Less, G., Pavlovec, R., Pignatti, J., Samso, J.M., Schaub, H., Sirel, E., Strougo, A., Tambareau, Y., Tosquella, Y., and Zakrevskaya, E., 1998. Larger foraminiferal biostratigraphy of the Tethyan Paleocene and Eocene. *Bulletin de la Société Géologique de France*, 169, pp. 281-299.
- Seyrafian, A., and Mojikhalifeh, A.R., 2005. Biostratigraphy of the Late Paleogene-Early Neogene succession, north- central border of the Persian Gulf. *Carbonates Evaporites*, 20(1), pp. 91–97. <https://doi.org/10.1007/BF03175452>.
- Sharbazheri, K.M., Ghafor, I.M., and Mohammed, Q.A., 2009. Biostratigraphy of the Cretaceous/Tertiary boundary in the Sirwan Valley (Sulaimani Region, Kurdistan, NE Iraq), *Geologica carpathica*, 60(5), pp. 381-396. <https://doi.10.2478/v10096-009-0028-x>.
- Sharbazheri, K. M., Ghafor, I.M., and Mohammed, Q.A., 2011. Biostratigraphy of the Cretaceous/Paleogene boundary in the Dokan area, Sulaimaniyah, Kurdistan Region, NE Iraq, *Iraqi Bulletin of Geology and Mining*, 7(3), pp. 1-24.
- Shreif A., Boukhary M., Abul-Nasr R. A., and Obaidalla, N.A., 2019. Ypresian Nummulites and their stratigraphic significance from El-Guss Abu Said plateau, Farafra Oasis,

Western Desert, Egypt. *Arabian Journal of Geosciences*, (12)72, pp. 1-14.
<https://doi.org/10.1007/s12517-018-4195-x>.

Sirel, E., Özgen-Erdem, N., and Kangal, Ö., 2013. Systematics and biostratigraphy of Oligocene (Rupelian-early Chattian) foraminifera from lagoonal-very shallow water limestone in the eastern Sivas basin (central Turkey). *Geologia Croatica*, 66(2), pp. 83-110. <https://doi.org/10.4154/GC.2013.07>.

Surdashi, A.M., 1997. Depositional and post depositional deformation of Naopurdanlimestone Unit from Chuwarta- Mawat area, Northeast of Iraq. *Iraqi Geological Journal*, 30 (2), pp. 118-12.

Plate 1




Plate 1: a. *Alveolina elliptica* (Sowerby, 1840), oblique section, sample 33; b. *Assilina granulosa* (d'Archiac, 1850), axial section, sample 26; c. *Assilina dandotica* (Davies, 1937), axial section, sample 32; d. *Alveolina solida* (Hottinger, 1960), axial section, sample 35; e. *Alveolina frumentiformis* (Schwager, 1883), oblique section, sample 33; f. *Alveolina subovata* (Wan, 1990), oblique section, sample 35; g. *Nummulites planulatus* (Lamarck, 1804), equatorial section, sample 34; h. *Nummulites globulus* (Leymerie, 1846), axial section, sample 30; i. *Discocyclus ranikotensis* (Davies, 1927), equatorial section, sample 38; j. *Alveolina laxa* (Hottinger, 1960), axial section, sample 1; k. *Alveolina oblonga* (d'Orbigny, 1826), axial section, sample 33; l. *Alveolina fornasinii* (Checchia-Rispoli, 1907), axial section, sample 35; m. *Assilina laminosa* (Gill, 1953), axial section, sample 36; n. *Nummulites exilis* (Douvillé, 1919), axial section, sample 30; o. *Assilina spira* (Roissy, 1805), axial section, sample 37; p. *Assilina dandotica* (Davies, 1937), axial section, sample 27; q. *Nummulites mamillinus* (Douvillé, 1924), axial section, sample 30.

Plate 2




Plate 2: a. *Discocyclus sella* (d'Archiac, 1850), equatorial section, sample 27; b. *Alveolina munieri* (Hottinger, 1960), axial section, sample 33; c. *Assilina exponens* (Sowerby, 1840), axial section, sample 34; d. *Discocyclus archiaci* (Schlumberger, 1903), equatorial section, sample 18; e. *Lepidocyclus* sp., equatorial section, sample 28; f. *Discocyclus varians* (Kaufmann, 1867), equatorial section, sample 28; g. *Discocyclus* sp., equatorial section, sample 26; h. *Morozovella formosa* (Bolli 1957), axial section, sample 14; i. *Nummulites perforatus* (Montfort, 1808), axial section, sample 26; j. *Nummulites discorbinus* (Schlotheim, 1820), sub-axial section, sample 26; k. *Discocyclus dispansa* (Sowerby, 1840), equatorial section, sample 28; l. *Nummulites globulus* (Leymerie, 1846), sub-axial section, sample 34; m. *Nummulites djokdjokartae* (Martin, 1881), sub-axial section, sample 27.




Plate 3: a. *Discocyclus archiaci* (Schlumberger, 1903), axial section, sample 4; b. Coral, equatorial section, sample 1; c. *Alveolina ellipsoidalis* (Schwager, 1883), oblique section, sample 2; d. Gastropoda, axial section, sample 4; e. *Heterostegina* sp., axial section, sample 10; f. *Asterocyclus matanzensis* (Cole 1957), equatorial section, sample 12; g. *Nummulites fabianii* (Prever, 1905), sub-axial section, sample 10; h. *Heterostegina* sp., equatorial section, sample 10; i. *Sphaerogypsina* sp., equatorial section, sample 5; j. *Pyrgo* sp., oblique section, sample 8; k. *Lithophylum* sp., oblique section, sample 1; l. Bryozoan, equatorial section, sample 13; m. *Quinqueloculina* sp., equatorial section, sample 6; n. *Miliolina*, equatorial section, sample 1; o. *Discocyclus* sp., equatorial section, sample 5; p. *Subbotina linaperta*, equatorial section, sample 11; q. calcareous algae, oblique section, sample 11.




Plate 4: a. *Lockhartia tipper*, equatorial section, sample 34; b. *Alveolina sirelii*, oblique section, sample 32; c. *Alveolina regularis* (Hottinger, 1960), oblique section, sample 35; d. *Alveolina pissiformis*, axial section, sample 32; e. *Alveolina rakoveci gueroli* (Drobne, 1977), oblique section, sample 33; f. *Orbitolites complanatus* (Lamarck), axial section, sample 33; g. Calcareous algae, sample 11; h. *Alveolina pastiliicata*, oblique section, sample 33; i. *Orbitolites complanatus* (Lamarck), equatorial section, sample 35; j. *Alveolina globosa* (Leymerie), oblique section, sample 35; k. *Nummulites mamillatus* (Fichtel and Moll, 1798), sub-axial section, sample 35; l. *Nummulites solitaires*, equatorial section sample 35; m. *Nummulites catari*, equatorial section, sample 3; n. *Nummulites ptukhiani* (Z.D. Kacharava, 1969), sub-axial section, sample 10; o. *Nummulites alsharhani*, sub-axial section sample 11.