

Mohammed Salih Mahdy Hussan

New Results of Fuzzy Continuous Functions on Fuzzy Topological spaces

Mohammed Salih Mahdy Hussan

Baghdad, Iraq.

Al-Mustansiriyah University

College of Education

Department of Mathematics

Abstract

In this paper we give further results about concerning certain to fuzzy continuous functions defined from fuzzy topological space to another fuzzy topological space and to show the relationships between fuzzy continuous functions where we confine our study to some of their types such as, fuzzy θ -continuous function, fuzzy strong θ -continuous and fuzzy δ -continuous function.

Keywords: fuzzy topological space, fuzzy δ -closure, fuzzy θ -closure, fuzzy θ -continuous function, fuzzy strong θ -continuous, fuzzy δ -continuous function.

نتائج جديدة للدوال المستمرة الضبابية على الفضاءات التبولوجية الضبابية

محمد صالح مهدي حسين

بغداد، العراق / الجامعة المستنصرية / كلية التربية / قسم الرياضيات

الخلاصة

في هذا البحث نعطي نتائج أخرى حول بعض الأنواع للدوال المستمرة الضبابية من فضاء التبولوجي ضبابي إلى فضاء تبولوجي آخر وإظهار العلاقات بين الدوال المستمرة الضبابية حيث نتحصر دراستنا حول بعض الأنواع مثل

fuzzy θ -continuous function, fuzzy strong θ -continuous, fuzzy δ -continuous function.

الكلمات الدلالية

الفضاء التبولوجي الضبابي، fuzzy δ -closure, fuzzy θ -closure, fuzzy θ -continuous function,

fuzzy strong θ -continuous, fuzzy δ -continuous function.

Mohammed Salih Mahdy Hussan

1.Introduction

The study of fuzzy sets was initiated with the famous paper of Zadeh [12] in 1965, and thereafter the paper of Chang [3] in 1968 paved the way for the subsequent tremendous growth of the numerous fuzzy topological concepts. The concepts of fuzzy δ -closure and fuzzy θ -closure in fuzzy topological space were introduced by Ganguly and Saha [10] and Mukherjee and Sinha [6] respectively. In this paper we investigate some properties of fuzzy strong θ -continuous, fuzzy θ -continuous and fuzzy δ -continuous.

2.Preliminaries

The definitions and results which are used in this paper concerning fuzzy topological spaces have already taken certain standard shape. A fuzzy set $\tilde{A}$ in a fuzzy topological space $(X, \tilde{T})$ (fts X , for short) is said to be quasi-coincident (q -coincident, for short) with a fuzzy set $\tilde{B}$, denoted by $\tilde{A} q \tilde{B}$ if and only if there exists $x \in X$ such that $\mu_{\tilde{A}}(x) + \mu_{\tilde{B}}(x) > 1$ [4]. A fuzzy set in $(X, \tilde{T})$ is said to be quasi-neighborhood (q -nbd, for short) of a fuzzy point X_r or $\tilde{p}$ (where X is the support and r is the value of the fuzzy point $0 < r \leq 1$) if and only if there exists a fuzzy open set $\tilde{B}$ such that $\tilde{p} q \tilde{B} \leq \tilde{A}$ [4]. For two fuzzy sets $\tilde{A}, \tilde{B}$ in X , $\tilde{A} \subseteq \tilde{B}$ iff $\tilde{A}$ is not q -coincident with $\tilde{B}^c$ (complement of $\tilde{B}$) to be denoted by $\tilde{A} \not q \tilde{B}^c$. By $cl(\tilde{A})$ and $int(\tilde{A})$ we shall denote respectively the fuzzy closure and fuzzy interior of a fuzzy set $\tilde{A}$ in a fts X . A fuzzy point $\tilde{p} \in cl(\tilde{A})$ if and only if each q -nbd of $\tilde{p}$ is q -coincident with $\tilde{A}$ [4].

Definition 2.1 [4]. A fuzzy set $\tilde{A}$ of a fuzzy topology space $(X, \tilde{T})$ is called

- (a) A fuzzy regular closed set of X , if $cl(int(\tilde{A})) = \tilde{A}$
- (b) A fuzzy regular open set of X , if $int(cl(\tilde{A})) = \tilde{A}$.

Remark 2.2 [2],[7]. Every fuzzy regular open set is a fuzzy open set and every fuzzy regular closed set is a fuzzy closed set.

Definition 2.3 [11]. A fuzzy point $\tilde{p}$ in $(X, \tilde{T})$ is said to be a fuzzy (δ -cluster point) of a fuzzy set $\tilde{A}$ if for each fuzzy open q -nbd $\tilde{U}$ of $\tilde{p}$, $int(cl(\tilde{U})) q \tilde{A}$.

The set of all fuzzy (δ -cluster points) of $\tilde{A}$ is called the fuzzy (δ -closure) of $\tilde{A}$ and is denoted by

New Results of Fuzzy Continuous Functions on Fuzzy Topological spaces

Mohammed Salih Mahdy Hussan

$[\tilde{A}]_\delta$. A fuzzy set $\tilde{A}$ is fuzzy (δ -closed) if and only if $(\tilde{A} = [\tilde{A}]_\delta)$ and the complement of a fuzzy set (δ -closed set) is fuzzy (δ -open) set.

Remark 2.4 [1],[5] [6]. Every fuzzy regular open set is a fuzzy δ -open set and a fuzzy δ -open set is

a union of fuzzy regular open sets and hence is a fuzzy open set and for a fuzzy open set $\tilde{A}$ in a fts

$$(X, \tilde{T}), \text{cl}(\tilde{A}) = [\tilde{A}]_\delta = [\tilde{A}]_0.$$

Definition 2.5 [6]. A fuzzy point $\tilde{p}$ is said to be fuzzy θ -cluster point of a fuzzy set $\tilde{A}$ if and only if for every open q-nbd. $\tilde{U}$ of $\tilde{p}$, $\text{cl}(\tilde{U})$ is quasi-coincident with $\tilde{A}$. The set of all fuzzy θ -cluster points of $\tilde{A}$ is called the fuzzy θ -closure of $\tilde{A}$ and will be denoted by $[\tilde{A}]_\theta$.

A fuzzy set $\tilde{A}$ will be called fuzzy θ -closed if and only if $\tilde{A} = [\tilde{A}]_\theta$ and the complement of fuzzy θ -closed set is a fuzzy θ -open.

Definition 2.6 [4]. Let $(X, \tilde{T})$ be a fts, and A is any subset of X ; then the family $\tilde{T}_A$, defined by $\tilde{T}_A = \{ \tilde{B} / A : \tilde{B} \in \tilde{T} \}$, which is obviously a fuzzy topology for A , is called relative fuzzy topology, or the relativization of $\tilde{T}$ to A . Such a fuzzy topological space $(A, \tilde{T}_A)$ is called a subspace of $(X, \tilde{T})$.

3. Characterizations of fuzzy θ – continuity, fuzzy strong θ – continuity, and fuzzy δ -continuity.

Definition 3.1 [8]. A function $f : (X, \tilde{T}) \longrightarrow (Y, \tilde{T}')$, is said to be

- (a) Fuzzy θ -continuous (f. θ .c, for short) if for each fuzzy point $\tilde{p}$ in $(X, \tilde{T})$ and each fuzzy open q-nbd. $\tilde{V}$ of $f(\tilde{p})$, there exists fuzzy open q-nbd. $\tilde{U}$ of $\tilde{p}$ such that $f(\text{cl}(\tilde{U})) \subseteq \text{cl}(\tilde{V})$.

New Results of Fuzzy Continuous Functions on Fuzzy Topological spaces

Mohammed Salih Mahdy Hussan

(b) Fuzzy strong θ -continuous (f.s. θ .c, for short) if for each fuzzy point $\tilde{p}$ in $(X, \tilde{T})$ and each fuzzy open q-nbd. $\tilde{V}$ of $f(\tilde{p})$, there exists fuzzy open q-nbd. $\tilde{U}$ of $\tilde{p}$ such that $f(\text{cl}(\tilde{U})) \subseteq \tilde{V}$.

Definition 3.2 [9]. A function $f : (X, \tilde{T}) \longrightarrow (Y, \tilde{T}')$, is said to be fuzzy δ -continuous at a fuzzy point $\tilde{p}$ in $(X, \tilde{T})$, if for each fuzzy open q-nbd. $\tilde{V}$ of $f(\tilde{p})$, there exists an open fuzzy nbd. $\tilde{U}$ of $\tilde{p}$ such that $f(\text{int}(\text{cl}(\tilde{U}))) \subseteq \text{int}(\text{cl}(\tilde{V}))$.

Also, the next theorem appears in [6], [8] with uncompleted proof and hence we give the complete details of the proof.

Theorem 3.3. For a function $f : (X, \tilde{T}) \longrightarrow (Y, \tilde{T}')$, each of the following statements equivalent:

- (a) f is fuzzy θ -continuous function.
- (b) For each fuzzy set $\tilde{A}$ in X , $f([\tilde{A}]_0) \subseteq [f(\tilde{A})]_0$.
- (c) For each fuzzy set $\tilde{B}$ in Y , $[f^{-1}(\tilde{B})]_0 \subseteq f^{-1}([\tilde{B}]_0)$.
- (d) For each θ -closed set $\tilde{B}$ in Y , $f^{-1}(\tilde{B})$ is a fuzzy θ -closed in X .
- (e) For each θ -open set $\tilde{B}$ in Y , $f^{-1}(\tilde{B})$ is a fuzzy θ -open in X .
- (F) For each open set $\tilde{B}$ in Y , $[f^{-1}(\tilde{B})]_0 \subseteq f^{-1}(\text{cl}(\tilde{B}))$.

Proof:

(a) $\Rightarrow$ (b). Let $\tilde{p}$ be a fuzzy point $\in [\tilde{A}]_0$ and let $\tilde{U}$ any open q-nbd. of $f(\tilde{p})$. Then there exists an open q-nbd. $\tilde{V}$ of $\tilde{p}$ such that $f(\text{cl}(\tilde{V})) \subseteq \text{cl}(\tilde{U})$, i.e., f is a fuzzy θ -continuous.

Now $\tilde{p} \in [\tilde{A}]_0$, $\text{cl}(\tilde{V}) \cap \tilde{A} \Rightarrow f(\text{cl}(\tilde{V})) \cap f(\tilde{A})$

$$\text{cl}(\tilde{U}) \cap f(\tilde{A}) \Rightarrow f(\tilde{p}) \in [f(\tilde{A})] \Rightarrow \tilde{p} \in f^{-1}([f(\tilde{A})]_0).$$

New Results of Fuzzy Continuous Functions on Fuzzy Topological spaces

Mohammed Salih Mahdy Hussan

Thus, $[\tilde{A}]_\theta \subseteq f^{-1}([\tilde{A}]_\theta)$ so that $f([\tilde{A}]_\theta) \subseteq [f(\tilde{A})]_\theta$

(b) $\Rightarrow$ (c). By (b) $f([f^{-1}(\tilde{B})]_\theta) \subseteq [ff^{-1}(\tilde{B})]_\theta \subseteq [\tilde{B}]_\theta$.

Which implies that $[f^{-1}(\tilde{B})]_\theta \subseteq f^{-1}([\tilde{B}]_\theta)$.

(c) $\Rightarrow$ (d). We have $[\tilde{B}]_\theta = \tilde{B}$

Now by (c) $\Rightarrow [f^{-1}(\tilde{B})]_\theta \subseteq f^{-1}([\tilde{B}]_\theta) = f^{-1}(\tilde{B}) \Rightarrow [f^{-1}(\tilde{B})]_\theta \subseteq f^{-1}(\tilde{B})$

For each fuzzy θ -closed set $\tilde{B}$ in Y . $f^{-1}(\tilde{B})$ is a fuzzy θ -closed in X .

(e) $\Rightarrow$ (f). Since $\tilde{B}$ be a fuzzy open in Y , by remark (2.4) $\text{cl}(\tilde{B}) = [\tilde{B}]_\theta$ and we have from (c)

$[f^{-1}(\tilde{B})]_\theta \subseteq f^{-1}([\tilde{B}]_\theta)$. We have $[f^{-1}(\tilde{B})]_\theta \subseteq f^{-1}(\text{cl}(\tilde{B}))$

(f) $\Rightarrow$ (a). Let $\tilde{p}$ be a fuzzy point in X and $\tilde{V}$ be a fuzzy open q-nbd. of $f(\tilde{p})$. Then

$$\begin{aligned} \mu_{(1-\text{cl}(\tilde{V}))}(x) \text{ is a fuzzy open in } Y. \text{ By (f), we have } [f^{-1}(1 - \text{cl}(\tilde{V}))]_\theta &\subseteq f^{-1}(\text{cl}(1 - \text{cl}(\tilde{V}))) \\ &= 1 - f^{-1}(\text{int}(\text{cl}(\tilde{V}))). \end{aligned}$$

Then there exists a fuzzy open q-nbd. $\tilde{U}$ of $\tilde{p}$, such that $\text{cl}(\tilde{U}) \not\subseteq (1 - f^{-1}(\text{cl}(\tilde{V})))$ so that

$f(\text{cl}(\tilde{U})) \subseteq \text{cl}(\tilde{V})$. Hence, f is a fuzzy θ -continuous function.

Both theorems appear in [8], [5] respectively, without a complete proof. We presented the proof completely for its importance as we had done in the previous theorems.

Theorem 3.4. For a function $f : (X, \tilde{T}) \longrightarrow (Y, \tilde{T}')$, the following are equivalent:

- (a) f is fuzzy strong θ -continuous.
- (b) $f([\tilde{A}]_\theta) \subseteq \text{cl}(f(\tilde{A}))$ for every fuzzy set $\tilde{A}$ in X .
- (c) $[f^{-1}(\tilde{B})]_\theta \subseteq f^{-1}(\text{cl}(\tilde{B}))$ for every fuzzy set $\tilde{B}$ in Y .
- (d) The inverse image of every fuzzy closed set in Y is fuzzy θ -closed in X , i.e., every fuzzy closed set $\tilde{B}$ in Y , $f^{-1}(\tilde{B})$ is fuzzy θ -closed in X .

New Results of Fuzzy Continuous Functions on Fuzzy Topological spaces

Mohammed Salih Mahdy Hussan

(e) The inverse image of every fuzzy open set in Y is fuzzy θ -open in X , i.e., every fuzzy open set $\tilde{B}$ in Y , $f^{-1}(\tilde{B})$ is a fuzzy θ -open in X .

Proof:

(a) $\Rightarrow$ (b). Let $\tilde{p} \in [\tilde{A}]_0$ and $\tilde{V}$ be fuzzy open q-nbd. of $f(\tilde{p})$.

By (a), there exists a fuzzy open q-nbd. $\tilde{U}$ of $\tilde{p}$ such that $f(\text{cl}(\tilde{U})) \subseteq \tilde{V}$. Now, we have $\tilde{p} \in [\tilde{A}]_0$

$$\Rightarrow \text{cl}(\tilde{U}) \text{ q } \tilde{A} \Rightarrow f(\text{cl}(\tilde{U})) \text{ q } f(\tilde{A}) \Rightarrow \tilde{V} \text{ q } f(\tilde{A}) \Rightarrow f(\tilde{p}) \in \text{cl}(f(\tilde{A})) \Rightarrow \tilde{p} \in f^{-1}(\text{cl}(f(\tilde{A})))$$

Hence, $[\tilde{A}]_0 \subseteq f^{-1}(\text{cl}(f(\tilde{A})))$ and so $f([\tilde{A}]_0) \subseteq \text{cl}(f(\tilde{A}))$.

(b) $\Rightarrow$ (c). Let $\tilde{B}$ be a fuzzy set in Y . By (b) $f([f^{-1}(\tilde{B})]_0) \subseteq \text{cl}(f(f^{-1}(\tilde{B}))) \Rightarrow f([f^{-1}(\tilde{B})]_0) \subseteq \text{cl}(\tilde{B})$

$$\Rightarrow f^{-1}(f([f^{-1}(\tilde{B})]_0)) \subseteq f^{-1}(\text{cl}(\tilde{B})) \Rightarrow [f^{-1}(\tilde{B})]_0 \subseteq f^{-1}(\text{cl}(\tilde{B}))$$

(c) $\Rightarrow$ (d). Let $\tilde{B}$ be a fuzzy closed in Y .

By (c), we have $[f^{-1}(\tilde{B})]_0 \subseteq f^{-1}(\text{cl}(\tilde{B})) = f^{-1}(\tilde{B})$ (since $\tilde{B}$ closed), which implies that

$$f^{-1}(\tilde{B}) = [f^{-1}(\tilde{B})]_0. \text{ Hence, } f^{-1}(\tilde{B}) \text{ is fuzzy } \theta\text{-closed}$$

(c) $\Rightarrow$ (e). Let $\tilde{B}$ be a fuzzy open in Y . Then $\mu_{(1-\tilde{B})}(x)$ is a fuzzy closed and by (d),

$$f^{-1}(1 - \tilde{B}) = 1 - f^{-1}(\tilde{B}) \text{ is fuzzy } \theta\text{-closed. Hence, } f^{-1}(\tilde{B}) \text{ is fuzzy } \theta\text{-open.}$$

(e) $\Rightarrow$ (a). Let $\tilde{p}$ be any fuzzy point in X and $\tilde{V}$ be a fuzzy open q-nbd. of $f(\tilde{p})$. By (e),

$$f^{-1}(\tilde{V}) \text{ is a fuzzy } \theta\text{-open in } X.$$

$$\text{Now, } f(\tilde{p}) \text{ q } \tilde{V} \Rightarrow \tilde{p} \text{ q } f^{-1}(\tilde{V}) \Rightarrow \tilde{p} \notin 1 - f^{-1}(\tilde{V})$$

Hence $\mu_{(1-f^{-1}(\tilde{V}))}(x)$ is fuzzy θ -closed set such that $\tilde{p} \notin 1 - f^{-1}(\tilde{V})$.

New Results of Fuzzy Continuous Functions on Fuzzy Topological spaces

Mohammed Salih Mahdy Hussan

Then, there exists fuzzy open q -nbd. $\tilde{U}$ of $\tilde{p}$, such that $\text{cl}(\tilde{U}) \not\subseteq (1 - f^{-1}(\tilde{V}))$, which implies

$f(\text{cl}(\tilde{U})) \subseteq \tilde{V}$. This shows that f is fuzzy strong θ -continuous function.

Theorem 3.5. Let $f : (X, \tilde{T}) \longrightarrow (Y, \tilde{T}')$, then the following implications holds:

- (a) f is a fuzzy δ -continuous function.
- (b) $f([\tilde{A}]_\delta) \subseteq [f(\tilde{A})]_\delta$ for every fuzzy set $\tilde{A}$ in X .
- (c) $[f^{-1}(\tilde{B})]_\delta \subseteq f^{-1}([\tilde{B}]_\delta)$ for every fuzzy set $\tilde{B}$ in Y .
- (d) For every fuzzy δ -closed set $\tilde{B}$ in Y , $f^{-1}(\tilde{B})$ is a fuzzy δ -closed in X .
- (e) For every fuzzy δ -open set $\tilde{B}$ in Y , $f^{-1}(\tilde{B})$ is fuzzy δ -open in X .
- (f) For every fuzzy regular open set $\tilde{B}$ in Y , $f^{-1}(\tilde{B})$ is δ -open in X .

Proof:

(a) $\Rightarrow$ (b). We show that for a subset $\tilde{A}$ of X , $f([\tilde{A}]_\delta) \subseteq [f(\tilde{A})]_\delta$

Let $\tilde{p}$ be a fuzzy point in Y such that $\tilde{p} \in f([\tilde{A}]_\delta)$. Then $\tilde{p} = f(\tilde{q})$, where $\tilde{q} \in [\tilde{A}]_\delta$.

It may be seen that if for a fuzzy subset $\tilde{A}$ of X , $\tilde{p} \in f(\tilde{A})$ then there is an $x \in \text{supp } \tilde{A}$ such that $f(x) = y$; Let $\tilde{U}$ be a regularly open q -nbd. of $\tilde{p}$; there exists a regularly open q -nbd. $\tilde{V}$ of $\tilde{q}$ such that $f(\tilde{V}) \subseteq \tilde{U}$. Now, $\tilde{q} \in [\tilde{A}]_\delta$, which implies to $\tilde{V} \not\subseteq \tilde{A}$.

If possible $f(\tilde{V}) \not\subseteq f(\tilde{A})$. Then for all $y \in \tilde{V}$, $f(\tilde{V})(y) + f(\tilde{A})(y) \leq 1$, i.e., $\tilde{V} \not\subseteq \tilde{A} \Rightarrow$ a contradiction

Thus, $f(\tilde{V}) \subseteq f(\tilde{A})$, i.e., $\tilde{U} \subseteq f(\tilde{A})$, i.e., $\tilde{p} \in [f(\tilde{A})]_\delta$, i.e., $f([\tilde{A}]_\delta) \subseteq [f(\tilde{A})]_\delta$.

(b) $\Rightarrow$ (c). By (b) $f([f^{-1}(\tilde{B})]_\delta) \subseteq [f(f^{-1}(\tilde{B}))]_\delta$, so that: $f([f^{-1}(\tilde{B})]_\delta) \subseteq [\tilde{B}]_\delta$ i.e. $[f^{-1}(\tilde{B})]_\delta \subseteq f^{-1}([\tilde{B}]_\delta)$.

New Results of Fuzzy Continuous Functions on Fuzzy Topological spaces

Mohammed Salih Mahdy Hussan

(c) $\Rightarrow$ (d). Let $\tilde{B}$ be a fuzzy δ -closed set in Y . Then $[\tilde{B}]_{\delta} = \tilde{B}$, by (c) $[f^{-1}(\tilde{B})]_{\delta} \subseteq f^{-1}([\tilde{B}]_{\delta}) = f^{-1}(\tilde{B})$.

Thus, $f^{-1}(\tilde{B})$ is a fuzzy δ -closed in X .

(d) $\Rightarrow$ (e). Let $\tilde{B}$ be a fuzzy δ -open set in Y .

Then $\tilde{B}^c$ is a fuzzy δ -closed in Y . By (d), $f^{-1}(\tilde{B})$ is a fuzzy δ -closed in X . but $f^{-1}(\tilde{B}^c) = (f^{-1}(\tilde{B}))^c$

$f^{-1}(\tilde{B})$ is a fuzzy δ -open in X .

(e) $\Rightarrow$ (f). Every regular open fuzzy set is δ -open fuzzy set also. Thus (e) $\Rightarrow$ (f)

(f) $\Rightarrow$ (a). Let $\tilde{B}$ be a regular open q -nbd. of $f(\tilde{p})$ such that $\tilde{p}$ be a fuzzy point in X . by (f) $f^{-1}(\tilde{B})$ is

δ -open in X and $\tilde{p} \in f^{-1}(\tilde{B})$. But every a δ -open fuzzy set is a union of fuzzy regular open sets [Remark (2.4)] and hence is an open fuzzy set.

Hence, there exists a fuzzy regular open sets $\tilde{A}$ such that $\tilde{p} \in \tilde{A} \subseteq f^{-1}(\tilde{B})$.

Now, $f(\tilde{A}) \subseteq \tilde{B}$ so $f(\text{Int}(\text{cl}(\tilde{A}))) \subseteq \text{Int}(\text{cl}(\tilde{B}))$, So, f is a fuzzy δ -continuous function.

4. THE COMPOSITION OF FUZZY CONTINUOUS FUNCTIONS

First, we study the composition of fuzzy θ -continuous, fuzzy strong θ -continuous, and

fuzzy δ -continuous.

Theorem 4.1. If $f : (X, \tilde{T}) \longrightarrow (Y, \tilde{T}')$ and $g : (Y, \tilde{T}') \longrightarrow (Z, \tilde{T}'')$ are a fuzzy θ -continuous functions, then so is $g \circ f : (X, \tilde{T}) \longrightarrow (Z, \tilde{T}'')$.

Proof:

New Results of Fuzzy Continuous Functions on Fuzzy Topological spaces

Mohammed Salih Mahdy Hussan

Let $\tilde{C}$ be a fuzzy open set in $(Z, \tilde{T}'')$ since g is a fuzzy θ -continuous function.

Therefore $[g^{-1}(\tilde{C})]_{\theta} \subseteq g^{-1}(\text{cl}(\tilde{C}))$ (Theorem (3.3), part f). But $[g^{-1}(\tilde{C})]_{\theta}$ is the set of all a fuzzy

θ -cluster point of $g^{-1}(\tilde{C})$, there exist a fuzzy point (called fuzzy θ -cluster point) of a fuzzy set $g^{-1}(\tilde{C})$ s.t for every q -nbd. $\tilde{U}$ of $\tilde{p}$, $\text{cl}(\tilde{U})$ is q – coincident with $g^{-1}(\tilde{C})$ in $(Y, \tilde{T}')$.

Now, we have f is a fuzzy θ -continuous function, $[f^{-1}(g^{-1}(\tilde{C}))]_{\theta} \subseteq f^{-1}(g^{-1}(\text{cl}(\tilde{C})))$

$[(gof)^{-1}(\tilde{C})]_{\theta} \subseteq (gof)^{-1}(\text{cl}(\tilde{C}))$ (since $f^{-1}(g^{-1}(\text{cl}(\tilde{C}))) = (gof)^{-1}(\text{cl}(\tilde{C}))$).

Hence (gof) is a fuzzy θ -continuous function (Theorem (3.3) (f) $\Rightarrow$ (a)).

Theorem 4.2. If $f : (X, \tilde{T}) \longrightarrow (Y, \tilde{T}')$ and $g : (Y, \tilde{T}') \longrightarrow (Z, \tilde{T}'')$ are a fuzzy strong θ -continuous functions, then so is $gof : (X, \tilde{T}) \longrightarrow (Z, \tilde{T}'')$.

Proof:

Let $\tilde{C}$ be a fuzzy open set in $(Z, \tilde{T}'')$ since g is a fuzzy strong θ -continuous function. Hence $g^{-1}(\tilde{C})$ is a fuzzy θ -open in $(Y, \tilde{T}')$ (Theorem (3.4), part (e)). Now f is a fuzzy strong θ -continuous function.

$f^{-1}(g^{-1}(\tilde{C}))$ is a fuzzy θ -open in $(X, \tilde{T})$. But, $f^{-1}(g^{-1}(\tilde{C})) = (gof)^{-1}(\tilde{C})$

This means that $(gof)^{-1}(\tilde{C})$ is a fuzzy θ -open in $(X, \tilde{T})$. Hence, (gof) is a fuzzy strong θ -continuous function.

Theorem 4.3. If $f : (X, \tilde{T}) \longrightarrow (Y, \tilde{T}')$ and $g : (Y, \tilde{T}') \longrightarrow (Z, \tilde{T}'')$ are fuzzy δ -continuous function, then so is $gof : (X, \tilde{T}) \longrightarrow (Z, \tilde{T}'')$.

Proof:

Let $\tilde{V}$ be a fuzzy regularly open set in $(Z, \tilde{T}'')$ since g is a fuzzy δ -continuous. Hence $g^{-1}(\tilde{V})$ is a union of a fuzzy regularly open sets in $(Y, \tilde{T}')$. Every a fuzzy regular open set is fuzzy δ -open set (Remark (2.4)). Therefore $g^{-1}(\tilde{V})$ is a fuzzy δ -open set.

New Results of Fuzzy Continuous Functions on Fuzzy Topological spaces

Mohammed Salih Mahdy Hussan

Assume that $g^{-1}(\tilde{V}) = \bigcup_{\alpha \in \Omega} \tilde{W}_{\alpha}$, where $\tilde{W}_{\alpha}$ is fuzzy regular open in $(Y, \tilde{T}')$.

Now, $f^{-1}(g^{-1}(\tilde{V})) = f^{-1}\left[\bigcup_{\alpha \in \Omega} \tilde{W}_{\alpha}\right] = \bigcup_{\alpha \in \Omega} f^{-1}(\tilde{W}_{\alpha})$ (since If $\tilde{A}_i \subset X$, for every $i \in I$,

then $f\left(\bigcup_{i \in I} \tilde{A}_i\right) = \bigcup_{i \in I} f(\tilde{A}_i)$. Now, f is fuzzy δ -continuous, so $f^{-1}(\tilde{W}_{\alpha})$ is fuzzy δ -open in $(X, \tilde{T})$.

Hence, $\bigcup_{\alpha \in \Omega} f^{-1}(\tilde{W}_{\alpha})$ is fuzzy δ -open in $(X, \tilde{T})$. So $f^{-1}(g^{-1}(\tilde{V}))$ is fuzzy δ -open in $(X, \tilde{T})$

and $f^{-1}(g^{-1}(\tilde{V})) = (g \circ f)^{-1}(\tilde{V})$. This means that $(g \circ f)^{-1}(\tilde{V})$ is fuzzy δ -open in $(X, \tilde{T})$,

Hence $(g \circ f)$ is a fuzzy δ -continuous function (Theorem (3.5) (f) $\Rightarrow$ (a)).

5. THE RESTRICTION OF FUZZY CONTINUOUS FUNCTIONS

In the see from, study the restriction of fuzzy continuous functions of fuzzy θ -continuous, fuzzy strong θ -continuous, and fuzzy δ -continuous and we state and proof with the following theorem.

Theorem 5.1. If $f : (X, \tilde{T}) \longrightarrow (Y, \tilde{T}')$ a fuzzy δ -continuous function and A is open set in X , then

$f/A : (A, \tilde{T}_A) \longrightarrow (Y, \tilde{T}')$ is a fuzzy δ -continuous function.

Proof:

Let $\tilde{V}$ be a fuzzy regular open set in $(Y, \tilde{T}')$

Since f is a fuzzy δ -continuous function, then $f^{-1}(\tilde{V})$ is a fuzzy δ -open set in $(Y, \tilde{T})$ (Theorem (3.5) (f))

Now, $(f/A)^{-1}(\tilde{V}) = f^{-1}(\tilde{V}) \cap A$, but A is open set in X , then $f^{-1}(\tilde{V}) \cap A$.

New Results of Fuzzy Continuous Functions on Fuzzy Topological spaces

Mohammed Salih Mahdy Hussan

Hence, $(f/A)^{-1}(\tilde{V})$ is a fuzzy δ -open in A so (f/A) is a fuzzy δ -continuous function (Theorem (3.5) part (f) $\Rightarrow$ (a)).

Theorem 5.2. If $f : (X, \tilde{T}) \longrightarrow (Y, \tilde{T}')$ is a fuzzy θ -continuous function and A is open set in X then $f/A : (A, \tilde{T}_A) \longrightarrow (Y, \tilde{T}')$ is a fuzzy θ -continuous functions.

Proof:

Let $\tilde{B}$ be a fuzzy open set in $(Y, \tilde{T}')$, Since f is a fuzzy θ -continuous function

Then $[f^{-1}(\tilde{B})]_0 \subseteq f^{-1}(\text{cl}(\tilde{B}))$ is implies (Theorem (3.3) (f))

$$\begin{aligned} \text{Now: } [(f/A)^{-1}(\tilde{B})]_0 &= [A \cap f^{-1}(\tilde{B})]_0 \\ &\subseteq A \cap f^{-1}(\text{cl}(\tilde{B})) \end{aligned}$$

But, $A \cap f^{-1}(\text{cl}(\tilde{B})) = (f/A)^{-1}(\text{cl}(\tilde{B}))$. Therefore, $[(f/A)^{-1}(\tilde{B})]_0 \subseteq (f/A)^{-1}(\text{cl}(\tilde{B}))$

Hence, f/A is a fuzzy θ -continuous function (Theorem (3.3)(f) $\Rightarrow$ (a)).

Theorem 5.3. If $f : (X, \tilde{T}) \longrightarrow (Y, \tilde{T}')$ is a fuzzy strong θ -continuous function and A is open set in X , then $f/A : (A, \tilde{T}_A) \longrightarrow (Y, \tilde{T}')$.

Proof:

Let $\tilde{B}$ be a fuzzy open set in $(Y, \tilde{T}')$, Since f is a fuzzy strong θ -continuous function

Then $f^{-1}(\tilde{B})$ is a fuzzy θ -open in $(X, \tilde{T})$ (Theorem (3.4) (e)).

Now $(f/A)^{-1}(\tilde{B}) = A \cap f^{-1}(\tilde{B})$ but A is open set in X , then $f^{-1}(\tilde{B}) \cap A$ is θ -open in A .

Hence, $(f/A)^{-1}(\tilde{B})$ is a fuzzy θ -open in A , so f/A is a fuzzy strong θ -continuous function (Theorem (3.4) part (e) $\Rightarrow$ (a)).

6. RELATIONSHIP BETWEEN FUZZY CONTINUOUS FUNCTIONS.

We introduce and prove the following theorem where the relationship between fuzzy θ -continuous, fuzzy δ -continuous and fuzzy strongly θ -continuous.

Theorem 6.1. Every fuzzy strong θ -continuous function is a fuzzy δ -continuous function.

Proof:

Let $f : (X, \tilde{T}) \longrightarrow (Y, \tilde{T}')$ be a fuzzy strong θ -continuous and let $\tilde{p}$ a fuzzy point in $(X, \tilde{T})$.

Let $\tilde{V}$ be a fuzzy open set in $(Y, \tilde{T}')$ containing $f(\tilde{p})$, then there exists a fuzzy open set $\tilde{U}$ in $(X, \tilde{T})$ containing $\tilde{p}$, such that $f(\text{cl}(\tilde{U})) \subseteq \tilde{V}$, Now $f(\text{int}(\text{cl}(\tilde{U}))) \subseteq f(\text{cl}(\tilde{U})) \subseteq \tilde{V}$ but, $\tilde{V} \subseteq \text{cl}(\tilde{V})$,

$$\tilde{V} = \text{int}(\tilde{V}) \subseteq \text{int}(\text{cl}(\tilde{V})). \text{ So, } f(\text{int}(\text{cl}(\tilde{U}))) \subseteq \text{int}(\text{cl}(\tilde{V}))$$

Which means that f is a fuzzy δ -continuous function.

Theorem 6.2. Every fuzzy strong θ -continuous function is a fuzzy θ -continuous function.

Proof:

Let $\tilde{p}$ be a fuzzy point in $(X, \tilde{T})$ and let a fuzzy open q-nbd. $\tilde{V}$ of $f(\tilde{p})$.

We must prove, there exists a fuzzy open q-nbd. $\tilde{U}$ of $\tilde{p}$ such that $f(\text{cl}(\tilde{U})) \subseteq \text{cl}(\tilde{V})$. Since f is

a fuzzy strong θ -continuous function. Therefore there exists a fuzzy open q-nbd. $\tilde{U}$ of $\tilde{p}$.

$$\rightarrow f(\text{cl}(\tilde{U})) \subseteq \tilde{V}. \text{ Now: } \tilde{V} \subseteq \text{cl}(\tilde{V}) \Rightarrow f(\text{cl}(\tilde{U})) \subseteq \tilde{V} \subseteq \text{cl}(\tilde{V}) \Rightarrow f(\text{cl}(\tilde{U})) \subseteq \text{cl}(\tilde{V}).$$

Hence, f is a fuzzy θ -continuous function.

Theorem 6.3.[6] Every fuzzy δ -continuous function is fuzzy θ -continuous function .

References

- [1] Allam, A. A., and Zahran, A. A., "On Fuzzy δ - Continuity and α - near Compactness in Fuzzy Topological Spaces", Vol.50, PP.103-112, 1992.
- [2] Azad, K. K., "On Fuzzy Semicontinuity, Fuzzy Almost continuity and Fuzzy Weakly continuity", J. Math. Anal. Appl., Vol.82, PP.14-32, 1981.
- [3] Chang, C. L., "Fuzzy Topological space", J. Math. anal. And Appl., Vol.24, PP.182-190, 1968.
- [4] Ming, P. P. and Mong, L. Y., "Fuzzy Topology I. Neighborhood Structure of a Fuzzy point and Moor-smith Convergence", J. Math. Anal. and Appl., Vol.76, PP.571-599, 1980.
- [5] Mukherjee, M. N. and Ghosh, "Some Stronger forms of Fuzzy Continuous Mapping on Fuzzy Topology Spaces: J. Fuzzy Sets and System, Vol.38, PP375-387, 1990.
- [6] Mukherjee, M. N. and Sinha, S. P., "On Some Near – Fuzzy Continuous Function Between Fuzzy Topological Spaces", J. Math. Anal. and Appl., Vol.34, PP. 245-254, 1990.
- [7] Othman, H. A. and Latha, S., "New Results of Fuzzy Alpha-Open Sets Fuzzy Alpha-Continuous Mappings", J. Contemp. Math. Sciences, Vol.4, No.29, PP.1415-1422, 2009.
- [8] Park, J. H., Lee, B. Y. and Choi, J. R., "Fuzzy θ -Connectedness", J. Fuzzy Set and Systems, Vol.59, PP.237-244, 1993.
- [9] Ray, A.D. and Chettri, P., "On Fuzzy Nearly Compact Regular Open Topology", Vol.4, PP.59-68, 2009.
- [10] S. Ganguly and S. Saha , A not on δ - continuity and δ -connected sets in fuzzy set theory Simon Stevin, Vol. 62, PP,127-141, 1988.
- [11] Saha, S., "Fuzzy δ -Continuous Mappings", J. Math. and Appl., Vol.126, PP.130-142, 1987.
- [12] Zadeh, L., "Fuzzy Sets", Inform. Control, Vol.8, PP.338-353, 1965.