



Tikrit Journal of Pure Science

ISSN: 1813 – 1662 (Print) --- E-ISSN: 2415 – 1726 (Online)

Journal Homepage: <http://tjps.tu.edu.iq/index.php/j>



SOME RESULTS ON STRONGLY π -REGULAR RING

Sinan O. Al-Salihi, Emad I. Jassim

Department of Mathematics, College of Education for pure science, Tikrit University, Tikrit, Iraq

ARTICLE INFO.

Article history:

-Received: 14 / 6 / 2022

-Accepted: 16 / 7 / 2022

-Available online: 26 / 4 / 2023

Keywords: regular ring , π -regular ring , strongly π -regular ring

Corresponding Author:

Name: Emad I. Jassim

E-mail: emad.i.jassim@st.tu.edu.iq

drsinan2001@gmail.com

Tel:

©2022 COLLEGE OF SCIENCE, TIKRIT UNIVERSITY. THIS IS AN OPEN ACCESS ARTICLE UNDER THE CC BY LICENSE

<http://creativecommons.org/licenses/by/4.0/>



ABSTRACT

In this paper we study the strongly π -regular ring (for short st. π -reg. rg.) and some properties also give some new results of st. π -reg. rg. and its connection with other rings.

1. Introduction

Let R be a ring we study concept of st. π -reg. ring., which introduced 1954 by Azumaya [2], and we give background theorems and corollary which we need in this paper also give some new results of st. π -reg. rg. and its connection with other rg.s. An element $a \in R$ is called regular element if there exists some $b \in R$ such that $aba = a$. A ring is called regular ring if every element is regular.

2. Strongly π -reg. ring.

Definition 2.1 [2]:

we call a st. π -regular if it is both right π -regular and left π -regular.

Now it can readily be seen that a power a^n of a is right (or left) reg. iff here exists an element b of s.t. $a^{n+1}b = a^n$ (or $ba^{n+1} = a^n$), where $a, b \in R$.

Theorem 2.2 [2]:

Under the assumption that R is of bounded index, the following four conditions are equivalent to $\forall$ other:

- (1) R π -reg.,
- (2) R is right π -reg.,
- (3) R is left π -reg.,
- (4) R is st. π -reg..

Lemma 2.3 :

Let b, c satisfy $a^{n+1}b = a^n, ca^{m+1} = a^m$ for some $n, m \in \mathbb{Z}$. Then they satisfy $a^{m+1}b = a^m, ca^{n+1} = a^n$ too.

Proof: When $m \geq n$ $a^{m+1} = a^m$ follows immediately from $a^{n+1}b = a^n$. Suppose now $m < n$. Then $a^m = ca^{m+1}$ implies $a^m (= c^2a^{m+2} = \dots) = c^{n-m}a^n$, and so we obtain $a^{m+1}b = c^{n-m}a^{n+1}b = c^{n-m}a^n = a^m$.

Similarly, we can verify the validity of $ca^{n+1} = a^n$.

■

Proposition 2.4[2] :

Every St. π -reg. element is π -reg..

Proposition 2.5 [5]:

Let R be a st. π -reg. ring. Then for all $a \in R$, there exists a positive integer n s.t. $a^n = eu = ue$ for some $e \in Id(R)$ and some $u \in U(R)$, where $Id(R)$ and $U(R)$ denote the set of idempotent of R and the set of units of R , respectively.

Definition 2.6 [8]:

A central idempotent in A is an idempotent in the central of A .

Theorem 2.7 [5]:

Let $\mathcal{R}$ be a rg. with central idempotent. Then $\mathcal{R}$ is st. π -reg. iff $N(\mathcal{R}) = J(\mathcal{R})$ and $\mathcal{R}/N(\mathcal{R})$ is reg., where $N(\mathcal{R}), J(\mathcal{R})$ denoted the set of all nilpotent and the Jacobson of $\mathcal{R}$ respectively.

Definition 2.8 [1]:

A ring $\mathcal{R}$ is called an exchange ring if for every $a \in \mathcal{R}$, there exists $e \in Id(\mathcal{R})$ such that $e \in a\mathcal{R}$ and $1 - e \in (1 - a)\mathcal{R}$. ($Id(\mathcal{R})$ means the set of all idempotent in $\mathcal{R}$).

Remark 2.9 [1]:

Every st. π -reg. rg. is an exchange rg..

Theorem 2.10 [1]:

Let $\mathcal{R}$ be an exchange ring and let a be a reg. element of $\mathcal{R}$ is st. π -reg., then a is a unit-reg element of $\mathcal{A}$.

Definition 2.11 [4]:

Let I be an ideal of a ring $\mathcal{R}$. We say that I is a st. π -reg. ideal of $\mathcal{R}$ in case for any $a \in I$ if there exist $n \in \mathbb{N}$ and $b \in I$ s.t. $a^n = a^{n+1}b$.

Theorem 2.12 [4]:

Let I be an ideal of a rg. $\mathcal{R}$. Then the following are equivalent:

- (1) I is st. π -reg..
- (2) Every element in I is st. π -reg. element.

Proposition 2.13 [6]:

Every right (or left) π -reg. rg. $\mathcal{R}$ is st. π -reg..

Remark 2.14:

The factor ring of the integers with respect to the ideal generated by the integer 4 is a st. π -reg. rg. which is not a reg. rg..

Theorem 2.15 [7]:

Let $\mathcal{R}$ be a rg. and I an ideal of $\mathcal{R}$.

(1) If $\mathcal{R}$ is a st. π -reg. rg. then so is $\mathcal{R}/I$ is st. π -regular ring.

(2) Assume that I is a reg. ideal of $\mathcal{R}$. Then, $\mathcal{R}$ is a st. π -reg. rg. Iff so is $\mathcal{R}/I$.

Proposition 2.16 [7]:

Let $\mathcal{R}$ be a rg. and P be a prime ideal of $\mathcal{R}$. If $\mathcal{R}/P$ is st. π -reg., then so is $\mathcal{R}_P$.

Definition 2.17:

Let $\mathcal{R}$ be a rg. and let $a \in \mathcal{R}$, the element a is called w-idempotent if for some positive integer n , a^n is an idempotent, i.e. $(a^n)^2 = a^n$.

Remark: The property that a is an w-idempotent is equivalent to the property that $\exists$ distinct positive integer n, m s.t. $a^n = a^m$.

On the other hand if there exists positive integer n, m with $n > m$ with $a^n = a^m$. Then there is some $t > 0$ s.t. $t(n - m) > m$.

Let $k = t(n - m) = m$ and let $f = a^{m+k} = a^{t(n-m)}$ then

$$a^m = a^n = a^m \cdot a^n a^{-m} = a^m a^{t(n-m)}$$

Thus

$$f = a^{t(n-m)} = a^{m+k} = a^k \cdot a^m = a^k a^m a^{t(n-m)} = a^k a^m a^{k+m} = f^2$$

$\therefore a$ is w-idempotent.

Theorem 2.18:

Let a be a st. reg. element of a ring $\mathcal{R}$. There exists one and only one element c s.t. $ac = ca$, $a^2c = (ca^2) = a$ and $ac^2 (= c^2a) = c$, and in particular a is reg. element. For any element b s.t. $a^2b = a$, c coincides with ab^2 . Moreover, c commutes with every element which is commutative with a .

Proof: Let b, d be two elements s.t. $a^2b = a, da^2 = a$. Then

$$(1) \quad ab = ba^2b = da,$$

So that

$$(2) \quad ab^2 = dab = d^2a.$$

From (1) we have also

$$(3) \quad aba = da^2 = a = a^2b = ada.$$

Now put $c = ab^2$. It follows then from (1), (2), (3), that

$$ac = adab = ab = da \quad daba = ca, \quad a^2c = aca = aba = a,$$

$$ac^2 = dac = dab = c, \text{ as desired.}$$

Suppose next c' be any element which satisfies the same equalities as c : $ac' = c'a, a^2c' = a, a^2c' = c'$. Then, be replacing b, d in (2) by c, c' respectively, we get $c = ac^2 = c'^2a = c'$, showing the uniqueness of c .

For the proof of the last assertion, let z be any element s.t. $az = za$. Then we have first $caz = cza = cza^2c = ca^2zc = azc = zac$, i.e., z commutes with $ca = ac$. It follows from this now $cz = c^2az = czca = cazc = zcac = zc$, and this completes the proof. ■

corollary 2.19 [2]:

Let a be a st. π -reg. element of $\mathcal{A}$. Suppose that a^n is right reg.. Then a^n is in fact st. reg., and moreover there exists an element c s.t. $ac = ca$ and $a^{n+1}c = a^n$.

Corollary 2.20 [3]:

Let $\mathcal{R}$ be a st. π -reg. rg. and $s \in \mathcal{R}$. Then $\exists n \geq 1$ and $a \in \mathcal{R}$ s.t. $s^n = s^{2n}a$, $sa = as$ and $a^2s^n = a$.

Theorem 2.21:

Let $\mathcal{R}$ be a rg. and $\{S_i\}_{i \in I}$ a collection of st. π -reg. subrg.s. Then $\bigcap_{i \in I} S_i$ is st. π -reg..

Proof: Let $s \in S = \bigcup_{i \in I} S_i$. Using one of the S_i we can find $n \geq 1$ and $a \in S_j$ s.t.

$s^n = s^{2n}a, sa = as$ and $a^2s^n = a$. Now consider S_j . For some $m \geq 1$ and $b \in S_j$ there is a solution for $s^{nm} = s^{2mn}b, s^{nm}b = bs^{nm}, b^2s^{nm} = b$. Further $s^{nm} = s^{2nm}a^m, s^{nm}a = as^{nm}$ and $a^{2m}s^{nm} = a^m$.

By corollary 2.26, $b = a^m \in S_j$. From $a = a^2s^n$ it follows that $a = a^m s^{(m-1)n} \in S_j$ if $m \geq 1$. If $m = 1$, $b = -a$ already. In any case $a \in S_j$. ■

Lemma 2.22 [2]:

Let a be a st. π -reg. element of index n , and c an element s.t. $ac = ca$ and $a^{n+1}c = a^n$ (as in corollary 2.19). Then $a - a^2c$ is a nilpotent element of index n .

We now obtain from corollary (2.19) and lemma (2.22), immediately the following.

Theorem 2.23:

Let $\mathcal{R}$ be a ring and let $a \in \mathcal{R}$ be a st. π -reg. element. Then there exists elements $u \in \mathcal{R}$ and $h \in \mathcal{R}$ s.t.

1. u is invertible. 2. $uh = hu = a$ 3. h is w -idempotent.

Proof: $\leftarrow$ By corollary (2.19), $\exists c \in \mathcal{R}$ and $n \in \mathcal{R}$ s.t. $a^{n+1}c = a^n$ and $ca = ac$. Then we have $a^n = a^{n+1}c = a^{n+2}a^2 = \dots = a^{2n}c^n = a^n c^n a^n$.

Let $w = a^n c^n$.

Then $w^2 = w$ and the elements a, c and w commute with $\forall$ other.

We also have $acw = ac(a^n c^n) = (a^{n+1}c)c^n = a^n c^n = w$

and $a^n w = a^n c^n a^n = a^n$.

Let $u = aw + (1 - w)$

and $h = w + a(1 - w)$ then $uh = hu$.

And $uh = [aw + (1 - w)][w + a(1 - w)] = aw^2 + a(1 - w)^2 = aw + a - aw = a$.

Also $h^n = [w + a(1 - w)]^n = w^n + a^n(1 - w)^n = w + a^n(1 - w) = w + a^n - a^n w = w$.

Thus g is an w -idempotent.

Finally ; let $z = [cw + (1 - w)]$ then $zu = uz$ and $uz = [aw + (1 - w)][cw + (1 - w)] = acw^2 + (1 - w)^2 = w + (1 - w) = 1$.

Therefore u is invertible. ■

Corollary 2.24 :

Let $\mathcal{R}$ be a st. π -reg. ring and let $a \in \mathcal{R}$, then $\exists$ elements $u \in \mathcal{R}$ and $h \in \mathcal{R}$ s.t. 1. u is invertible. 2. $uh = hu = x$. 3. h is an w -idempotent. Moreover, if A is a rg. s.t. for every element $a \in A$ $\exists$ elements $u \in A$ and $h \in A$ satisfying conditions (1),(2) and (3), then A is st. π -reg..

Proof: $\leftarrow$ The first assertion directly from theorem (2.23).

The second assertion s.t. let $a \in A$ and there exists elements $u \in A$ and $h \in A$ satisfying condition (1) , (2) and (3) for integer $n > 0$ s.t.

$$h^{2n} = h^n. \text{ Then } a^n = u^n h^n = u^{2n} u^{-n} h^{2n} = a^{2n} u^{-n} = a^{n+1} (a^{n-1} u^{-n})$$

And thus S is st. π -reg..

Remark 2.25:

We list have other useful relations of the elements use in the proof of theorem (2.23).

Let a is st. π -reg. elements and a rg. $\mathcal{R}$, and let $n \in N$ and a, c and w in $\mathcal{R}$ be the same in the proof of Theorem 2.23.

Thus we have $a^{n+1}c = a^n$, $ac = ca$

$$w = a^n c^n$$

$$acw = w,$$

$$a^n w = w$$

and a, c and w commute with $\forall$ other

$$\text{set } u = aw + (1 - w)$$

$$v = aw - (1 - w)$$

Then u and v and invertible with inverse

$$u^{-1} = cw + (1 - w)$$

$$v^{-1} = cw - (1 - w)$$

Finally , $a(1 - w)$ is nilpotent with $(a(1 - w))^n = 0$ It is st. π -reg.. ■

It is clearly consequence of corollary (2.24), is another proof of the result that $J(\mathcal{R})$, the Jacobson radical of $\mathcal{R}$ is nil when $\mathcal{R}$ is st. π -reg.. Since 0 is the only idempotent in $J(\mathcal{R})$, nilpotent elements only

w -idempotent in $J(\mathcal{R})$.

$\leftarrow$ If $a \in J(\mathcal{R})$ and h is an w -idempotent in the decomposition of a , then h is also in $J(\mathcal{R})$. Hence h (and hence a) is nilpotent.

In the following we will present the very important theorem.

Theorem 2.26 :

Let $\mathcal{R}$ be a st. π -reg. rg. if 2 is a unit in $\mathcal{R}$, then for all element of $\mathcal{R}$ can be expressed as a sum of two units.

Proof: Suppose $a \in \mathcal{R}$. Then as in the proof of Theorem 2.23, $\exists c \in \mathcal{R}$ and $n \in N$ s.t. $ac = ca$ and $a^{n+1}c = a^n$.

Let elements w, u, v, u^{-1} and v^{-1} in $\mathcal{R}$ be define as in remarks following colloary 2.24, Since v commutes with

$a(1 - w)$, we have that $2^{-1}v + a(1 - w)$ is a unit.

$$\text{Thus } 2^{-1}u + [2^{-1}v + a(1 - w)] =$$

$$2^{-1}(aw + (1 - w)) + 2^{-1}[aw - (1 - w)] +$$

$$a(1 - w) = aw + a(1 - w) = a.$$

Hence a is the sum of two units. ■

Now, Let $\mathcal{R}$ be a rg., and let $U(\mathcal{R})$ denoted the subrg. of $\mathcal{R}$ generated by the units of $\mathcal{R}$.

Thus , Theorem 2.24, shows that if $\mathcal{R}$ is st. π -reg.. And 2 is a unit of $\mathcal{R}$, then $U(\mathcal{R}) = \mathcal{R}$.

Proposition 2.27:

Let $\mathcal{R}$ be a st. π -reg. rg. and let A be a subrg. of $\mathcal{R}$.

If $U(\mathcal{R}) \leq A$, then A is st. π -reg..

Proof: $\leftarrow$ Let $a \in A$. Thus $a = uh$, where $u \in \mathcal{R}$, $h \in \mathcal{R}$, u is invertible, $uh = hu$, and h^n is idempotent for some integer $n > 0$. Then $u \in A$, $u^{-1} \in A$ and so $h = u^{-1}a \in A$. Thus the factorization in $\mathcal{R}$ is also a factorization in A , and so by corollary (2.24), A is st. π -reg.. ■

Proposition 2.28:

Every element a in a st. π -reg. rg. $\mathcal{R}$ is unit st. π -reg.; i.e. a has a generalized invers of st. π -reg. which is invertible.

Proof: Let $a \in \mathcal{R}$. Then as in the proof of theorem (2.23), $\exists c \in \mathcal{R}$ and $n \in N$ s.t. $ac = ca$ and $a^{n+1} = a^n$ let elements w and u in $\mathcal{R}$ be defined an in the remarks following corollary (2.24).

$$\text{Then } a^{n+1}u^{-1} = a^{n+1}[cw + (1 - w)]$$

$$= a^n acw + a^{n+1} - aa^n w$$

$$= a^n w + a^{n+1} - a^{n+1}$$

$$= a^n. \quad \blacksquare$$

Definition 2.29 [5]:

A ring $\mathcal{R}$ is said to be reduced if it has no nonzero nilpotent elements.

Proposition 2.30:

Let $\mathcal{R}$ be a reduced st. π -reg. rg.. Then $\mathcal{R}$ is a reg. rg..

Proof : Let $a \in \mathcal{R}$. Then as in the proof of Theorem 2.23, $\exists c \in \mathcal{R}$ and $n \in \mathcal{R}$ s.t. $ac = ca$ and $a^{n+1} = a^n$.

Let $w = a^n c^n$, then as in the remark following corollary 2.24 ,

$a(1 - w)$ is nilpotent.

$$\text{Hence } a = a^{n+1}c^n = a(a^{n-1}c^n)a. \quad \blacksquare$$

References

- [1] Ara, P. "Strongly π -regular rings have stable range one." Proceedings of the American Mathematical Society, vol. 124, no. 11, 1996.
- [2] Azumaya, G. Strongly π -regular rings Journal of the Faculty of Science Hokkaido University. Ser. 1 Mathematics, vol. 13, no. 1, 1954.
- [3] Burgess, W. D., and Menal, P. "On Strongly π -regular rings and homomorphisms into them." Communications in Algebra, vol. 16, no. 8, 1988.
- [4] Chen, H., and Chen, M. "On Strongly π -regular ideals." Journal of Pure and Applied Algebra, vol. 195, no. 1, 2005.
- [5] Chin, A. "A note on Strongly π -regular rings" Acta Mathematica Hungarica, vol. 102, no. 4, 2004.
- [6] El Khalfaoui, R., and Mahdou, N. "On stable range one property and Strongly π -regular rings" Afrika Matematika, vol. 31, no. 5, 2020.
- [7] Hirano, Y. "Some studies on Strongly π -regular rings" Mathematical Journal of Okayama University, vol. 20, no. 2, 1978.
- [8] Pawloski, R. M. Computing the cohomology ring and Ext-algebra of group algebras. The University of Arizona, 2006.

بعض النتائج حول الحلقات المنتظمة القوية من نمط π

عماد ابراهيم جاسم ، سنان عمر الصالحي

قسم الرياضيات ، كلية التربية للعلوم الصرفة ، جامعة تكريت ، تكريت ، العراق

الملخص

في هذا البحث قمنا بدراسة الحلقة المنتظمة القوية من نمط π - وبعض الخصائص التي تعطي ايضا بعض النتائج الجديدة حول الحلقة المنتظمة القوية من نمط π - وارتباطه بحلقات اخرى .