

Emergence of Extended spectrum β - Lactamases (ES β L) in clinical Isolates of *Klebsiella Pneumoniae* in Baghdad.

Nabaa A. Jasim, Riad A. Delool.

Emergence of Extended spectrum β - Lactamases (ES β L) in clinical Isolates of *Klebsiella Pneumoniae* in Baghdad.

Nabaa A. Jasim, Riad A. Delool.

Department of Biology, university of Almustansiriyah, college of science.

Received 8 April 2012 ; Accepted 13 May 2012

Abstract

The spread of Gram- negative bacteria with plasmid-born Extendend- Spectrum β - Lactamases (ES β Ls) had become aworldwide problem, that enzymes confer resistance to penicillin,cephalosporin the first, second and third generations and aztreonam via hydrolysis of the antibiotic ,that sample collected from urine and sputum sample .five isolates from forty-four of *Klebsiella pneumoniae* was production ES β L (11.36%).

Tested for susceptibility using Disk diffusion method ,and determination of (MIC)by standard agar dilution method ,and phenotypic detection of ES β L by using disc approximation method.

Keyword: *klebsiella pneumoniae*, Extended spectrum β -Lactamases, disc approximation method, minimum inhibitory concentration.

إنتشار إنزيمات البييتالاكتاميز الواسعة الطيف في عزلات بكتريا *Klebsiella*

Pneumoniae السريرية في بغداد.

نبأ علي جاسم ، رياض عبد الحسين دلول

كلية العلوم ،الجامعة المستنصرية ،قسم علوم الحياة

الخلاصة

ان انتشار انزيمات البييتالاكتاميز الواسعة الطيف المشفرة بلازميديا في البكتريا السالبة لصبغة غرام اصبحت مشكلة عالمية. هذه الانزيمات تكون مقاومة للبنسلينات وسيفالوسبورينات الجيل الاول والثاني والثالث والازتريونام عن طريق تحليلها لهذه المضادات الحيوية. العينات جمعت من الادرار والقشع ، خمس عينات موجبة من بكتريا *Klebsiella Pneumoniae* كانت منتجة لانزيمات ES β L (11.36%) من مجموع العينات البالغة اربعة واربعين عينة. تم اجراء فحص الحساسية وتحديد التركيز المثبط الادنى وتشخيص انتاجها لانزيمات ES β L مظهرها باستخدام طريقة الاقراص المتاخمة.

الكلمات الدالة :- الكلبسيلا الرئوية، انزيمات البييتالاكتاميز الواسعة الطيف، طريقة الاقراص المتاخمة، التركيز المثبط الادنى.

Introduction

The spread of antibiotic resistant bacteria has become alarg problem in most countries including Iraq. This bacteria is usually due to the production of extended spectrum β - Lactamase enzyme (1).

The excessive use of the oxyimino- cephalosporins in clinical practice had Resulted in *Klebsiella* species and other member of the Enterobacteriaceae showed diaminished susceptibility (2). *Klebsiella pneumoniae* is anormalflora bacteria, become opportunistic when transfer to urinary tract. its conceder the second cause of Nosocomial infection, (3). it

Emergence of Extended spectrum β - Lactamases (ES β L) in clinical Isolates of *Klebsiella Pneumoniae* in Baghdad.

Nabaa A. Jasim, Riad A. Delool.

showing increase of resistance when it found with another bacteria it showed co-resistance to (Aminoglycoside, Quinolones). That resistance increas with produce (ES β L) enzyme (4).

This study performed to investigate by phenotypic detection of ES β L in *Klebsiella pneumonia* isolated from patients in hospitals in Baghdad.

Material and method

Bacterial isolates:

One thousand and five hundred (1500) sample of urin and one hundred (100) sample from sputum were collected from Baghdad hospitals .

Diagnosis Isolates:

Forty- four isolates of *Klebsiella pneumoniae* were collected, That isolates diagnosis by morphological diagnosis, Biochemical tests and using Api 20Esystem.

The positive sample was culture in blood Agar, MacConkey Agar, Eosin methylen blue Agar and were stored in medium containing 15% glycerol in brain heart infusion broth and stored in deep freezing at -20C°.

Antimicrobial Susceptibility Testing

Organisms were tested by disc diffusion method using Muller-Hinton agar as described by(5).

Phenotypic Detection of ES β L

Detection of ES β L in positive sample was ascertained using disc approximation method (6). Using Mulller- Hinton agar and three disc containing 30 μ g for cefotaxime , ceftriaxone were Placed 30 mm (center to center) around a disc containing amoxicillin plus clavulanic acid (Augmentin) an increase in the zone diameter of Augmentin compared with other antibiotic considered positive for ES β LS.

Results

All sample collcted from urin and sputum (1600) were analysed for *K.pneumoniae* by identication according to morphological and biochemical tests .which showed (44) positive sample of *K.pneumoniae* Distribution according to clinical source in Table (I). This sample showed high resistance to antibiotic in this study, and was showed high MIC Value to antibiotic , the result showed five positive sample (11.36%) from 44 samples of *K.pneumoniae* have ability to producing ES β L enzyme.

Emergence of Extended spectrum β - Lactamases (ES β L) in clinical Isolates of *Klebsiella Pneumoniae* in Baghdad.

Nabaa A. Jasim, Riad A. Delool.

Table 1 :distribution of *K.pneumoniae* according to clinical source.

source	Total sample	Isolate	
		NO	%
Urine	1500	25	1.6
sputum	100	19	19

Discussion

The phenotypic Identification of producing ES β L enzyme can do by tow steps. The first is antimicrobial susceptibility test and the second steps is synergy between oxyimino cephalosporine and clavulanate (7). Disk approximation Method used to phenotypic method, the result showed 5 isolat (11. 36)% from 44 isolat of *K.pneumoniae* producing ES β L . is show in figure (1).




Figure (1):ESBL detection by using disc approximation method.

CTX:Augmentin CRO:Ceftriaxone CTX:Cefotaxime

Our result agree with Yan (8) , and disagree with Lal et al (2007) (9).our study showed high resistance to antibiotic, 100% to (ceftazidime, cefotaxime , ceftriaxone Aztreonam, carbencillin, piperacillin, Ampicillin), 80% to cefepime and ciprofloxacin 40% to Augmantin and streptomycin, 60% to Kanamycin. 20% to Amikacin, 0% to meropenem and Imipenem as show in figure (2)

Emergence of Extended spectrum β - Lactamases (ES β L) in clinical Isolates of *Klebsiella Pneumoniae* in Baghdad.

Nabaa A. Jasim, Riad A. Delool.




Figure (2):Antibiotic resistance of ES β L produce *K.pneumoniae*

CTX: Cefotaxime

PRL: pipracillin

AMC: Amoxicillin /clavulanic acid

Py: Carbenicillin

K:kanamycin

AM:Ampicillin

CRO: ceftriaxone

CAZ: ceftazidime

FEP: Cefepime

AK: Amikacin

ATM: Azetreonam

IMP: Imipenem

MPM: Meropenem

S: Streptomycin

CIP: Ciprofloxacin

The Resistance to carbencillin, pipracillin may be produce by bacteria enzyme broking that antibiotic(10).

The resistance to ceftriaxone, ceftazidime, cefotaxime by producing ES β L enzyme (11) resistance to cefotaxime, ceftriaxone, ceftazidime. Agree with Amin *et al* (2009)(12). While resistance to ceftriaxone was disagree with Nasehi *et al* (2010)(13). The resistance to Imipenem & meropenem was agree with Lim *et al*. (2009)(14)

Our study determination the minimum inhibitory concentration for four β -lactam antibiotic (Imipenem, ceftazidime, Ceftriaxone, cefotaxim.) . as show in table:2.

**Emergence of Extended spectrum β - Lactamases (ES β L) in
clinical Isolates of *Klebsiella Pneumoniae* in Baghdad.
Nabaa A. Jasim, Riad A. Delool.**

Table 2: Minimum Inhibitor concentration of ESBL –Producing *K. Pneumoniae*

Antibiotic NO. of Isolates	Imipenem	Ceftazidime	Ceftriaxone	cefotaxime
Break point $\mu\text{g/ml}$	≥ 16	≥ 32	≥ 64	≥ 64
K 22	0.05	128	512	512
K 26	0.05	32	256	128
K28	0.05	32	512	256
K 29	0.05	64	512	512
K36	0.1	265	1024	1024

The Imipenem MIC Value agree with nasehi *et al* (2010)(13).while ceftazidime MIC value Was disagree with GrÖber *et al* (2009)(15).and our study showed MIC to cefotaxime and cefriaxone disagree with Jones et al (2009) (16).

References

- (1) Dashti, A.A; jadaon,.M; Gomaa,. H.; Noronha, B.and udo, E. (2010). Transmission of *Klebsiella pnemoniae* clone harbouring genes for CTX-M-15 like and SHV-12 enzyme inaneonatal intensive care unite of Kuwaiti hospital. Journal of Medical Microbiology.
- (2) Heritage, J.; ZAli, F.H; Binzi, G and Hawkey, P.M (1999): Evolution and spread of SHV extended. Spectrum —Lactamses in gram- negative . bacteria . J Antimicrob chemother, 44
- (3) Struve, C and Krogfel, K.A. (2004). Pathogenic potential of environmental *Klebsiella pneumoiae* isolates. Environ microbial., 584-590.
- (4) Dwivedi, M.; Mishra., A. A.; Azim, A.; Singh, B., Baronina, K. A; Pasad, K.N; Dhole, T.N and Dwivedi, U.N. (2009). Ventilator – associated pneumoniae caused by carbapenem-resistant Enterobacteriaceae carrying multiple metallo- Lactamase gen. Indian Journal of Pathology. And microbiology. 52(3): 339-342.
- (5) CLSI ,(Clinical and Laboratory standards institute)(2007).performance standard for antimicrobial susceptibility testing :seventeenth informational supplement M100-S17 .Vol.27 No (1).
- (6) Collee,J.G;Fraser ,A.G;Marmion ,B.P and Simmons ,A. (1996).Mackie and McCartney Chrchill.livingston.USA
- (7) Rawat, D and Nair, D (2010). Extended spectrum -Lactamases gram. Negatve becteria . J glob infect Dis., (3) : 263-274.
- (8) Yan, J.J.; Ko, W.C. and Wu, J.J (2001). Identification of a plasmid Encoding SHV-12 , TEM-1, and Variant of IMP-2 metallo -B-Lactamases, IPM-8, from a clinical Isolate of *Klebsiella pneumoniae*. Antimicrob Agents chemother., 45(8): 2368-2371
- (9) Lal. P.; Kapil, A.; Das, B. K and Jood, S.(2007). Occurrence of TEM and SHV gene extended spectrum beta lactamase (ES β L) producing *Klebsiell* spp. Isolate from tertiary care hospital. Indian J med Re., 125:173-178
- (10) Pagani, G.; Radice, M.; Amoroso , A.; famiglieti, A., and gutkind, G (2006) Abstr. 44th . interscience conference on Antimicrobial Agent and chemother abst. C1-293 washington. D.C.

Emergence of Extended spectrum β - Lactamases (ES β L) in clinical Isolates of *Klebsiella Pneumoniae* in Baghdad.

Nabaa A. Jasim, Riad A. Delool.

- (11) Woodford, N.; ward, M.E.; kanfmann, M. E.; Fagan. J.; james, D.J; Johson , A.D. ;warner ,R and Livemove, D.M.(2006).molecular characterization of *E-coli* isolaltes producing CTX-M-15-extended spectrum lactmas in the united kingdon .Health protection Agency .,57:154-155.
- (12) Amin, A.; Ghumro,R. B.;Hussain ,s.and Hameed,A. (2009).prevalence of antibiotic resitance among clinical isolaltes of *klebsiella pneumomia* Isclated from tertiary care hospitalin in pakistan. Malaysia Journal of microbiology., 5(2): 81-86.
- (13) Lim, K. T.; Yeo, C.C.; Xasin, R.m.; Balan, G. and thong, K.L (2009). Charactenzation of Multidrag resistant and extended spectrum -Lactamases producing *Klebsiella Pneumoniae* strain from Malaysia hospital. Journal of medical Microbiolgy., 58:1463-1469.
- (14) Nashei, L. Shahcheraghi, F; Nikbin, V. S. and Nematzaden,s (2010). PER. CTX- M, TEM and SHV Beta. Lactamases in Clinical Isolates tahrn, Iran, Iranian Journal of Bisc Medical Sciences., 13(3): 111-118.
- (15) GrÖber, S.: Linke, D.; schutz, W.; fladerer, C. Madtunng, J and pfeifer., Y. (2009) . Emergence of carbapenem-non. Susceptible extended- spectrum . B-lactamase producing *klebsiella pneumoniae* isolated at the university hospital of Tübingen, Germany. Journal of Medical microbiology., 58: 912-922.
- (16) Jones, C. H.; Tuckman , M.; keeney , D.; Ruzin, A and Brad ford , P.A (2009). Characterization and sequence Analysis of ESL Encading genes from *Escherichia coil*, *Klebsiella pnemnomiae* isolated collected during tiygeclin phase 3 clinical trials. Antimicrob Agents chemother., 53 (2): 456-475.

