

The Role of Some CT Criteria in the Diagnosis of Solitary Pulmonary Cavitory Lesion

Mohammed Abd Kadhim *, Sundus Abd-Alrazzaq Salman**,
Husham Jubran Mousa***

ABSTRACT:

BACKGROUND:

The term cavity is a gas-filled space that appears on imaging as an area of low attenuation surrounded by a variously thickened wall within a pulmonary consolidation, mass, or nodule. Multidetector computed tomography (CT) of the chest is the current technique of choice for evaluating lung cavities.

OBJECTIVE:

To assess the role of some computed tomography criteria in differentiating benign from malignant solitary cavitory lung lesion.

PATIENT AND METHODS:

This was a cross sectional study done in the Computed Tomography unit of Al-Immamain Al-Kadhmain Medical city in Baghdad, Iraq between October, 2016 and June, 2017. All patients presented with solitary pulmonary cavitory lesions detected by chest x-ray referred for different reasons. Examination of the chest was performed by multi-detector CT (Somatom definition edge, SIEMENS medical system, Germany (256 slices)) with 2 sets of CT examination one before and another after giving IV nonionic iodinated contrast medium (Ultravist 370 mg /ml), 1.5 ml/kg Body weight. The final diagnosis was obtained depending on the sputum culture for AFB, bronchoscopy and biopsy, bronchoscopy and brush cytology and true cut biopsy.

RESULTS:

The study included 50 patients with solitary pulmonary cavitory lesions, 54% were male and 46% were females. Final diagnosis was 80% diagnosed as benign and 20% as malignant lesions. Enhancement was significant among 20% of studied patients. The common associated CT scan features were pleural effusion (24%), consolidation (12%), consolidation and tree in Bud (10%), pleural effusion and mediastinal lymph nodes (8%), tree in bud (10%), mediastinal lymph node and consolidation (8%), ground glass opacification, honeycombing and tree in bud (6%), pleural effusion and tree in bud (4%), fibrosis (2%) and encysted hydro-pneumothorax (2%). There was a highly significant association of significant enhancement and Pleural effusion with malignant solitary cavitory lesions ($p<0.001$ and $p=0.003$). A highly significant association was observed between increased mean thickness of lesion and malignant solitary cavitory lesion ($p<0.001$)

CONCLUSION:

The common computerized tomography characteristics of malignant solitary pulmonary cavitory lesions were significant enhancement, pleural effusion and increased diameter and thickness of pulmonary cavities.

KEYWORDS: solitary pulmonary cavity, CT scan.

INTRODUCTION:

The term cavity is a gas-filled space that appears on imaging examinations as an area of lucency or low attenuation surrounded by a variously thickened wall (greater than 4 mm) within a pulmonary consolidation, mass, or

nodule and usually result from the drainage of necrotic lesions via the bronchial tree ⁽¹⁾. Pulmonary cavities are one of the most common features of the lung and are often encountered in chest imaging examinations, and their differential diagnosis includes diverse malignant and non-malignant diseases ⁽²⁾. Various pathogenic mechanisms underlie the formation of cavitory lesions and include: inadequate local blood supply creating central necrosis, infarction from occlusion of regional nutritional vessels, and blockage of a bronchus resulting in necrosis distal to the obstruction ⁽³⁾. Many malignant tumors and benign diseases can develop or

*Professor in Medical Collage/ Al-Nahrain University, Consultant Radiologist in Al-Imamian Al-Kadhmyian Medical city/ Baghdad Iraq.

**Radiologist in Al-Imamian Al-Kadhmyian Medical City/ Baghdad Iraq.

***Specialist Radiologist in Al-Sader Teaching Hospital/ Basra Iraq.

CT CRITERIA IN THE DIAGNOSIS OF PULMONARY CAVITARY LESION

present with cavities, and it is important to differentiate a cavitory malignant tumor from a benign lesion ⁽⁴⁾. Up to 15% of primary lung cancers cavitate, mostly of squamous cell pathology ⁽⁵⁾. Although the cavitation in pulmonary metastases is not as common as in primary pulmonary cancer, cavitating pulmonary metastases have been reported with primary lung, colon, stomach, head and neck, urinary bladder, kidney, uterine cervix, bone, pancreas, and breast cancers ⁽³⁾.

Multidetector computed tomography (CT) of the chest is the current technique of choice for evaluating lung cavities, as it provides precise information on size, shape, location of lesions, and other characteristics that may not be evident on CXRs ^(6, 7). It is an important tool in the evaluation of mediastinal lymph nodes (LNs), and evaluating associated complications. It also can be used to differentiate the various etiologies and identifying pleural/ airway/ diaphragmatic pathologies and evaluating visualized bones. Contrast-enhanced CT (CECT) is the investigation of choice for evaluation of mediastinal LNs and also aids in depicting pleural enhancement in empyema ⁽⁸⁾.

AIM OF THE STUDY:

To assess the role of some computed tomography criteria in differentiating benign from malignant solitary cavitory lung lesion.

PATIENTS AND METHODS:

The Study was a cross sectional analytic study carried out in the Computed Tomography unit of Al-Immamain Al-Kadhmain Medical city in Baghdad, Iraq through the period from 1st October, 2016 to 30th June, 2017. All patients presented to Computed Tomography unit with solitary pulmonary cavitory lesions detected by chest x-ray referred from clinicians for different reasons. Research approval was taken from Institutional Review Board/ Al-Nahrain college of Medicine. Oral informed consent was taken from patients before enrolling in the study.

Inclusion criteria: adult patients with single cavitory lung lesion detected by chest x-ray regardless of symptoms. Exclusion criteria: multiple cavitory lung lesions, patients with known lung carcinoma on treatment, patients with known pulmonary tuberculosis on treatment and patients receiving chest radiotherapy.

Examination protocol: the examination of the chest was performed with multi-detector CT (Somatom definition edge, SIEMENS medical system, Germany (256 slices)). All the patients

were examined in the supine position in caudo-cranial direction during breath holding period of 5-10 sec, Slice thickness 5mm, KVp 100-120, mA 200-300. All patients had 2 sets of CT examination one before and another after giving IV nonionic iodinated contrast medium (Ultravist370 mg /ml), the dose of contrast medium in an adult patient is given as 1.5 ml/kg Body weight and the magnitude of enhancement was assessed by measuring CT number pre- and post-contrast administration and was considered significant if the increase in CT number is more than 15 HU. Images were viewed in 2 different window settings: lung window (WL= -600, WW= -1600), and mediastinum window (WL=40, WW=350). Sagittal and coronal reformatted images were obtained from initial axial CT data.

Statistical analysis: all patients' data entered using computerized statistical software; Statistical Package for Social Sciences (SPSS) version 22 was used. Descriptive statistics presented as (mean $\pm$ standard deviation) and frequencies as percentages. Chi square test was used for comparison between categorical data (Fishers exact test was used when expected variables were less than 20% of total). Independent sample t-test was used to compare between two means. In all statistical analysis, level of significance (p value) set at ≤ 0.05 .

RESULTS:

The study included 50 patients with x-ray diagnosis of solitary pulmonary cavitory lesions, 54% were male and 46% were females with male to female ratio as 1.1:1. The mean age of the studied group was 50.7 ± 11.9 years; 4% of them were in age group <30 years, 10% of them were in age group 30-39 years, 34% of them were in age group 40-49 years, 22% of them were in age group 50-59 years and 30% of them were in age group ≥ 60 years. The clinical presentations of patients with solitary pulmonary cavitory lesions were: productive cough (94%), fever (70%), weight loss (56%), sweating (20%), anorexia (16%), and dyspnea (16%).

The final diagnosis of the 50 patients was: 80% of patients with solitary cavitory lesions were diagnosed by CT scan as benign and 20% of them were diagnosed as malignant lesions. Regarding specific diagnosis, the most frequent diagnosis was TB (60%), followed by; squamous cell carcinoma (14%), lung abscess (10%), hydatid cyst (10%), adenocarcinoma (4%), and metastasis (2%) as shown in table 1.

CT CRITERIA IN THE DIAGNOSIS OF PULMONARY CAVITARY LESION

Table 1: The final diagnosis of patients with solitary pulmonary cavitory lesions.

Variable	No.	%
Diagnosis		
TB	30	60.0
Squamous cell carcinoma	7	14.0
Lung abscess	5	10.0
Hydatid cyst	5	10.0
Adenocarcinoma	2	4.0
Metastasis (adenoca.)	1	2.0
Total	50	100.0
Final diagnosis		
Benign	40	80.0
Malignant	10	20.0
Total	50	100.0

CT findings: The mean diameter of solitary cavitory lesions was 38.5 ± 11.7 mm; with range of 15 ± 58.5 mm. Mean thickness of solitary cavitory lesions was 8.6 ± 8.1 mm; with range of 3 ± 27.2 mm. The CT enhancement finding was significant among 20% of studied patients. The common associated CT scan features were pleural effusion (24%), consolidation (12%), consolidation and tree in Bud (10%), pleural effusion and mediastinal lymph nodes (8%), tree

in bud (10%), mediastinal lymph node and consolidation (8%), ground glass opacification, honeycombing and tree in bud (6%), pleural effusion and tree in bud (4%), fibrosis (2%) and encysted hydro-pneumothorax (2%). There was a highly significant association between significant enhancement and pleural effusion with malignant solitary cavitory lesions ($p < 0.001$ and 0.003 respectively) as shown in table 2.

CT CRITERIA IN THE DIAGNOSIS OF PULMONARY CAVITARY LESION

Table 2: The patients' CT scan characteristics distribution according to CT scan findings.

Variable	Benign		Malignant		Total		
	No.	%	No.	%	No.	%	
Enhancement							
Significant	1	10	9	90	10	20	0.001** Highly significant
Non significant	38	95	2	5	40	80	
Associated features							
Pleural effusion	9	22.5	3	30.0	12	24.0	0.003** Significant
Consolidation	3	7.5	3	30.0	6	12.0	Non Significant
None	7	17.5	0	-	7	14.0	Non Significant
Consolidation & tree in bud	5	12.5	0	-	5	10.0	Non Significant
Pleural effusion and mediastinal lymph nodes	0	-	4	40.0	4	8.0	Non Significant
Tree in bud	5	12.5	0	-	5	10.0	Non Significant
Mediastinal lymph nodes and consolidation	4	10.0	0	-	4	8.0	Non Significant
Ground glass, honeycombing and tree in bud	3	7.5	0	-	3	6.0	Non Significant
Pleural effusion and tree in bud	2	5.0	0	-	2	4.0	Non Significant
Fibrosis	1	2.5	0	-	1	2.0	Non Significant
Encysted hydro-pneumothorax	1	2.5	0	-	1	2.0	Non Significant

*Chi-square test; **Fishers exact test.

The mean diameter of malignant cavitory lesions was significantly higher than mean diameter of benign cavitory lesions ($p < 0.001$). A highly significant association was observed between increased mean thickness of lesion and malignant solitary cavitory lesion ($p < 0.001$) as shown in table 3.

Table 3: The mean diameter and thickness of benign and malignant lesions in CT.

Variable	Benign (mm)	Malignant (mm)	P
	Mean±SD	Mean±SD	
Diameter (mm)	35.03±10.12	52.4±5.5	<0.001* Highly significant
Thickness (mm)	5.3±4.9	21.5±4.4	<0.001* Highly significant

*Independence sample t-test.

DISCUSSION:

It's important in clinical and radiological field to distinguish a cavitory malignant lesion from a benign lesion ⁽⁹⁾. A continuous higher number of admissions to hospitals and health centers related with or mimicking tuberculosis has highlighted the issue of precise detection of tuberculosis and cancers. There are numerous interferences between both illnesses like they both are prevalent, have high incidence, include lung parenchyma or more all, described by closed presentations. Despite this, there are numerous dissimilarities between these two diseases like they have different etiology, diverse results, and treatment. Delay in the detection and treatment of lung malignancies lead to poorer result and lower survival ⁽¹⁰⁾.

The mean age of the studied patients with solitary pulmonary cavitory lesion was 50.7 years with high proportions of patients over 40 year's age. This finding is similar to that reported in a previous study in USA which stated that solitary pulmonary cavitory lesion were more prevalent among elderly patients and many radiology abnormalities especially among smokers were obvious after age of 50 years ⁽⁶⁾. Other studies in India and USA reported that malignant cavitory lesion had higher incidence with increase of patient's age ⁽¹¹⁾. The incidence of lung cancer increases with age, with 60% of patients being over the age of 65 ⁽¹²⁾.

In this study the common significant CT characteristics of malignant solitary pulmonary cavities were significant enhancement, and pleural effusion. These findings are consistent with the results of many previous studies like Sim et al ⁽¹³⁾ study in UK, Li et al ⁽⁹⁾ study in China and Ödev et al ⁽¹⁴⁵⁾ study in Turkey. Truong et al ⁽¹⁵⁾ documented that pleural effusion is more commonly found among patients with malignant solitary pulmonary nodules. CT scan contrast enhancement >15 HU is 99% predictive for benign lesions ⁽¹⁶⁾. Malignant tumors are generally relatively more hypervascular in comparison to benign lesions and the intensity of enhancement is directly related to the vascularity led to the conclusion that the stronger the enhancement the more malignant potential of the nodule ⁽¹⁷⁾. This enhancement cut off of 15HU resulted in an excellent sensitivity of 98% but unfortunately it was not very specific for malignancy, only about 50–60%. The general conclusion based on this report was that benign lesions usually enhance no more than 15 HU, whereas most of the malignant nodules develop more intensive enhancement, usually

over 20 HU ⁽¹⁸⁾. In a different study by Yi et al ⁽¹⁹⁾ with the enhancement cut-off of 30 HU, sensitivity for malignant nodules was 99% with a negative predictive value of 97%.

The appearance and thickness of a cavity wall can play useful role in diagnosis. Benign cavities tend to have smooth, thin walls, usually less than 4 mm at its broadest point, whereas nodules containing cavities with irregular, thick walls (exceeding 16 mm) have been found to be malignant in up to 95% of cases ⁽¹⁷⁾. The present study showed that means diameter and thickness of malignant cavitory lesion were significantly higher than benign cavitory lesions. This finding coincides with results of Figueroa et al ⁽²⁰⁾ study in USA and Nin et al ⁽²¹⁾ study in UK which reported that increase of diameter and thickness of solitary pulmonary cavities is significantly related to malignancy. Another study conducted in South Korea ⁽²²⁾ revealed that wall thickness is significantly differentiating between malignant and benign cavitory lung lesions. Inconsistently, previous Chinese study ⁽²³⁾ documented that cavitory wall thickness is unreliable for differentiating between malignant and benign lesions. Malignant pulmonary cavitation characterized by irregular or round appearance with discrepancy in wall thickness. The thickness above 24 mm in thickness with consolidation is predictive of malignancy ⁽²⁴⁾. The use of cavity wall thickness to discriminate among infectious etiologies of pulmonary cavities is even more problematic. While some infections, such as pneumocystis pneumonia, coccidioidomycosis, and Echinococcus, have been classically associated with thin walled cavities, the absence of comparative studies with systematic, objective measurements of cavity wall thickness among infectious etiologies severely limits the use of cavity wall thickness as a diagnostic tool in discriminating among infectious causes of cavities ⁽²⁵⁾.

CONCLUSION:

The common computerized tomography characteristics of malignant solitary pulmonary cavitory lesions were significant enhancement, pleural effusion and increased diameter and thickness of pulmonary cavities.

REFERENCES:

1. Kemal O, İbrahim G, Tamer A, et al. Cystic and Cavitory Lung Lesions in Children: Radiologic Findings with Pathologic Correlation. J Clin Imaging Sci. 2013; 3: 60.

2. Kim NR, Han J. Pathologic review of cystic and cavitory lung diseases. *Korean J Pathol* 2012; 46:407-14.
3. Vourtsi A, Gouliamos A, Mouloupoulos L, et al. CT appearance of solitary and multiple cystic and cavitory lung lesions. *Eur Radiol* 2001;11:612-22.
4. Honda O, Tsubamoto M, Inoue A, Johkoh T, Tomiyama N, Hamada S, Mihara N, Sumikawa H, Natsag J, Nakamura H. Pulmonary cavitory nodules on computed tomography: differentiation of malignancy and benignancy. *Journal of computer assisted tomography*. 2007 ;31:943-49
5. Truong MT, Sabloff BS, Ko JP: Multidetector CT of solitary pulmonary nodules. *Radiol Clin North Am*, 2010; 48: 141–55.
6. Gadkowski LB, Stout JE. Cavitory pulmonary disease. *Clin Microbiol Rev* 2008;21:305-33.
7. Ryu JH, Swensen SJ. Cystic and cavitory lung diseases: focal and diffuse. In *Mayo Clinic Proceedings* 2003; 78:744-52.
8. YUANGANG Q, QING Z, YONG H and DAOQING W. Manifestations and pathological features of solitary thin-walled cavity lung cancer observed by CT and PET/CT imaging. *Oncol Lett*. 2014 Jul; 8: 285–90.
9. Li BG, Ma DQ, Xian ZY, Guan J, Luo KJ, Fan QW, Peng L, He W. The value of multislice spiral CT features of cavitory walls in differentiating between peripheral lung cancer cavities and single pulmonary tuberculous thick-walled cavities. *The British journal of radiology*. 2012;85:147-52.
10. Bhatt ML, Kant S, Bhaskar R. Pulmonary tuberculosis as differential diagnosis of lung cancer. *South Asian journal of cancer*. 2012 ;1:36.
11. Jemal A, Siegel R, Xu J, Ward E. Cancer statistics, 2010. *CA: a cancer journal for clinicians*. 2010 ;60:277-300.
12. Marom EM, Martinez CH, Truong MT, Lei X, Sabloff BS, Munden RF, Gladish GW, Herbst RS, Morice RC, Stewart DJ, Jimenez CA. Tumor cavitation during therapy with antiangiogenesis agents in patients with lung cancer. *Journal of Thoracic Oncology*. 2008;3:351-57.
13. Sim YT, Poon FW. Imaging of solitary pulmonary nodule—a clinical review. *Quantitative imaging in medicine and surgery*. 2013;3:316.
14. Odev K, Guler I, Altinok T, et al. Cystic and cavitory lung lesions in children: radiologic findings with pathologic correlation. *Journal of clinical imaging science*. 2013;3:60.
15. Truong MT, Ko JP, Rossi SE, Rossi I, Viswanathan C, Bruzzi JF, Marom EM, Erasmus JJ. Update in the evaluation of the solitary pulmonary nodule. *Radiographics*. 2014;34:1658-79.
16. Zhang M, Kono M. Solitary pulmonary nodules: evaluation of blood flow patterns with dynamic CT. *Radiology*. 1997;205:471-78.
17. Choromańska A, Macura KJ. Evaluation of solitary pulmonary nodule detected. *Pol J Radiol*, 2012; 77: 22-34.
18. Swensen SJ, Viggiano RW, Midthun DE et al: Lung nodule enhancement at CT: multicenter study. *Radiology*, 2000; 214: 73–80.
19. Yi CA, Lee KS, Kim EA et al: Solitary pulmonary nodules: dynamic enhanced multi-detector row CT study and comparison with vascular endothelial growth factor and microvessel density. *Radiology*, 2004; 233: 191–99.
20. Figueroa, Elyn Riedel E and S. Glickman MS. Clinical and radiographic differentiation of lung nodules caused by mycobacteria and lung cancer: a case-control study *BMC Infection Disease* 2015;15:482.
21. Nin CS, de Souza VV, Alves GR, et al. Solitary lung cavities: CT findings in malignant and non-malignant disease. *Clinical radiology*. 2016;71:1132-36.
22. Park Y, Kim TS, Yi CA, et al. Pulmonary cavitory mass containing a mural nodule: differential diagnosis between intracavitory aspergilloma and cavitating lung cancer on contrast-enhanced computed tomography. *Clinical radiology*. 2007;62:227-32.
23. Tripathi S, Zhen X. Differentiation of Benign and Malignant Solitary Pulmonary Nodule: Literature Review. *Advances in Lung Cancer*. 2015;4:17.
24. Erasmus JJ, McAdams HP, Farrell MA, Patz Jr EF. Pulmonary nontuberculous mycobacterial infection: radiologic manifestations. *Radiographics*. 1999;19:1487-503.
25. Souza Jr AS, Gasparetto EL, Davaus T, Escuissato DL, Marchiori E. High-resolution CT findings of 77 patients with untreated pulmonary paracoccidioidomycosis. *American Journal of Roentgenology*. 2006 ;187:1248-52.