

Overweight and Obesity Among Children and Adolescents in Baladrouz City

Wathiq Hassan Farman (BSc)¹ and Hameed M Majeed (PhD)²

Abstract

Background: Obesity in children and adolescents is one of the most serious public health challenges of the 21st century because of the effects of future morbidity. Its prevalence has increased significantly in most countries, especially in low and middle-income countries, especially in urban areas.

Objective: To know the distribution of this epidemic among our children and draw a real picture of intermediate school students in Baladrouz region.

Patients and Methods: A cross-section study, involved the students of the first and the second stages, the information and the measurements were taken from 1798 students (910 males, 888 females) their ages were between 11-14 years, over one month duration (December of 2017), during this period we visit a were 10 Intermediate schools for both genders (six of them in the center of the region and four in the surrounding villages those belong to the region), included the age, weight, height, counting the body mass index, waist circumference, Neck circumference, systolic blood pressure and diastolic blood pressure in addition to the questionnaire form and the information of the school card.

Results: The percentages of the overweight and the obesity among males were as follows: 26.9%, 17.4% for intermediate, first and second stages respectively, while in females it was 47.9%, 33.3% respectively. The percentage of weight gain and central obesity of the waist circumference was 27.0%, 24.1% for the first and second stages respectively. Either in females, as follows: 21.5%, 12.8% for the first and second stages respectively. There was an increasing in the percentages of systolic blood pressure among female for all stages of the study, which amounted to 60.3%, 48.7% either in males was 70.4%, 64.8% for the first and second stages respectively. The percentage of normal diastolic blood pressure was 54.5%, 49.0% for males, while in females it was 72.5%, 65.7% for the first and second stages respectively.

Conclusion: The study showed high rates of overweight and obesity among middle school students, and was higher in females than males.

Keywords: Overweight, Obesity, Children of the Intermediate schools, Baladrouz.

Corresponding Author: wathk9298@gmail.com

Received: 8th November 2018

Accepted: 30th December 2018

¹ College of Education for Pure Science- University of Diyala -Diyala - Iraq.

² Directorate of Education Diyala- Diyala – Iraq.

Introduction

Obesity among children, adolescents and adults has emerged as one of the most serious public health concerns in the 21st century. The worldwide prevalence of childhood obesity has increased unusually over the past 3 decades. The growing prevalence of childhood obesity has also led to appearance of obesity-related comorbid disease entities at an early age. Childhood obesity can adversely affect nearly every organ system and often causes serious consequences, including hypertension, dyslipidemia, insulin resistance, dysglycemia, fatty liver disease and psychosocial complications. It is also a major contributor to increasing healthcare expenses. For all these reasons, it is important to prevent childhood obesity as well as to identify overweight and obese children at an early stage so they can begin treatment and attain and maintain a healthy weight[1]. Obesity is most often defined by the body mass index (BMI), a mathematical formula of weight-for-height index. BMI is measured by dividing the body weight in kilograms to height in meters squared (kg/m^2) [2,3,4]. In the pediatric age group, gender-specific BMI-for-age percentile curves are used to define overweight and obesity. Children and adolescents with a BMI over the 85th but less than the 95th percentile for age and gender are considered overweight and those with a BMI greater than the 95th percentile are considered obese. Children and adolescents with a BMI greater than the 99th percentile are considered severely obese [5,3,6]. High blood pressure has developed highly prevalent in children and adolescents

in recent years [7,8]. the notion of prehypertension has been brought to light for children and adolescents too. Elevated blood pressure at a young age is a predictor of hypertensive disease later in life [9,10] and it is related with cardiac and renal alterations as well as target organ damage [11,12]. Considering the potential future impact of uncontrolled high blood pressure in this segment of the population, excessive attention should be paid to establishing suitable prevention and treatment programs. Neck circumference (NC) has also been used as a potential proxy for obesity and cardiovascular disease in adults. Very few investigators¹¹ have attempted to use NC to screen for high BMI in children; therefore, the objectives of this study were to examine the correlation between NC and BMI in children, to examine the ability of NC to identify correctly children with high BMI, and to determine the best NC cut point for identifying children of various ages as overweight/obese [13].

Patients and Methods

The Study involved the students of the first and the second stages, within Baladrouz region-Diyala, the information and the measurements were taken from 1798 students (910 males, 888 females) their ages were between 11-14 years, in one month duration (December of 2017), during this period the visits were to 10 Intermediate schools for both genders (six of them in the center of the region and four in the surrounding villages those belong to the region), included the age, weight, height, counting the body mass

index, waist circumference, Neck circumference, systolic blood pressure and diastolic blood pressure in addition to the questionnaire form and the information of the school card.

Anthropometric measurements

The anthropometric measurements of the students were measured by tape measure tool for the height (the students stood on the ground and the height from head to toe was measured) and waist circumference measurements (below the last rib level or at the umbilicus level), balance apparatus for the body weight (kg) measurement by putting it on the ground and the student stood straightly on it and the body mass index of the students was counted for both genders (males and females) by using the following equation: [14] Body mass index = weight (kg) / height (m²).

Blood pressure measurement

The mercury sphygmomanometer was used for the systolic and diastolic blood pressure (mmHg) measurement by fitting the left arm to a solid stent and rolling the cuff about the

arm and inflated it to cover 70% of the arm, adjacent the elbow.

Statistical analysis

The results were statistically analyzed by (Independent - Samples T Test) analysis and application of SPSS program version 20. A P value < 0.05 was considered statistically significant.


Results

The present study measures the body mass index, waist circumference, classified according to Centers for Disease Control (CDC) and Prevention to underweight, natural, overweight and obesity and systolic and diastolic blood pressure classified according to The Fourth Report on the Diagnosis. The Table (1) shows mean anthropometric measurement values and systolic and diastolic blood pressure and The mean BMI, WC, NC in both males and females, The rise BMI, SBP, DBP in females was higher than that of males at significant level (P value < 0.05).

Table (1): Comparison of anthropometric measurements, systolic and diastolic blood pressure between male and female.

Measures	Stages \ Age (years)	Male		Female	
		Number	Mean ± SD	Number	Mean ± SD
BMI (kg/m ²)	First(11-12)	433	20.03±4.33	459	21.08±4.37
	Second(13-14)	477	20.85±4.89	429	21.86±4.04
WC (cm)	First(11-12)	433	71.27±11.92	459	70.47±4.99
	Second(13-14)	477	73.79±13.14	429	71.87±10.04
NC(cm)	First(11-12)	433	30.92±5.37	459	30.22±2.21
	Second(13-14)	477	31.54±3.66	429	30.57±2.17
SBP (mmHg)	First(11-12)	433	107.30±13.92	459	112.91±15.44
	Second(13-14)	477	109.24±14.84	429	117.69±14.74
DBP(mmHg)	First(11-12)	433	63.49±11.64	459	68.61±10.96
	Second(13-14)	477	64.34±11.27	429	71.11±11.00

*SD: Standard deviation, BMI: Body mass index, WC: Waist circumference, NC: Neck circumference, SBP: Systolic blood pressure, DBP: Diastolic blood pressure .



Figure(1): BMI for SBP & DBP in males.




Figure (2): BMI for SBP & DBP in females

The Table (2) refers to the distribution of the overweight and obesity according to the body mass index for the males of the first and the second stages which reached (12.4 % ,14.5%) , (4.1 % , 13.2 %) respectively

while the percentages of the overweight and obesity for the females of the first and the second stages reached (35.9% ,12.0 %) , (22.1 % ,11.1 %) respectively.

Table (2): The percentages of overweight and obesity distribution among the intermediate schools' students according to the body mass index measurement.

Gender	Stages/Age(years)	No.	U.w		Natural		O.w		Ob	
			Mean	%	Mean	%	Mean	%	Mean	%
Male	First (11-12)	433	14.60	7.22	18.00	65.81	23.33	12.43	28.56	14.54
	Second (13-14)	477	15.00	10.27	18.71	72.32	24.02	4.19	30.53	13.21
Female	First (11-12)	459	14.44	5.01	17.94	47.05	23.21	35.91	29.82	12.03
	Second (13-14)	429	14.83	1.87	19.44	64.81	25.07	22.14	30.68	11.18

*(%): Percentage, (No.): Number, (U.w.): Underweight, (O.w.): Overweight, (Ob.): Obesity

The Table (3) refers to the distribution of the overweight and the central obesity according to the waist circumference for the males of the first and second stages which reached (14.7 % , 9.4%) , (19.4 % , 7.5 %) respectively while the percentages

of the overweight and the central obesity for the females of the first and second stages reached (13.5 % , 8.0%) , (10.7 % , 2.1 %) respectively.

Table (3): The percentages of overweight and obesity distribution among the intermediate schools' students according to the waist circumference measurement.

Gender	Stages/Age(years)	No.	U.w		Natural		O.w		Ob	
			Mean	%	Mean	%	Mean	%	Mean	%
Male	First (11-12)	433	58.41	37.72	65.96	38.10	80.11	14.78	93.77	9.40
	Second (13-14)	477	58.27	30.42	68.01	42.55	83.08	19.49	98.09	7.54
Female	First (11-12)	459	59.13	35.72	63.66	42.70	74.81	13.51	89.42	8.06
	Second (13-14)	429	58.62	36.15	66.89	51.04	78.77	10.72	89.83	2.09

* (%): Percentage, (No.): Number, (U.w.): Underweight, (O.w.): Overweight, (Ob.): Obesity

The Table (4) refers to the distribution of the prehypertension and hypertension according to the systolic blood pressure for the males of the first and second stages which reached (18.4 % , 11.0%) , (21.1 % ,

13.3%) respectively while the percentages of the prehypertension and hypertension for the females of the first and second stages reached (20.0 % , 19.6%) , (23.3 % , 27.9 %) respectively.

Table (4): The percentages of prehypertension and hypertension distribution among the intermediate schools' students according to the systolic blood pressure measurement.

Gender	Stages/Age(years)	No.	Natural		Prehypertension		Hypertension	
			Mean	%	Mean	%	Mean	%
Male	First (11-12)	433	99.93	70.43	119.5	18.49	132.5	11.08
	Second (13-14)	477	101.3	64.88	120.8	21.74	127.5	13.38
Female	First (11-12)	459	102.8	60.34	121.2	20.12	135.1	19.62
	Second (13-14)	429	104.6	48.71	122.1	23.32	135.98	27.97

* (%): Percentage, (No.): Number

The Table (5) refers to the distribution of the prehypertension and hypertension according to the diastolic blood pressure for the males of the first and second stages which reached (39.1% , 6.3 %) , (33.9 % ,

16.9%) respectively while the percentages of the prehypertension and hypertension for the females of the first and second stages reached (24.4 % , 3.0 %) , (26.8 % , 7.4%) respectively.

Table (5): The percentages of prehypertension and hypertension distribution among the intermediate schools' students according to the diastolic blood pressure measurement.

Gender	Stages/Age(years)	No.	Natural		Prehypertension		Hypertension	
			Mean	%	Mean	%	Mean	%
Male	First (11-12)	433	54.67	54.51	72.95	39.15	90.05	6.34
	Second (13-14)	477	55.83	49.06	71.81	33.97	83.15	16.98
Female	First (11-12)	459	72.54	72.54	80.47	24.41	91.23	3.11
	Second (13-14)	429	65.73	65.73	81.12	26.85	91.85	7.43

* (%): Percentage, (No.): Number

Discussion

The obesity is the main determine to increase the blood pressure in the childhood, the high prevalence of the fat in the body is an pointer for the some complications those belong to the obesity such as hypertension. According to the body mass index, the percentage of the overweight and obesity for the males was higher in the first stage 26.9% followed by the second stage 17.4% while the percentage of the overweight and obesity for females was higher in the first stage 47.9% followed by the second stage 33.3%. The study of James et al 15 found that BMI is more associated with blood pressure than weight and height variables, as well as a good representative of some factors that are believed to have an effect on blood pressure, including diet. Several studies confirm a link between obesity and hypertension. The Flynn et al 16 study found that there is a threefold increased risk in obese children to develop hypertension than healthy children. According to the waist circumference, the percentage of the overweight and central obesity for the males was higher in the first stage 27.0% followed by the second stage 24.1 %, in females the percentage of the overweight and central obesity was higher in

the first stage 21.5% followed by the second stage 12.8%. Chen et al 17 showed a significant upward trend to increase blood pressure with increased waist circumference and BMI.

According to the systolic blood pressure, the percentage of the prehypertension and hypertension for the males was higher in the second stage 35.1 % followed by the first stage 29.5 % ,the percentage of the prehypertension and hypertension for the females was higher in the second stage 51.2 % followed by the first stage 39.6 %. According to the diastolic blood pressure, the percentage of the prehypertension and hypertension for the males was higher in the second stage 50.9 % followed by the first stage 45.4% while the percentage of the prehypertension and hypertension for the females was higher in the second stage 34.2 % followed by the first stage 27.4 % . study Hosseini et al 18, in 2015 showed that both the SBP and DBP rose steadily with increasing, for BMI. A positive relationship study also found increased waist circumference and increased systolic and diastolic blood pressure levels [19] . The results of the Kelishadi et al 20 study showed

that neck circumference was significantly associated with some cardiovascular risk factors, including hypertension, weight gain, general and central obesity.

Conclusions

The study showed high rates of overweight and obesity among middle school students, and was higher in females than males.

Recommendations

Health awareness of the dangers of weight gain and obesity by holding lectures or seminars for students and their parents.

References

- [1] GungorNk, Overweight and Obesity in Children and Adolescents: J Clin Res PediatrEndocrinol. 2014;6(3):129-143.
- [2] Hubbard VS. Defining overweight and obesity: what are the issues? Am J ClinNutr. 2000;72:1067-1068.
- [3]Nicolai JP, Lupiani JH, Wolf A J. An Integrative approach to obesity. In: Rakel D (ed).Integrative Medicine (3rd ed). W.B. Saunders (Elsevier), Philadelphia, PA2012:364-375.
- [4]Klish WJ. Clinical evaluation of the obese child and adolescent. In:Motil KJ, Geffner M (section eds) and Hoppin AG (Deputy ed).Up to date. www.uptodate.com©2013UpToDate.
- [5]Lustig RH and Weiss R. Disorders of energy balance. In: Sperling MA (ed) PediatricEndocrinology (third edition). Saunders Elsevier, Philadelphia, PA 2008:788-838.
- [6]BMICurves.<http://www.cdc.gov/growthcharts>.
- [7]Moore WE, Eichner JE, Cohn EM, Thompson DM, Kobza CE, Abbott KE. Blood pressure screening of school children in a multiracial school district: the Healthy Kids Project. Am J Hypertens.2009; 22: 351-356.
- [8]Rafrat M, Gargari BP, Safaiyan A. Prevalence of prehypertension and hypertension among adolescent high school girls in Tabriz, Iran. Food Nutr Bull.2010; 31: 461-465
- [9]Tirosh A, Afek A, Rudich A.Progression of normotensive adolescents to hypertensive adults:a study of 26,980 teenagers. Hypertension.2010; 56: 203-209.
- [10] Williams CL, Hayman LL, Daniels SR. Cardiovascular health in childhood. A statement for health professionals from the Committee on Atherosclerosis, Hypertension, and Obesity in the Young (AHOY) of the Council on Cardiovascular Disease in the Young, American Heart Association. Circulation.2002; 106: 143-160.
- [11]Genovesi S,Pieruzzi F,Giussani M. Analysis of heart period and arterial pressure variability in childhood hypertension:key role of baroreflex impairment.Hypertension.2008;51:1289-1294.
- [12]McNiece KL, Gupta-Malhotra M, Samuels J.Left ventricular hypertrophy in hypertensive adolescents: analysis of risk by 2004 National High Blood Pressure Education Program Working Group staging criteria. Hypertension.2007; 50: 392-395.
- [13]NafiuO O, Burke C, Lee J, Voepel-Lewis T, Malviya S and Kevin K.Neck

Circumference as a Screening Measure for Identifying Children With High Body Mass Index .Pediatrics. 2010;126:e306 .

[14]Cintra IP ; Passos MAZ ; Santos LC ; Machado HC and Fisberg M. Waist - to - Height Ratio Percentiles and Cutoffs for Obesity: A Cross-sectional Study in Brazilian Adolescents. J HEALTH POPUL NUTR. 2014; 32 (3): 411- 419.

[15]James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, et al. 2014 evidencebased guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8).JAMA. 2014;311:507-520.

[16] Flynn JT, Daniels SR, Hayman LL. American Heart Association Atherosclerosis, Hypertension and Obesity in Youth Committee of the Council on Cardiovascular Disease in the Young. Update: ambulatory blood pressure monitoring in children and adolescents: a scientific statement from the American Heart Association. Hypertension.2014; 63(5): 1116–1135 .

[17]Chen XU, Liu YU,Xizhuo Sun, Zhaoxia Yi,Honghui Li, .Comparison of body mass index, waist circumference, conicity index, and waist-to-height ratio for predicting incidence of hypertension: the rural Chinese cohort study. Journal of Human Hypertension.2018;32(3):228-235. doi: 10.1038/s41371-018-0033-6. Epub 2018 Feb 7.

[18]Hosseini M, Baikpour M, Yousefifard M, Fayaz M, Koohpayehzadeh J,

Ghelichkhani P.Blood pressure percentiles by age and body mass index for adults. Excli J. 2015 ;14:465-477.

[19] Choy C , Chan W , Chen T , Shih C , Wu L and Liao C. Waist circumference and risk of elevated blood pressure in children: a cross- sectional study BMC Public Health. 2011; 11 (613): 1–6.

[20]Kelishadi R,Heidari-Beni M, Qorbani M, Motamed-Gorji N, Motlagh M E, Ziaodini H, Taheri M , Ahadi Z, Aminae T, Heshmat R. Association of neck and wrist circumferences with cardio-metabolic risk in children and adolescents: The CASPIAN-V Study. Nutrition;32-38. doi: 10.1016/j.nut.2017.06.009.