

Prevalence of Fasciolosis and Cystic Echinococcosis in Slaughtered Cattle in Abattoir of Al-Nassiriyah City

Wesam J.Hansh

General Directorate of Education in Thi-Qar Province / Ministry of Education

Thi-Qar/ Iraq

wjhansh81@gmail.com

Qasim M. Al-Taher

Dept. Science (Biology)

College of Basic Education / University of Sumer

Thi-Qar/ Iraq

qasim20020@gmail.com

Muthenna M.T. Alebady

General Directorate of Education in Thi-Qar Province / Ministry of Education

Thi-Qar/ Iraq

Muthenna96@gmail.com

Abstract—The present study was conducted from January to May 2018 to identify the infection rate of fasciolosis and cystic echinococcosis in cattle slaughtered in the slaughterhouse of Nassiriyah Municipality, Thi-Qar province. During this study, 524 (247 males and 277 females) cattle were submitted for postmortem inspection. Chi-square is used to analyze the data. The prevalence of fasciolosis in the examined cattle was (7.25%) while the rate of infection with cystic Echinococcosis was (8.40%) with no significant difference. Infection with fasciolosis and cystic echinococcosis was higher in females than males. The current study showed significant relevance between the sex of cattle and the prevalence of both fasciolosis and cystic Echinococcosis ($P \leq 0.05$). The prevalence of fasciolosis and cystic Echinococcosis were significantly affected among the age groups in cattle ($P \leq 0.05$).

Keywords—Prevalence, Fasciolosis, Cystic Echinococcosis.

I. INTRODUCTION

Parasitism is vital in many agroecological areas and a critical issue to the agriculture sector and, thus, the global economy (Vercruyse and Claerebout, 2001). Parasitic diseases can decrease domestic animals' productivity in ruminants and cause death, which diminishes the supply of available human food, is a critical financial issue for breeders, and cause local economic losses (Acha and Szyfres, 2001). The high infection of diseases of helminthiasis (including lungworms, hydatidosis, and fasciolosis) in domestic animals is causing considerable damage to the disposal of organs and carcasses and decreasing the output and performance of animals (Getachew et al., 2012). Liver trematodes are one of the widespread parasitic diseases of humans and animals caused by the *Dicrocoelium dendriticum* and *Fasciola* genus, which live in the bile ducts of ruminants and humans (Martinez et al., 2013). Fasciolosis is a zoonotic disease induced by *Fasciola hepatica* and *Fasciola gigantica*, which is still an economically severe parasitic infection of animals (Hodzic et al., 2013). Sheep, cattle, and humans are definitive hosts, while snails from the *Lymnaea* genus act as

intermediate hosts (Karyakarte and Damle, 2004). Fasciolosis in cattle and buffaloes is caused by *Fasciola gigantica* in the Thi-Qar province, identified by polymerase chain reaction (PCR), while *Fasciola hepatica* is not diagnosed in the same province (AL-Salhey, 2009). Cystic echinococcosis, also known as hydatidosis, is a global parasitic disease induced by the larval stage (metacestode or hydatid cyst) of tapeworms belonging to the *Echinococcus* genus. There are four species within this genus. These are called *E. granulosus*, *E. multilocularis*, *E. oligarthrus*, and *E. vogeli* (Nakao et al., 2007). In Iraq, cystic echinococcosis is induced by *E. granulosus*, which is considered a hyperendemic disease (Molan, 1993). The definitive host is a dog (also other canines), and the intermediate host is sheep, cattle, pigs, goats, camels, and humans are accidental intermediate hosts (Karyakarte and Damle, 2004). The study demonstrates in Thi-Qar province that the majority of hydatid cysts samples which are isolated from humans, sheep, cattle, buffaloes, and camels, are caused by sheep strain (genotype 1) of *E. granulosus* using polymerase chain reaction (PCR) and sequence analysis of ribosomal DNA internal transcribed spacer subunit 1 (rDNA – ITS1) and mitochondrial Cytochrome c oxidase subunit 1 (mtCOX1) genes (Hansh, 2016). Inspection records of slaughtered animals have been applied as valuable sources for estimating epidemiological aspects of assured diseases in some countries (Mirani et al., 2000; Kara et al., 2009). Previous studies show that fasciolosis and cystic echinococcosis are endemic diseases in some parts of Iraq. Therefore the present study aims to identify the prevalence of fasciolosis and cystic echinococcosis in cattle slaughtered in the abattoir of Nassiriyah municipality.

II. MATERIALS AND METHODS

A. Study Area

The present study was conducted in the abattoir of Nassiriyah Municipality, located on the riverside of Euphrates in Thi-Qar province/Iraq, to investigate the

prevalence of Fasciolosis and Cystic Echinococcosis in slaughtered cattle. The period of study was conducted from January to May 2018. Abattoir was visited weekly (once a week)during the study period.

B. Data Collection

Five hundred twenty-four cattle (247 males and 277 females) were examined during the study period. The number of livers infected with Fasciolosis and livers and lungs infected with cystic echinococcosis of slaughtered cattle at the abattoir were recorded. Each organ in slaughtered cattle was examined macroscopically by visual inspection to detect hydatid cysts and sometimes to use an incision to investigate small hydatid cysts or by palpating bile ducts through using an incision to detect the infection of fasciolosis—other information on the slaughtered cattle including recording sex and age of the examined cattle. The prevalence of infection was calculated as a proportion by dividing the number of infected cattle by the total number of examined cattle.

C. Statistical Analysis

The registered data and the relevance between the impact of sex and age on the infected organs in cattle presented in the findings were analyzed using Chi-square tests in SPSS software. Differences were considered significant at P-value (≤ 0.05).

III. RESULTS

Prevalence of Fasciolosis and Cystic Echinococcosis

During this study, 524 slaughtered cattle were inspected at postmortem examination to detect the prevalence of infection. The total number of fasciolosis and cystic echinococcosis infections was 38(7.25%) and 44(8.40%), respectively. Statistically, no significant difference ($P \geq 0.05$) was noticed between the prevalence of both diseases, Table (1).

TABLE 1: PREVALENCE OF FASCIOLOSIS AND CYSTIC ECHINOCOCCOSIS IN CATTLE.


Disease	No. Examined Cattle	No. infected Cattle	% of Infection
Fasciolosis	524	38	7.25
Cystic Echinococcosis (Liver and Lungs)		44	8.40

The present study showed combined infections with fasciolosis and cystic echinococcosis in cattle livers; sometimes, the liver is infected with fasciolosis, and the

lungs are infected with cystic echinococcosis. (Macroscopic image 1, Table 2). According to statistical analysis, there is no significant difference between mixed infections.

TABLE 2: PREVALENCE OF COMBINED INFECTIONS IN CATTLE.

No. Examined Cattle	Infection of the liver with Fasciolosis and Cystic Echinococcosis	Infection of the liver with Fasciolosis and lung with Cystic Echinococcosis
524	8(1.53)%	3(0.57)%



Macroscopic image 1(A and B) combined infections of *Fasciolagigantica* (a) and hydatid cysts (b).

The present results showed that the Infection with fasciolosis was higher in females 35(12.64%) than in males 3(1.21%) with a significant statistical difference. Similarly, the prevalence of cystic echinococcosis was higher in females 43(15.52%) than in males 1(0.40%) with clear statistical variation, Table (3).

TABLE 3: THE EFFECT OF SEX ON CATTLE'S PREVALENCE OF FASCIOLOSIS AND CYSTIC ECHINOCOCCOSIS.

Sex	No. Examined Cattle	Fasciolosis		Cystic Echinococcosis	
		No. Infected	% of Infection	No. Infected	% of Infection
Male	247	3	1.21	1	0.40
Female	277	35	12.64	43	15.52

The present study indicated that the age group above >4 years recorded a high rate of infection 29(20.86%) in cattle infected with fasciolosis, and the age group < 2 years recorded a low rate of infection 4(1.67%) with an obvious statistical variation. Similarly, cystic echinococcosis occurred at a significantly higher rate in the age group above

>4 years 38(27.34%) compared to the infection within other age groups, Table (4).

TABLE 4: THE EFFECT OF AGE ON THE PREVALENCE OF FASCIOLOSIS AND CYSTIC ECHINOCOCCOSIS IN CATTLE.

Age group (year)	No. Examined Cattle	Fasciolosis		Cystic Echinococcosis	
		No. Infected	% of Infection	No. Infected	% of Infection
< 2	239	4	1.67	1	0.42
2 – 4	146	5	3.42	5	3.42
>4	139	29	20.86	38	27.34

IV. DISCUSSION

The present study showed no significant difference between the prevalence of fasciolosis (7.25%) and cystic echinococcosis (8.40%) in cattle. This result is comparable with studies of other workers. Kadir and Rasheed (2008) found that the highest rate of cystic echinococcosis was seen in cattle (4.38%), while the rate of liver fluke was (2.63%) in Kirkuk slaughterhouse, and also Ahmed and Rasheed (2013) reported that the rate of infection by hydatid cyst and liver fluke in cattle was (5.5%) and (3.2%) respectively in Kirkuk abattoir. Alatabiet *al.* (2017), in AL-Najaf slaughterhouse, recorded that the incidence of liver fluke was (0.71%) and hydatid cysts was (1.59%) and these results are less than the findings of the current study. Mohamadzadeh *et al.* (2016) revealed that the prevalence in cattle was (2.1%) and (1.65%) for cystic echinococcosis and fasciolosis, respectively, in Kazerun abattoir in Iran. The results of this study were relatively similar to the study of Ezatpouret *al.* (2014), who recorded in Iran the prevalence of cystic echinococcosis (9.4%) and fasciolosis (7.6%) in cattle. In Turkey, Balkaya and Simsek (2010) recorded a very high prevalence of cystic echinococcosis (34.3%), more than fasciolosis (21%). The current study disagrees with Mellauet *al.* (2010), who indicated in Tanzania that the prevalence was higher in cattle infected with fasciolosis (8.6%) than in cystic echinococcosis (3.2%). Boralet *al.* (2013) recorded that in Egypt, the prevalence of fasciolosis 2.67 % in cattle, more than cystic echinococcosis (0.61%). The differences in the prevalence may be linked to geographical allocation, environmental variation, sample size, and study time.

The prevalence of fasciolosis and cystic echinococcosis in an area is influenced by several factors comprising hosts, parasites, and environmental effects. The parasites and their intermediate and definitive hosts facilitate the life cycle

maintenance and can cause the incidence of these diseases. Environmental factors such as temperature, moisture, and rainfall especially affect the hatching of *Fasciolagigantica* ova, the viability of metacercaria, and the population of snails. Furthermore, temperature and moisture affect the survival of *Echinococcusgranulosus* ova. Phiriet *al.* (2005) mentioned that the seasonal variation affected the distribution pattern of *F. gigantica*. In addition, the prevalence of fasciolosis is increased in areas with a high rainfall range than in areas with very little rainfall. Getawet *al.* (2010) referred to several factors which cause the prevalence of cystic echinococcosis, such as the random spread of abattoirs, the absence of the rigorous healthy condition in meat inspection, and the spread of stray dogs and their feeding on infected viscera with hydatid cysts and subsequent contamination of pasture and grazing fields.

The current study exhibited that the livers of cattle exposed to infection by *F.gigantica* together with hydatid cysts. Furthermore, the liver of the cattle is infected with *F.gigantica* and the lungs with hydatid cysts. The reasons of combined infection may occur because the liver is an important site for these parasites, both metacercaria of *F.gigantica* and eggs of *E.granulosus* encyst in the small intestine and penetrate the intestinal mucosa and then reach the liver by portal circulation and lodge in biliary tracts (*F.gigantica*) and formed hydatid cysts of *E.granulosus*. Another reason is that the mechanisms of immunological defense, which is stimulated against infection by *F.gigantica*, may not prevent the occurrence of infection by hydatid cysts and vice versa. Hussienet *al.* (2015) recorded the presence of both fasciolosis and cystic echinococcosis jointly in cattle livers, with a recorded prevalence rate (of 3.93%) in Ethiopia.

The current study showed that fasciolosis and cystic echinococcosis was prevalent at higher rates in females of the infected cattle than in the males, with a significant difference. This variation between the infection rates in females and males may be because the males are being slaughtered at an early age, nutrition type, and most male cattle are brought up in unique places. Therefore, all these reasons reduce the rates of infection in males. At the same time, the females remain for a longer period used for reproductive purposes. Rahmianet *al.* (2012) identified that females are most affected by cystic echinococcosis than males due to living for long ages. Iqbalet *al.* (2012) stated other reasons which affect the susceptibility of the females, such as slaughtering at a late age and the stress of pregnancy, parturition, and lactation. The finding of the current study is comparable to results reported by Daryaniet *al.* (2006) in Iran, which recorded prevalence of fasciolosis in female cattle (38.4 %) is more than in males (12.2 %), and Azamiet *al.* (2013) showed the prevalence of cystic

echinococcosis in females of cattle (9.1%) and males (3.1%) in Iran.

According to statistical analysis, a significant difference was observed among age groups and the percentage of infection with fasciolosis and cystic echinococcosis in the cattle. Cattle above 4 years are highly susceptible to infection than other age groups. This result agrees with the findings of Molina *et al.* (2005), who showed that the prevalence of fasciolosis in cattle and buffaloes was higher in an age of older animals. Debas and Ibrahim (2013) recorded the prevalence of cystic echinococcosis in the age group above 5 years (31.2%), whereas they recorded (8.8%) in the age group less than 5 years. The reasons for incidence in aged animals may be because they remain for longer, especially the females used for reproductive purposes, production of milk, and other benefits, and this causes weakness in resistance to diseases. Ibrahim *et al.* (2008) demonstrated that age variation could be again linked with differences in infection exposure because aged animals may have been exposed to more infective stages.

V. CONCLUSION

Fasciolosis and cystic echinococcosis in cattle are endemic parasitic diseases in Thi-Qar province/ Iraq. The prevalence documented in this study may contribute to determining the prevention programs. Meat inspection at postmortem is a suitable method for providing helpful information about the infections of these diseases.

ACKNOWLEDGMENT

The authors would like to express special thanks to staff members of the abattoir of Nassiriyah Municipality for their assistance during this study.

REFERENCES

- Acha, N. and Szyfres, B. (2001). Zoonoses and communicable diseases common to man and animals. 3rd ed. Washington, PP:7 – 9.
- Ahmed, L. M and Rasheed, S. A. (2013). Distribution of liver and lung helminthic infections among slaughtered animals in Kirkuk abattoir. J. Genet. Environ. Resour. Conserv., 1(1):36 - 40.
- Alatabi, A. C.; Al-dujaily, A. H and Al-mialy, A. J. (2017). Study the Rate of Hydatid Cysts, Liver Fluke, Pneumonia, and Hepatitis in Al-Najaf Slaughter House, Al-Najaf, Iraq. Kufa, J. Vet. Med. Sci., 8(2): 137 – 142.
- AL-Salhey, W. J. H. (2009). Epidemiological and Pathological Study on *Fasciolagigantica* of Cattle and Buffaloes at Thi-Qar Province. M.Sc. Thesis, Univ. Thi-Qar, pp:138.
- Azami, M.; Anvarinejad, M.; Ezatpour, B. and Alirezaei, M. (2013). Prevalence of hydatidosis in slaughtered animals in Iran. *Turkiye Parazitol. Derg.*, 37: 102 – 106.
- Balka Balkaya, I. and Simsek, S. (2010). Prevalence and Economic Importance of Hydatidosis and Fasciolosis in Slaughtered Cattle in Erzurum province of Turkey. *Kafkas, Univ. Vet. Fak. Derg.*, 16(5): 793 – 797.
- Borai, M. G. E.; Nagi, A. A.; Gab-Allah, M. S.; El Mashad, A. I. and Moustafa, S. A. (2013). Comparative Pathological Studies On Parasitic Affections Of Liver IN Farm Animals. *Benha, Vet. Med. J.*, 25(2): 284-295.
- Daryani, A.; Alaei, R.; Arab, R.; Sharif, M.; Dehghan, M. H.; and Ziaei, H. (2006). Prevalence of Liver Fluke Infection in Slaughtered Animals in Ardabil Province, Northwestern Iran. *J. Ani & Vet. Adv.*, 5: 408 – 411.
- Debas, E. and Ibrahim, N. (2013). Prevalence and Economic Importance of Hydatidosis in cattle slaughtered at North Gonder Elfora Abattoir. *Europ. J. Appl. Sci.*, 5(1): 29 – 35.
- Ezatpour, B.; Hasanvand, A.; Azami, M.; Mahmoudvand, H. and Anbari, K. (2014). A slaughterhouse study on prevalence of some helminths of cattle in Lorestan province, West Iran. *Asian, Pac. J. Trop. Dis.*, 4(5): 416 – 420.
- Getachew, H.; Guadu, T.; Fentahun, T. and Chanie, M. (2012). Small ruminants Hydatidosis: occurrence and Economic Importance in Addis Ababa Abattoir. *Global Veterinaria* 8(2): 160-167.
- Getaw, A.; Beyene, D.; Ayana, D.; Megersa, B. and Abunna, F. (2010). Hydatidosis: Prevalence and its economic importance in ruminants slaughtered at Adama municipal abattoir, Central Oromia, Ethiopia. *Act. Trop.*, 113: 221-225.
- Hansh, W. J. (2016). Biological Study and Molecular Identification on Hydatidosis in Intermediate Hosts Depending on Sequence Analysis for *rDNA – ITS1* and *mtCOX1* Genes in Thi-Qar Province. Ph.D. Thesis, Univ. Basrah, PP: 186.
- Hodzic, A.; Zuko, A.; Avdic, R.; Alic, A.; Omeragic, J. and Jazic, A. (2013). Influence of *Fasciola hepatica* on serum biochemical parameters and vascular and biliary system of sheep liver. *Ira. J. Parasitol.*, 8(1): 92-98.
- Hussien, D.; Kerala, F.; Hagos, Y and Teklu, A. (2015). Prevalence of Fasciolosis and Hydatidosis in Male Cattle Slaughtered at Butajira Municipal Abattoir, Southern Ethiopia. *Act. Parasitol. Global.*, 6 (1): 49-54.
- Ibrahim, M. M.; Al-Ghamdi, M. A. and Al-Gahmid, M. M. (2008). Helminths community of veterinary importance of livestock in relation to some ecological and biological factors. *Turkiye, Parazitoloji, Dergisi*, 32: 42 – 47.
- Iqbal, H. J.; Maqbool, A.; Lateef, M.; Khan, M. A.; Riaz, A.; Mahmood, A.; Atif, F. A.; Ali, Z. and Ahmad, M. S. (2012). Studies on hydatidosis in sheep and goats at Lahore, Pakistan. *J. Anim. Plant. Sci.*, 22: 894 – 897.
- Kadir, M. A. and Rasheed, S. A. (2008). Prevalence of some parasitic helminthes among slaughtered

ruminants in Kirkuk slaughterhouse, Kirkuk, Iraq. Iraqi, J. Vet. Sci., 22(2): 81 – 85.

Kara, M.; Gicik, Y.; Sari, B.; Bulut, B. and Arslan, M. O.(2009). A slaughterhouse study on prevalence of some helminthes of cattle and sheep in Malatya province, Turkey. J. Anim. Vet. Adv., 8:2200 – 2205.

Karyakarte, R. P. and Damle, A. S. (2004). Medical Parasitology. Books and Allied (P) Ltd. PP. 288.

Martinez-Valladares, M.; Robles-Perez, D.; Martinez- Perez, J.M.; Cordero-Perez, C.; Famularo-Mdel, R. and Fernandez-Pato, N.(2013). Prevalence of gastrointestinal nematodes and *Fasciolahepatica* in sheep in the northwest of Spain: relation to climatic condition and/or man-made environmental modifications.Parasit Vectors. 6:282.

Mellau, L. S. B.; Nonga, H. E. and Karimuribo, E. D. (2010). A Slaughterhouse Survey of Liver Lesions in Slaughtered Cattle, Sheep and Goats at Arusha, Tanzania. Res. J. Vet. Sci., 1 – 10.

Mirani, H. A.; Akthar, N.; Brohe, M. A.; Bughiu, S. and Oad, F. C.(2000).Hydatidosis in buffaloes at Larkana slaughterhouse. Pak. J. Biol. Sci., 3(8): 1331 – 1312.

Mohamadzadeh, T.; Shams, S.; Khanaliha, K.; Marhamatizadeh, M. H. and Vafa, A. (2016). A study on prevalence of some helminthic infections of the liver and

lungs among ruminants in abattoir of Fars province, Iran. Archives, Razi, Instit., 71(4): 245 – 251.

Molan, A. L. (1993). Epidemiology of Hydatidosis and Echinococcosis in Theqar province, southern Iraq. Japan. J. Med. Sci. Biol., 46: 29 – 35.

Molina, E. C.; Gonzage, E. A. and Lumbayo, L. A. (2005). Prevalence of infection with *F. gigantica* and its relationship to carcase and liver weight fluke and egg counts in Slaughter cattle and buffaloes in South Mindanao Philippines. Trop. Anim. Health. and Prod., 37: 215 – 221...

Nakao, M.; McManus, D. P.; Schantz, P. M.; Craig, P. S. and Ito, A. (2007). A molecular phylogeny of the genus *Echinococcus* inferred from complete mitochondrial genomes. Parasitology, 134: 713 – 722.

Phiri, A. M.; Phiri, I. k.; Sikasunge, C. S. and Monrad, J. (2005). Prevalence of fasciolosis in Zambian cattle observed at selected abattoirs with Emphasis on Age, Sex, and origin. J. Vet. Med. B., 52: 414 – 416.

Rahmani, K.; Radfar, M. H. and Adinehbeigi, K. (2012).Hydatidosis: prevalence and biometrical studies in sheep in Kerman area, southeast of Iran. Comp. Clin. Pathol., PP: 1 – 6.

Vercruysse, J. and Claerebout, E. (2001).Treatment vs. non-treatment of helminth infections in cattle: defining the thresholds.Vet.Parasitol., 98: 195-214.