

ANTIBODIES AGAINST ERYTHROCYTES ENHANCE INVASION OF MOUSE RETICULOCYTES BY *PLASMODIUM* *BERGHEI*⁺

Khalid Majeed Dakhel^{*}

Abstract:

Twenty mice were injected i.p. with 1×10^5 *P. berghei* infected red cells . Blood smears of these mice were prepared daily till the mice were died . Mice were categorized in three groups depending on similar type of course of parasitaemia after inoculation . Seven mice of Group A ($A_1 - A_7$) showed similar type of behavior and died due to low parasite infection . Another group of seven mice (B_1 to B_7) showed a gradual increase in parasitaemia till mice died due to high infection . Group C (C_1 to C_6) smears showed a vary rise and fall in infection and all the mice were cleared of parasite infection .

The antibody titre as detected by IFA , observed in these groups was 1 : 2048 , 1 : 1024 and 1 : 4096 in groups A , B and C respectively . The group C which showed protective immunity against *P. berghei* infection depicted strong fluorescence against normal erythrocyte membrane too . The fluorescence against normal RBCs membrane was observed very weak in group A , whereas it was negligible in group B .

المستخلص :

حقن ٢٠ حيواناً مختبرياً بواسطة 10x1 كريات دم حمراء مصابة بـ *P. berghei* و اخذت مسحات دم لهذه الحيوانات يومياً ولحين هلاك هذه الحيوانات وقسمت هذه الحيوانات الى ثلاثة مجاميع اعتماداً على الاستجابة للإصابة بعد الحقن وهي مجموعة A ($A_1 - A_7$) اعطت نفس السلوك وقد تم هلاك هذه الحيوانات نتيجة للإصابة بالطفيليات القليلة . المجموعة B ($B_1 - B_7$) اعطت زيادة تدريجية في نسبة الإصابة بالطفيليات ولحين هلاك الحيوانات . مجموعة C ($C_1 - C_6$) اعطت زيادة كبيرة وإصابة كاملة وكانت جميع الحيوانات خالية من الطفيليات .

نسبة الاجسام المضادة حددت بواسطة اختبار فحص الاجسام المناعية (IFA) وقد لوحظت النتائج التالية :

1 : 2048 , 1 : 1024 , 1 : 4096 للمجاميع الثلاثة A , B , C وعلى التوالي .
وقد اعطت المجموعة C مناعة كبيرة جداً للإصابة بالطفيليات وقد كانت A ضعيفة جداً .

Introduction

⁺ Received on 2007/5/2 , Accepted on 2008/11/30

^{*} Assist prof. -Community Health Dept. Nassiriah Technical Institute

Malaria continues to cause morbidity and mortality on a large scale in tropical countries . According to the report given by WHO [1] , human malaria parasite causes 300 – 500 million clinical cases which include young children under 5-6 years of age , pregnant women and non – immune travelers . Annual mortality caused by malaria ranges from 0.5 – 2.5 million including over a million death in children aged under five [2] .

Earlier studies have greatly influenced the knowledge about invasion of parasite in red cells [3] and the pioneer work suggested the role of receptors in recognition of susceptible host cells by merozoites for invasion .[4]

In vitro studies of *P . knowlesi* have shown the behavior of merozoites to be even more complex, since the process of invasion involves various sequential changes in shapes of both host and parasite cells [5]

A study was described that the interactions between *P . falciparum* merozoites and human RBC is mediated by specific parasite proteins and sialoglycoproteins (SGPS) on the surface of the host cell.[6]

The first *plasmodium* molecule found to be located in the microneme organelles of the merozoite is the 135 kDa *P . Knowlesi* Duffy binding proteins (pk DBP) [7] . The gene encoding this proteins was subsequently used as a probe to clone the corresponding gene from *P.vivax* (Pv DBP) [8].

It was suggested that the invasion of glycophorin A and B deficient erythrocytes by *P . falciparum* as trypsin or neuraminidase treated glycophorin B deficient cells were refractory to the parasite EBA-175 , a parasite protein appears to bind to glycophorin A . [9,10] suggested the EBA as transmembrane proteins characterized by conserved cysteine rich domain expressed in the micronemes of invasive merozoites .

Strong evidences have implicated EBA – 175 as being a parasite – ligand used in the recognition of human RBC by *p . falciparum* merozoite [11] .

Merozoite surface protein 2 (MSP-2) a highly polymeric in nature , generates antiparasite humoral response in mice [12] .

Vaccination of animals against sporozoite leads to formation of antibodies that inhibit sporozoite motility and block the invasion of blood vessels by sporozoite [13].

A study found that plasmodium – infected erythrocyte release a low molecular weight soluble factor that inhibits the activation of macrophages and dendritic cells [14].

The present work is an attempt to understand the preferential invasion of reticulocytes by the parasite in *p . berghei* infection .

Material and Methods

Parasite

Normal mice were injected intraperitoneally (i.p.) with parasitized red cell in citrate saline and the course of parasitaemia was monitored . Accordingly the mice were grouped in different categories .

NK 65 , a lethal strain of *P . berghei* was maintained in Balb / C mice swiss white mice . 20 mice each of 20-25 gms . were inoculated with 1×10^5 parasitized cells and daily course of infection was monitored by Giemsa stained blood smears . The percentage of total number of red blood cells :

Collection of Blood

The serum was collected in two ways :

- 1- A few drops of blood were collected from each mouse separately . It was allowed to clot at room temperature for one hour . The clot was broken and serum was isolated .
- 2- Mice were anesthetized by diethyl ether and bled by jugular vein incision . The blood from broken clot was then centrifuged at 1,000 g for 15 minutes and the serum was aspirated and collected .

Statistics analysis :

Chi (X^2) test was used for statically analysis in present study .

Indirect Fluorescent Antibody (IFA) test

IFA test was done according to slightly modified method (1972) [13] . The slides were counter stained with Evan's blue and studied under UV . Fluorescence microscope (Leica DMILS , Germany) .

Results :

Group A :

The group contained seven mice (A_1 to A_7) . All the mice of this group died due to infection and maximum parasitaemia remained between (24% - 60%) in most of mice (Fig . 1) . One of mice had only 7% maximum parasitaemia . The parasitaemia ranged between (3 - 15%) till day 10 and both reticulocytes and erythrocytes were infected . One mice died on day 12 , other two on day 14 , and one on day 16 . In the remaining three mice the parasitaemia further increased and reached about 48% to 60% followed by increase in reticulocytosis and decrease in parasitaemia . These mice were found dead due to infection on day 24 , 26 and 28 .

Group B :

This group also contained seven mice (B_1 to B_7) but on the contrary to group A the animals died to high parasitaemia (Fig . 2) . Maximum parasitaemia ranged between 58% to 82% . Both erythrocytes and reticulocytes were infected but with gradual increase in parasitaemia and appearance of reticulocytes . In two of the mice of this group more than 80% cell were infected . The animals died because of high parasitaemia between day 13 to day 17 .

Group C :

This group contained 6 mice (C_1 to C_6) . In this group the parasite was observed on 14th day of post inoculation in 4 mice . Whereas , two mice became infected on 5th day . There was Zig – Zag course of parasitaemia . In all the mice initially most of the erythrocytes were infected along with a few reticulocytes (Fig . 3) with the gradual increase in parasitaemia , reticulocytosis occurred and the post migrated from red cells to the reticulocytes and finally the parasite was exclusively confined to the reticulocytes . An increase in number of WBCs were observed during later days .

Futher smears showed decrease in parasitaemia between day 15 and day 20 and finally no parasite was observed in the smear of this group after 25th day .

Indirect Fluorescent Antibody (IFA) test :

IFA test was employed to detect the presence of antibodies in parasitized mice . The IFA slides were counter stained with Evan's blue to avoid any non-specific reaction .

The appearance of antimalarial antibodies in mice of different groups and the titre of antibodies are depicted in Table 1 .

Group A :

Antibodies appeared in one of the mice on day 9 whereas , on 10th day all the mice showed positive titre of 1 : 32 (Fig. 4) . Sera collected on day prior to the death of infected animals showed varying IFA titre . Four mice (A₁ , A₂ , A₆ and A₇) showed varying IFA titre between 1 : 256 and 1 : 1024 while others showed 1 : 2048 (Fig . 5) .

IFA that was significant ($P < 0.05$) with the infection percentage .

Group B :

In this group antimalarial antibodies were observed on day 9th in 2 mice (B₁ to B₄) whereas , by next day appeared in other 5 mice (Fig. 6) . The IFA titre ranged between 1 : 64 which died on day 13th and 1 : 512 which died on day 15th and 1 : 1024 which died on 16th and 17th day (Fig . 7) .

NO or almost negligible fluorescence was observed with normal red cell antigen . no significant ($P < 0.05$) between IFA and infection percentage .

Group C :

The antimalarial antibody was observed on day 10th post inoculation in all the 6 mice (Fig . 8) . IFA titre of 1 : 4096 was observed in the sera of these mice when sacrificed on 36th day . The IFA titre of this group showed in (Fig . 9) . A very strong IFA reaction was observed against all the stages of the parasite in case of parasitized erythrocytes or reticulocytes . The reaction with the host cell antigen revealed that more fluorescence was observed with erythrocytes as compared with the reticulocytes .

IFA that was significant ($P < 0.05$) with the infection percent .

Discussion

Susceptibility of Balb / C strain of white swiss mice to NK – 65 strain of *P.berghei* as an asynchronous parasite it was well established the suscep [14] .

Preferential invasion of reticulocytes by merozoites of *P . berghei* was also observed by others [15]; [16] ;[17] . *P. vivax* invades reticulocytes exclusively and reticulocyte binding gene expressed PvRBP-1 and PvRBP-2 , two proteins which are shown to be responsible for preferential invasion of *P . vivax* in reticulocytes . *P . berghei* has shown preference for reticulocytes , however , PvRBP gene probes did not cross hybridise to DNA of *P . falciparum* , *P . knowlesi* and *P . berghei* [18] .

In Vivo reticulocytes mature to red cells within 24-36 hours after release from bone marrow into circulation [19] . During the maturation process of erythrocytes a number of constituent proteins from the membranes of the reticulocytes are lost [20] ; [21] ; [22] . The phenomenon of invasion of red blood cells by the parasite is a multistep process involving attachment of the host cell and internalization of the parasite [23] ; [24] . The relationship between age of the host cell and invasion by *P.falciparum* indicated that young erythrocytes in particular have an increased susceptibility to the parasite [25] . In *P . berghei* the parasite parasitizes reticulocytes preferably initially and exclusively later , if it remains in the host cell for 2-3 weeks or more .

Although the IFA is useful in estimating exposure to malaria , previous work has indicated that many antibodies directed against some parasite antigens detected by IFA , depicted appearance of antibodies between 7th to 10th day of post inoculation . There was a gradual increase in antibody level in different mice . Some mice died due

to low parasitaemia with IFA titre of 1 : 2048 after 24 days whereas IFA titre of 1 : 1024 was observed in some mice which died due to high parasitaemia between 16th and 17th day , or some mice recovered from infection and the parasite was not observed in the smear with IFA titre of 1 : 4096 . An increase in WBC'S was observed during later days depicting some role of cell mediated immunity along with humoral immunity , both playing a major role in protection .

Finally humoral and cell mediated immunity along with some genetic factors may be responsible for protective immunity in these mice .

Table : 1 : Appearance of antimalarial antibodies in mice of different groups and the titre of antibody .

Group	Antibody appearance		Maximum Antibody Titre in sera collected *
	Day	No . of mice	
A	9 10	1 6	1 : 2048
B	7 8	2 5	1 : 1024
C	10	6	1 : 4096




Fig. 1 : Course of *P. berghei* in mice resulting in death of animals due to low parasitaemia.




Fig. 2 : Course of *P. berghei* in mice resulting in death animals due to high parasitaemia




Fig. 3 : Course of *P. berghei* infection in mice and subsequent protection of the animals




Fig. 4 : Histogram showing antimalarial antibodies in the sera of mice of group A. Animals showing 1 : 32 or more titre were positive.




Fig. 5 : Histogram showing IFA titre in the sera samples collected in group A mice.




Fig. 6 : Histogram showing antimalarial antibodies in the sera of mice of group B. Animals showing 1 : 32 or more titre were positive.




Fig. 7 : Histogram showing IFA titre in the last sera samples collected in group B mice.




Fig. 8 : Histogram showing antimalarial antibodies in the sera of mice of group C. Animals showing 1 : 32 or more titre were positive.




Fig. 9 : Histogram showing IFA titre in sera samples of group C mice.

References :

- 1- World Health organisation . Rolling back malaria . WHO World Health Report , 49-64. (1999)
- 2- White , N.J., Nosten , F ., Looareesuwan , S ., Watkins , W.M. , Marsh , K ., Snow , R.W., Kokwaro , G ., Ouma , J ., Hein , T.T., Moltneux , M.E ., Taylor , T.E ., Newbold , C.I. , Buehler , T.K., Danis , M., Greenwood , B.M., Anderson , R.M. and Ollirao , P..Averting a malaria disaster.*Lancet* , 353 , 1965-1967. (1999)
- 3- Ball , E.G., Anfinson , C.B., Geiman , Q.M., McKee , R.W. and Ormsbee , R.A..In vitro growth and multiplication of the malarial parasite , *P.knowlesi*.*Science* , 101 , 542 – 544. (1945)
- 4- McGhee , R.B. The adaptation of the avian malaria parasite *P.lophurae* to a continuous existence in baby mouse.*J.Infect.Dis* ., 88 , 86 – 97. (1951).
- 5- Dvorak , J.A., Miller , L.H ., Whitehouse , H.C.and Shiroishi , T.Invasion of erythrocytes by malaria merozoites.*Science* , 187 , 748 – 750. (1975).
- 6- Rodriguez , M.H.and Jungery , M.Glycoprotein recognition mediates attachment of *P.chabaudi* to mouse erythrocytes.*Infect.Immun* ., 55 , 187 – 192 .(1987)
- 7- Adams , J.H ., Hudson , D.E ., Torii , M ., Ward , G.E ., Wellens , T.E ., Aikawa , M.and Miller , L.H.The duffy receptor family of *plasmodium knowlesi* is located within the micronemes of invasion merozoites.*Cell* , 63 , 141 – 153. (1990)
- 8- Fang , X ., Kaslow , D.C ., Adams , J.H.and Miller , L.H.Cloning of the *plasmodium vivax* Duffy receptor.*Mol.Biochem.Parasitol* ., 44 , 125 – 132. (1991)
- 9- Dolan , S.A ., Proctor , J.L ., Alling , D.W ., Okubo , Y ., Wellems , T.E.and Miller , L.H.Glycophorin B as an EBA – 175 independent *plasmodium falciparum* receptor of human erythrocytes.*Mol.Biochem.Parasitol* ., 64 , 55 – 63. (1994)

- 10- Kappe , S.H ., Curley , G.P ., Neo , A.R ., Dalton , J.P.and Adams , J.H. Erythrocyte binding protein homologues of rodent malaria parasites.*Mol.Biochem.Parasitol* ., 89 , 137 – 148. (1997).
- 11- Rodriguez , L.E ., Urquiza , M ., Ocampo , M ., Suarez , J ., Curtidor , H ., Guzman , F ., Vargas , L.E ., Trivinos , M ., Rosas , M.and Patarroyo , M.E. *Plasmodium falciparum* EBA – 175 KDa protein peptides which bind to human red blood cells.*Parasitology* , 120 , 255 – 235. (2000).
- 12- Lawrence , N ., Stowers , A ., Mann , V ., Taylor , D.and Saul , A. Recombinant chimeric proteins generated from conserved regions of *Plasmodium falciparum* merozoite surface proteins 2 generate antiparasite humoral responses in mice.*Parasite immunology* , 22 , 211 – 221. (2000).
- 13- Vanderberg JP , Frevert U, intravital microscopy demonstrating antibody – mediated immobilisation of *P. berghei* sporozoites in vivo . *J . parasitol* . Aug ; 34 (9) : 991-6. 2004 .
- 14- J.C. Hafalla , U.Rai U , A. Morrot , D. Bernal – Rubio , F.zavala and A. Rudriquez . priming of CD8⁺ T cell responses following immunization with heat killed plasmodium sporozoites . *European journal of immunology* . may ; 36 (5) :179-86 . 2006 .
- 15- Ramakrishnan ,S.P. and Prakash , S.Studies on *P.berghei* n.sp.Vinke and Lips 1948.Morphology , periodicity and pathogenicity in blood induce infection in mice , rats and garden squirrels.*Indian J.Malarial* ., 4 , 369 – 375. .(1950)
- 16- Janse , C .J ., Mons , B ., Croon , J.J.A.B.and Vander Kaay , H.J.Long term in vitro culture of *P.berghei* and preliminary observations on gametocytogenesis.*Int.J.parasitol* ., 14 , 317 – 320. .(1984)
- 17- Janse , C.J ., Boorrsma , E.G ., Ramesar , J ., Grobbee , M.J.and Mons , B. Host cell specificity and schizogony of *P.berghei* under different in vitro conditions.*Int.J.parasitol* ., 19 , 509 – 514. (1989).
- 18- Galinski , M.R ., Medina , C.C ., Ingravallo , P ., Barnwell , J.W. A reticulocyte – binding protein complex of *P.vivax* merozoites.*Cell* 69 , 1213 – 1226. (1992).
- 19- Gronowicz , G ., Swift , H.and Steck , T.L. Maturation of the reticulocyte in vitro.*J.cell.sci* ., 71 , 177 – 197. (1984).
- 20- Patel , V.P ., Ciechanover , A ., Platt , O.and Lodish , H.F. Mammalian reticulocytes lose adhesion to fibronectin during maturation to reticulocytes.*Proc.Natl.Acad.Sci* ., USA , 82 , 440 – 444. (1985).
- 21- Rapoport , S.M. The reticulocyte.Boca Raton Florida : CRC Press. (1986).
- 22- Lazarides , E. From genes to structural morphogenesis : the genesis and epigenesis of a red blood cell.*Cell* , 51 , 345 – 356. (1987).
- 23- Ward , G.E ., Chitnis , C.E.and Miller , L.HStrategies for intracellular survival of microbes (ed.Russel.D .) Saunders , 155 – 190. .(1994).
- 24- Wilson , R.S.M. Biochemistry of red cell invasion.*Blood cells* , 16 , 237 – 252. (1990).
- 25- Pasvol , G ., Weatherall , D.J.and Wilson , R.J.M. The increased susceptibility of young red cells to invasion by the malarial parasite.*British Journal of Haematology* , 45 , 285 – 295. (1979).