



Effect of Some Environmental Factors and Antibiotics on the Growth of Different Species of Rhizobium

Sara Musbah Mohammed^{*}, Raad Hassani Sultan

Department of Biology, College of Education for Pure Sciences, University of Mosul, Mosul, Iraq.

Corresponding Author: sara.22esp4@student.uomosul.edu.iq

Citation: Mohammed SM, Sultan RH. Effect of Some Environmental Factors and Antibiotics on the Growth of Different Species of Rhizobium. Al-Kitab J. Pure Sci. [Internet]. 2024 July. 10 [cited 2024 July. 10];8(2):109-119. Available from: <https://doi.org/10.32441/kjps.08.02.p9>.

Keywords: Antibiotic, Herbicide, Rhizobia, and Isolation.

Article History

Received	10 May.	2024
Accepted	13 Jun.	2024
Available online	10 July	2024

© 2024. THIS IS AN OPEN-ACCESS ARTICLE UNDER THE CC BY LICENSE
<http://creativecommons.org/licenses/by/4.0/>



Abstract:

Six native rhizobial isolates from various cultural zones in Ninavah-Governate, Iraq, were used in this investigation. The following rhizobial strains were isolated from leguminous plant root nodules: *Rhododendron japonicum* SM29 from *Glycin max* L., *Rhizobium leguminosarum* bv. *trifolii* SM35 from *Trifolium alexanrinum* L., *Rhizobium leguminosarum* bv. *viciae* SM10 from *Vicia faba* L., *Ensifer ferdii* bv. *Fredii* SM13 from *Vigna unguiclata* L., *Ensifer meliloti* SM28 from *Medicago sativa* L., and *Rhizobium leguminosarum* bv. *phaseoli* SM42 from *Phaseolus vulgaris* L. Rhizobial bacteria were identified by the spherical, clear colonies seen after cultural investigation. Methyl red and Voges-Proskuar biochemical tests yielded negative results, but urease, catalase, indole, starch, Congo red, citrate, and motility tests yielded positive results. A high tolerance was found by the KNO₃ tolerance test.

Keywords: Antibiotic, Herbicide, Rhizobia, Isolation.

دراسة تأثير بعض العوامل البيئية والمضادات الحيوية على نمو اجناس مختلفة من

Rhizobium

سارة مصباح محمد^{*}، رعد حساني سلطان

قسم علم الأحياء، كلية التربية للعلوم الصرفة، جامعة الموصل، الموصل، العراق

sara.22esp4@student.uomosul.edu.iq, dr.raadsultan@uomosul.edu.iq

الخلاصة:

تم في هذه الدراسة عزل ستة عزلات من الرايزوبيا المحلية من مناطق مختلفة من محافظة نينوى-العراق. تم عزل هذه العزلات من العقد الجذرية من النباتات البقولية وكما يأتي:

Ensifer ferdii bv. *Fredii* SM13 من نبات الباقلاء, *Rhizobium leguminosarum* bv. *viciae* SM10 من نبات اللوبيا, *Ensifer meliloti* SM28 من نبات الجت, *Rhododendron japonicum* SM29 من نبات فول الصويا, *Rhizobium leguminosarum* bv. *trifolii* SM35 من نبات البرسيم والعزلة *Rhizobium leguminosarum* bv. *phaseoli* SM42 من نبات الفاصوليا. أظهر الفحص الزراعي مستعمرات دائرية شفافة دلالة على انها بكتريا الرايزوبيوم. أظهرت الفحوصات الكيموحيوية نتيجة سالبة لاختباري احمر المثيل وفوكس-بروسكر, في حين أظهرت اختبارات اليوريز والكتاليز والأندول وتحلل النشا و احمر الكونغو والسترات والحركة نتيجة موجبة ولجميع العزلات. اظهر اختبار تحمل ملح نترات البوتاسيوم درجة تحمل عالية لعزلات الرايزوبيا المحلية، حيث وصل الى 10% (وزن/حجم) وللعزلتين SM10 و SM28. كانت جميع العزلات مقاومة للأمبسلين 25 مايكروغرام/مل باستثناء العزلة SM42 حيث أظهرت حساسية لهذا المضاد الحيوي. أعطت جميع العزلات مقاومة عالية للمبيد العشبي الغليفوسات بتركيز 50 ملغم/مل.

الكلمات المفتاحية: عزل، رايزوبيا، مضادات حيوية، مبيدات عشبية.

1. Introduction:

To manage and lessen weed growth on agricultural grounds, chemicals have been employed extensively in agriculture [1]. Roughly one-third of agricultural land is treated with herbicides these days [2-4]. Because so few herbicides reach the intended organisms, the destiny of these chemicals has come under scrutiny [5, 6]. Thus, herbicides have an impact on the ecosystem's ability to operate, either directly or indirectly [7]. Glyphosate is a pre-emergence herbicide that is non-selective and is often used to eradicate undesired plants of all kinds, with a focus on weeds [8]. Since the recommended rate of glyphosate applied to the soil is based on the recommended rate for a particular location, the herbicide application must be done appropriately and by the required concentration [9]. Farmers use a lot of the chemical glyphosate to get rid of weeds. The pesticide's hazardous ingredient has the potential to infect *Rhizobium* bacteria, which fix nitrogen in the soil [10]. Additionally, using it in excessive doses can harm microbes, plants, and animals, which eventually deteriorates the food chain and results in nutritional deficiencies.

Particularly vitamins and minerals may also be hazardous to the body as a whole. [10] Glyphosate also has an impact on the functional characterization of plants, where beneficial bacteria are essential for the synthesis of iron and auxin siderophores, as well as for solubilizing zinc and phosphate uptake. [10] Glyphosate has a major effect on soil biology

and is poisonous to beneficial species such as earthworms and microflora. Herbicides containing glyphosate penetrate the soil through the surface of the soil or seep into it, influencing the populations of rhizobia [11]. Although the effect varies throughout species, there are multiple findings suggesting glyphosate herbicides can lower *Rhizobium* species populations [12]. Numerous studies have demonstrated that the nitrogen fixation process is adversely affected by nonselective glyphosate spraying, as it reduces the amount of rhizobia.

Antibiotics are small bioactive molecules that are naturally produced by microorganisms like bacteria and fungi during their secondary metabolism [13,14]. Antibiotic resistance is an ancient and naturally occurring phenomenon widespread in the environment. The rhizosphere contains a mixture of metabolically active microbial populations that compete in this environment about size, diversity, and biochemical activity. Production of antibiotics by some soil harbouring microorganisms, mainly bacteria and fungi, has been largely documented [15].

2. Materials and Methods

2.1 Extracting Rhizobial Bacteria from Root Nodules: According to Vincent [16], local rhizobial bacteria were identified from nodules produced on the hair roots of plants growing in several places in Nineveh Governorate/Iraq. Isolates began to be collected at the beginning of September for two months, and the plants were three to four weeks old. Leguminous plants come in six varieties: phaseolus, cowpea, alfalfa, soybeans, and clover. Rhizobia bacteria have a specialized symbiotic relationship with leguminous plants, as bacteria are only associated with a specific type of legume plants. The plant roots developed nodules and were washed with distilled water. After being immersed in 70% ethyl alcohol for three minutes, they were thoroughly cleaned three times with sterile distilled water. Subsequently, they were submerged in a sodium hypochlorite NaOCL solution with a 3:1 v/v water/sodium hypochlorite concentration.

After being submerged in it for fifteen minutes, the sample was thoroughly cleaned three times using sterile distilled water. It was then moved to a Petri plate with sterile filter paper inside to remove any remaining water from the nodes. Nodules were transferred to Yeast Extract Mannitol Agar (YEMA) and incubated for a duration of 24 to 48 hours at 28°C to guarantee sterilizing effectiveness. Using a sterile glass rod, sterilized nodules were crushed and then arranged in stripes on a Petri plate filled with YEMA media. The dish was then incubated for 24 to 48 hours at 28°C.

2.2 Cultural diagnosis of isolates from the area: Under the oil lens (x100) of a compound light microscope (Figure 1), a growth sample of the colonies was inspected for colony appearance and tested for Gram stain using a Stain Gram kit [17].




Figure 1: Isolated rhizobia colonies and cells

A: Colonies of rhizobia bacteria isolated from SM35
B: a smear of bacteria isolated from the root nodules of SM28, stained with gram stain under the oil lens

2.3 Tests using biochemistry to isolate *Rhizobium* bacteria: Bacterial strains obtained from the root nodules of the chosen plants were subjected to biochemical assays. The assays included Voges-Proskaur [18], Bromothymol blue [19], Congo red [16], motility [20, 21], urease, methyl red, starch degradation, catalase, Kovacs (indole) and citrate utilization [17].

2.4 Tolerance of the isolates to salt: By transferring a single culture of young rhizobial bacteria and spreading them out by plotting on YEMA medium that contains different concentrations of potassium nitrate KNO_3 as follows: 2, 4, 6, 8, and 10%, and then incubating at 28°C for a while of 24–48 hours, this test was carried out to determine the ability of rhizobial isolates to grow in the presence of different concentrations of potassium nitrate KNO_3 [22].

2.5 Testing for antibiotic resistance in rhizobial isolates: To determine the growth of rhizobial isolates in the presence of the following antibiotics ($\mu\text{g/ml}$): Ampicillin, 25, Rifampin, 5, Tetracycline, 10, and Neomycin, 10. These antibiotics were produced by the Turkey Bioanalyase Company.. Using a cotton swab, tested rhizobial isolates were distributed over Mueller-Hinton agar (M.H.A.) medium before the antibiotic discs were moved in. Using sterile forceps, vital cells were extracted, spread out on a medium, and cultured for twenty-four hours at 28°C . The presence of a clear halo surrounding the antibiotic tablets was used to identify isolates of rhizobial bacteria that were sensitive and resistant to different antibiotics [23, 24].

2.6 Glyphosate's Impact on the Rhizobial Isolates' Growth: The purpose of the experiment was to show how tolerant rhizobial isolates were to glyphosate, a pesticide. YEMA medium was used to create the calculated amount of herbicide at concentrations of 0, 5, 25, and 100 mg/ml . Sterile Petri dishes were filled with the medium. Herbicide-free YEMA media was made, transferred onto sterile Petri plates, and regarded as a control. The herbicide-containing Petri dishes were used to cultivate the bacteria. For two days, plates

were kept in the incubator at 28°C. The efficiency of the herbicide against rhizobial isolates was assessed by its effect on growth [25].

3. Results:

3.1 Analyzing culture and conducting biochemical testing: Following the separation of the bacteria from the plant's root nodules, convex, mucoid, semitransparent, and white colonies with smooth edges began to grow on the solid YEM medium. Under a light microscope, rhizobial cells were examined. Negative, bacilli-shaped Gram bacilli with no spores were found. **Table 1** provides a summary of the biochemical test results. The findings showed that the Vogus-Proskaour and methyl red tests yielded negative results for every isolated bacterium. On the other hand, positive results were obtained from the motility, urease, catalase, indole starch, Congo red, and bromothymol blue tests **Figure 2**.

Table 1: Biochemical tests for local rhizobial isolates.

	SM 10	SM13	SM28	SM2 9	SM3 5	SM4 2
Methyl red	-	-	-	-	-	-
Urease	+	+	+	+	+	+
Catalase	+	+	+	+	+	+
Indole	+	+	+	+	+	+
Starch	+	+	+	+	+	+
Congo red	+	+	+	+	+	+
BTB	+	+	+	+	+	+
V-P	-	-	-	-	-	-
Citrate	+	+	+	+	+	+
Motility	+	+	+	+	+	+

BTB: Bromothymol blue; V-P: Voges-Proskaur




Figure 2: Biochemical test.

A- Methyl red test, B- Urease test, C- Catalase test, D- Indole test, E- Starch test, F- Congo red test, G- BTB test, H- V-P test, I- Citrate test, J- Motility test.

The KNO₃ tolerance results are displayed in **Table 3**. Different rhizobial isolates responded differently to this stress in the YEMA medium with varying KNO₃ concentrations. Except for isolates SM35 and SM42, which displayed a decline in growth at a concentration of 6 and

10% (w/v), rhizobial isolates had very robust growth at KNO₃ concentrations of 2, 4, 6, and 8% (W/V). Two of the six rhizobial isolates, SM10 and SM28, demonstrated robust growth at a concentration of 10% (w/v).

Table 2: Effect of potassium nitrate on growth of Rhizobium

Isolate No.	KNO ₃ Concentrations % (w/v)				
	2	4	6	8	10
SM10	+++	+++	+++	+++	++
SM13	+++	+++	+++	+++	+
SM28	+++	+++	+++	++	++
SM29	+++	+++	+++	+++	+
SM35	+++	+++	++	+	+
SM42	+++	+++	++	+	+

(+++) very good growth, (++) good growth, (+) medium growth, (-) No growth

About the antibiotic sensitivity test, isolate SM42 demonstrated inhibition zones 13, 9, 13, and 11 mm to ampicillin (25µg/ml), rifampin (5µg/ml), tetracycline (10µg/ml), and neomycin (10µg/ml), respectively. All other isolates demonstrated resistance to ampicillin (25µg/ml). Tetracycline (10µg/ml) and Neomycin (10µg/ml) did not affect the isolates SM28 and SM29. While the remaining isolates displayed varying levels of sensitivity to various antibiotics, strain SM35 demonstrated resistance to Rifampin (5 µg/ml), isolates SM29 and SM35 were more susceptible to antibiotics, while isolate SM42 was more sensitive to these antibiotics

Figure 3.

Table 3: Local rhizobial isolates' susceptibilities and resistance to the drugs under study.

Isolate No.	Antibiotic concentrations µg/ml			
	Ampicillin 25	Rifampin 5	Tetracycline 10	Neomycin 10
SM 10	R	13*	15	12
SM 13	R	10	20	10
SM 28	R	10	13	R
SM 29	R	18	R	11
SM 35	R	R	17	13
SM 42	13	9	13	11

*: Millimeters (mm) represent the inhibition zone; R stands for resistant.




Figure 3: shows antibiotic testing on rhizobia isolates.
A-SM10, B-SM35 and C-SM42.

Table 5 displays the findings of the rhizobial isolates' tolerance to various glyphosate concentrations. Following a 72-hour incubation period, rhizobial isolates demonstrated their capacity to develop at glyphosate doses of 0, 25, and 50 mg/ml (Figure 4). On YEMA plates treated with Glyphosate at a concentration of 100 mg/mL, none of the rhizobial isolates grew.

Table 4: Effect of herbicide Glyphosate on rhizobial isolates

Isolate No.	Concentration of Glyphosate mg/ml			
	0	25	50	100
SM10	+++	+++	+++	-
SM13	+++	+++	+++	-
SM28	+++	+++	+++	-
SM29	+++	+++	+++	-
SM35	+++	+++	+++	-
SM42	+++	+++	+++	-

(+++) very good growth, (++) good growth, (+) medium growth, (-) No growth




Figure 4 shows the growth of the following rhizobial isolates after two days of incubation on YMA medium supplemented with glyphosate herbicide (mg/ml): A-25, B-50, and C-100.

4. Discussion:

Table 5 lists the isolated local strains, the names of leguminous plants in Table 5 were based on the scientific names according to plant taxonomists. The isolate numbers are the name of the model that the researchers chose for this isolate.

Table (5): Rhizobial isolates, plant sources, and rhizobial species occupied its nodules.

Isolate No.	Plant source	Genus and species of rhizobia
-------------	--------------	-------------------------------

SM10	<i>Vicia faba</i> L.	<i>Rhizobium leguminosarum</i> bv. <i>viciae</i>
SM13	<i>Vigna unguiculata</i> L.	<i>Ensifer ferdii</i> bv. <i>ferdii</i>
SM28	<i>Medicago sativa</i> L.	<i>Ensifer meliloti</i>
SM29	<i>Glycine max</i> L.	<i>R. japonicum</i>
SM35	<i>Trifolium alexandrinum</i> L.	<i>R. leguminosarum</i> bv. <i>trifolii</i>
SM42	<i>Phaseolus vulgaris</i> L.	<i>R. leguminosarum</i> bv. <i>phaseole</i>

Rhizobial isolates from the root nodules of leguminous plants used in this investigation had morphological traits common to Rhizobial genera. These isolates' traits included being bacilli and Gram stain-negative [17, 26, 27]. The tests for urease, catalase, indole, starch hydrolysis, motility, citrate consumption, Congo dye pick-up, and BTB were likewise positive for all isolates. In contrast, the isolates' findings from the Voges-Proskauer and methyl red tests were negative. According to this study, there were some variations in the biochemical examination of the rhizobial isolates compared to previous investigations [28]. Additionally, the results of the capacity of local rhizobial isolates to grow at various KNO₃ concentrations varied. This outcome is consistent with the findings of Bed and Naglot [22], who were able to identify *Rhizobium leguminosarum* bv. *trifolii*.

Variations in antibiotic sensitivity or resistance were shown by the test results. Results of this study were consistent with other researchers' results regarding resistance to Ampicillin, and sensitivity to Rifampin, Tetracycline, and Neomycin [29, 30, 31, 32]. There are many mechanisms in the resistance of rhizobial isolates to antibiotics and these mechanisms were considered to be active. Rhizobial bacteria have specific protein carriers on the outer membrane. Another mechanism is a modification of biosynthetic pathways inside rhizobial bacteria as well as the production of enzymes acting against the target which is an antibiotic. Permeability of the cell wall of rhizobial bacteria is another defined choice for rhizobia to avoid antibiotic effects [33]. Studying of resistance and sensitivity of rhizobial isolates to antibiotics has double benefits in genetic and academic studies [34]. Secondly, such studies provide insight into how rhizobia as a non-pathogenic bacteria, becomes antibiotic-resistant in soil. The explanation for this statement is that rhizobia can receive R plasmids from different bacterial species through a mechanism known as horizontal gene transfer which occurs between pathogenic and non-pathogenic bacterial conjugation [35].

The herbicide experiment performed on various rhizobium isolates also revealed that while the isolates were sensitive to the herbicide and displayed growth inhibition at a concentration of 100 mg/ml, all isolates demonstrated their ability to tolerate a concentration of 25 and 50 mg/ml of the herbicide [16]. The isolates grew best at doses of 25 and 50 mg/ml, with 100 mg/ml showing the greatest growth effect [22]. Therefore, the impact of the herbicide, its

concentration, and the local meteorological circumstances all affected the proliferation of rhizobial bacteria and their capacity to fix nitrogen [36].

5. Conclusion:

Different processes enable rhizobial bacteria to withstand high salt concentrations and pesticides. Poison-resistant plasmids and the production of efflux pumps are two examples of these methods. To completely comprehend the mechanisms underlying *Rhizobium*'s tolerance capacities and investigate the possible uses of these bacteria under these circumstances, more research is required.

Acknowledgement:

The researchers acknowledge the Department of Biology, College of Education for Pure Sciences, University of Mosul, Mosul, Iraq, for its valuable contribution to the enhancement of the quality of this research effort.

6. References:

- [1] Souza, M. C. O., Cruz, J. C., Cesila, C. A., Gonzalez, N., Rocha, B. A., Adeyemi, J. A., Nadal, M., Domingo, J. L., and Barbosa, F. Recent trends in pesticides in crops: A critical review of the duality of risks and benefits and the Brazilian legislation issue. *Environ. Res.* 2023; 115811.
- [2] Padgett, S. R., Re, D. B., Barry, G. F., Eichholtz, D. E., Xavier, D., Fuchs, R. L., Kishore, G. M., and Fraley, R. T. New weed control opportunities: development of soybeans with the Roundup Ready TM gene. In *Herbicide-Resistant Crops*, 53–84. (CRC Press, 2018).
- [3] Stalker, D. M., Kiser, J. A., Greg, B., Bruce, C., and Houck, C. M. Cotton weed control using the BXNTM system. In *Herbicide-Resistant Crops*, 93–106. (CRC Press, 2018).
- [4] Tudi, M. et al. Agriculture development, pesticide application, and its impact on the environment. *Int. J. Environ. Res. Public Health.* 2021;18 (3): 1112.
- [5] Pimentel, D. Amounts of pesticides reaching target pests: Environmental impacts and ethics. *J. Agric. Environ. Ethics.* 1995; 8: 17–29.
- [6] Richmond, M. E. Glyphosate: A Review of its Global Use, Environmental Impact, and Potential Health Effects on Humans and Other Species *J. Environ. Stud. Sci.* 2018; 8: 416-434.
- [7] Ruuskanen, S., Fuchs, B., Nissinen, R., Puigbò, P., Rainio, M., Saikkonen, K., Helander, M. Ecosystem consequences of herbicides: the role of the microbiome *Trends Ecol. Evol.* 2023
- [8] Harries, M., Flower, K. C., Scanlan, C. A., Rose, M. T., and Renton, M. Interactions between crop sequences, weed populations, and herbicide use in Western Australian broadacre farms: Findings of a six-year survey. *Crop Pasture Sci.* 2020; 71(5): 491–505.
- [9] Badowski, M. & Sadowski, J. Poziom pozostalosci 2, 4-Dw roslinach trwalych uzytkow zielonych. *Prog. Plant Prot.* 2008; 48(4): 1185–1189.

- [10] Singh, S. et al., Herbicide glyphosate: Toxicity and Microbial Degradation. Int. J. Environ. Res. Public Health 2020; 17(20): 7519.
- [11] Sharma, S., Kumar, S., Kumar, V., and Sharma, R. Pesticides and vegetables: Ecological and metabolic fate with their field and food significance. Int. J. Environ. Sci. Technol. 2021; 1–26.
- [12] Drouin, P., Sellami, M., Prevost, D., and Fortin, J. Antoun, H. Tolerance to agricultural pesticides of strains belonging to four genera of Rhizobiaceae. J. Environ. Sci. Health Part B. 2010; 45(8): 757–765.
- [13] O'Brien, J.; Wright, G.D. An ecological perspective of microbial secondary metabolism. Curr. Opin. Biotechnol. 2011, 22, 552–558.
- [14] Okada, B.K., Seyedsayamdost, M.R. Antibiotic dialogues: Induction of silent biosynthetic gene clusters by exogenous small molecules. FEMS Microbiol. Rev. 2017, 41, 19–33.
- [15] Shahriar, A., Kobra, A. T., Shomi, F. Y., Emran, T. B., Mallick, J., & Dutta, M. Presumptive correlation between phenotypic, genotypic and symbiotic diversities with antibiotic susceptibility traits of rhizobial strains from plant legumes. environment, 2020, 18, 19.
- [16] Vincent, J. M. A. Manual for the Practical Study of the Root Nodule Bacteria. I. B. P. Handbook No. 15 Oxford: Blackwell Scientific Publication, Oxford, 1970, p. 113–131.
- [17] Baghel, D., Gupta, S.B., & Chowdhury, T. Characterization and Evaluation of Native Rhizobium of Groundnut (*Arachis hypogaea* L.) and Soybean (*Glycine max* L.). Journal of Experimental Agriculture International, 2024, 46.5: 500-506.
- [18] Mahon, C. R., Lehman, D. C., and Manuselis, G. Textbook of Diagnostic Microbiology-e-Book. Elsevier Health Sciences 2018.
- [19] Pervin, S., Jannat B., Sanjee, S., and Farzana, T. Characterization of *rhizobia* from the root nodule and rhizosphere of *Lablab purpureus* and *Vigna sinensis* in Bangladesh. Turkish J. of Agri-Food Sci. Tech. 2017; 5(1): 14–17.
- [20] Collee, J. G., Marmion, B. P., Fraser, A. G., Simmons, A. Mackie, and McCartney, Practical Medical Microbiology. 14th ed., Chrchill Livingstone, New York, U.S.A. 1996, p. 263–98.
- [21] Harley, J. P., & Prescott, L. M., "Laboratory Exercises in Microbiology," 5th ed., WCB/McGraw-Hill, U.K. 2002.
- [22] Bed, M.K., and Naglot, A. Characterization of *Rhizobium* isolated from root nodules of *Trifolium alexandrinum*. J. Agri. Tech., 2011, 7(6): 1705–1723.
- [23] Ramaroson, M., Guillou, S., Rossero, A., Rezé, S., Anthoine, V., Moriceau, N., Zagorec, M. Selection procedure of bioprotective cultures for their combined use with high pressure processing to control spore-forming bacteria in cooked ham. Int. J. Food Microbiol. 2018; 276: 28–38.
- [24] Weinstein, M. P., Patel, J. B., Bobenchik, A., Campeau, S., Cullen, S. K., and Gallas, M. F. Clinical and laboratory standards institute performance standards for antimicrobial susceptibility testing. 2019; 88–9.

- [25] Shankar, P.V., Shaikh, N.R., and Vishwas, P.S. Effect of Different Herbicides on the Nodulation Property of Rhizobial Isolates. *Uni J. Environ Res. and Tech.* 2012; 2:293-299.
- [26] Hewedy, O. A., Eissa, R. A., Elzanaty, A. M., Nagaty, H. H., and Abd Elbary, M. I. Phenotypic and genotypic diversity of Rhizobia nodulating Faba beans from various Egyptian locations. *J. Bioprocess. Biotech.* 2014; 4(5): 1.
- [27] Niste, M., Vidican, R., Puia, C., Rotar, I., and Pop, R. Isolation and biochemical characterization of *Rhizobium leguminosarum* bv. *trifolii* and *Sinorhizobium meliloti* using API 20 NE and API 20 E. *Bull. USAMV Ser. Agric.* 2015; 72:173.
- [28] Bhargava, Y., Murthy, J. S. R., Rajesh Kumar, T. V., and Narayana Rao, M. Phenotypic, stress tolerance, and plant growth-promoting characteristics of rhizobial isolates from selected wild legumes of the semiarid region of Tirupati, India. *Adv. Microbiol.* 2016; 6(1):1–12.
- [29] Mihaylova, S.; Genov, N. and Moore, E. Susceptibility of environmental strains of *Rhizobium radiobacter* to antimicrobial agents. *World Appl. Sci. J.*, 2014; 31(5): 859-862.
- [30] Dhull, S.; Singh, K. and Gera, R. Intrinsic antibiotic resistance (IRA) of different rhizobial strains isolated from root nodules of *Cyamopsis tetragonoloba* L. Taub. *Chem. Sci. Rev. Lett.*, 2018; 6(21): 88-93.
- [31] Khalid, R.; Zhang, X.X.; Hayat, R. and Ahmed, M. Molecular characteristics of rhizobia isolated from *Arachis hypogaea* grown under stress environ. *Sustainability*, 2020; 12(15): 6259-6267.
- [32] Abdell-Hakim, M.M.; Dakhly, O.F and Sameh, A.M. Characterization of some plant growth promoting rhizobacteria isolated from sugar beet rhizosphere. *J. Mod. Res.*, 2022, 4: 6-13.
- [33] Naamala, J.; Jaiswal, S.K. and Dakora, F.D. Antibiotics resistance in *Rhizobium*: type, process, mechanism and benefit for agriculture. *Curr. Microbiol.*, 2016; 72(6): 804-816.
- [34] Prasad, C.K.; Vineetha, K.E.; Hassani, R. and Randhawa, G.S. Isolation and symbiotic characterization of aromatic amino acid auxotroph of *Sinorhizobium meliloti*. 2000. *Indian J. Exp. Biol.*, 83: 1041-1049.
- [35] Macedo, G.; Olesen, A.K.; Maccario, L.; Leal L.H.; Maas, D.H.P.; Heederik, D. and Schmitt, H. Horizontal gene transfer of an IncP1 plasmid to soil bacterial community introduced by *Escherichia coli* through manure amendment in soil microcosms. *Environ. Sci. Technol.*, 2022, 56(16): 11398-11408.
- [36] Sprout, S. L., Nelson, L., and Germida, J.J. Influence of metribuzin on the *Rhizobium leguminosarum* lentil (*Lens culinaris*) symbiosis. *Can J. Microbiol.*, 1992; 38:343–349.