

**Effect of Sublethal Concentrations of Cadmium on
some Physiological Alterations in Liza abu Fish**

Dr.Raaed S. Attee.Ph

Department of Animal production

College of Agriculture

Diyala University

Dr.Ph.Majdy F. Majeed

Department of Histology and anatomy

College of Veterinary Medicine - Basrah University

Asst. instructor .Taha K. Khadier

Department of Animal production

College of Agriculture - Diyala University

ABSTRACT :

The aim of this study was to obtain a holistic view of sublethal toxic responses of fish (Liza abu). After exposure to (0.1 & 0.5) mail- Cd in water for 7, 15, 30 days. The results h d that the distribution of Na'b and Ca+2 concentrations of s owe k+ ja jjjs Gills, but not results alteration of K concentration t e g .

Also these motels included chloride cells proliferation and caused the increase of chloride cells number comparison with control. Histological alternation showed hyperplasia and hypertrophy and earning of secondary lamellae.

INTRODUCTION :

The heavy metals such as cadmium are commonly present in fresh water. The natural background concentrations of (Cd) in aquatic environments is generally below 0.1 mail- (2). However through a variety of human activities such as combustion of fossil fuels, paper production, cement production and extractive metallurgy. Levels of cadmium have increased in aquatic ecosystem in number of industrialized areas of world (1), which it is water concentrations of (Cd) may be up to 100 my.! - and this led to increase susceptibility to disease (4). Cd is non- essential element and its biological function is unknown (3), and is toxic to aquatic organisms even at very low concentrations (2). In fish, uptake of Cd occurs across the

gastrointestinal tract and gills where it involved Ca+2 channels (23).

The gills freshwater fish represent the largest part of the total body surface area, and the bronchial epithelium the distinct between the water and the blood is only few micro's (5). This made gills are as sensitive target organ for waterborne pollutants (16), and this organ will allow for many toxicant such as cadmium enter to organism (18). However the exact routes by which metals pass through the gills are unknown (2). In gills they have three types of main cells which are pavement cells, mucous cells , and chloride cells (8), could be involved but attention has focused largely on chloride cells (CCS) because of their putative role in ions transport even though they occupy only about (7%) of total surface area (19). In fact, no study to date has proven conclusively that chloride cells on the gill ion transport cells strong circumstantial evidence favoring this interpretation that they have a higher metabolic rate than other epithelial cells (17).

Histological alternation in fish gills were used as a target organ for environmental chemical (5), so that the histologists investigation represent sensitive tool to detect the toxic effects of environmental pollutant in laboratory experimental (23).

The aim of present study was to detect the levels of Na^+ , K^+ , and Ca^{+2} concentrations of gills and record a number of chloride cells and show the histologists alternation after exposure to two concentrations of cadmium.

MATERIAL AND METHODS:

Liza abu was collected from fish farming station at Marine Science Center in University of Basrah. Fish was acclimated in laboratory for two weeks at 27°C in a large well-aired aquarium.

Fish average weight was 24.9 ± 1.5 grams. The sample size for each experiment was ten fishes. The fish was divided into three groups, the first one control kept in tap water at 27°C, while two and three groups were exposures for 30 days to sublethal concentrations of Cd in water (0.1 and 0.5 mg/L) respectively, after the exposure period, fish sample were removed from aquarium, and gill specimens of each individual were fixed for

two days by buffer formalin (10%) and routinely embedded in paraffin wax for light microscopy, sections were cut at (7mm) and stained with hematoxyline and eosin (H & E). The Na^+ , K^+ , were determined by flame and Ca^{+2} concentrations of gills atomic absorption (20). While chloride cells used the method of by use the left opercula were prepared for numerical chloride cells(20) .

RESULTS:

Exposure to cadmium (0.1 and 0.5 mg/L) during 7, 15, and 30day

resulted the following : gills after exposure to Na^+ , K^+ , and Ca concentration ons of gills various Cd concentrations are presented in table (1 , 2 and 3), furthermore, a significant decrease (P- 0.05) in Sodium and Calcium concentrations treatment groups compared with control group, While no significant difference in potassium concentrations compare with control was observed in gills for fish exposed to cadmium.

Table (4) showed number of Oprecular Chloride cells increased in a dose-dependent way, resulting in significantly more than (P- 0.05) of chloride cells in fish exposed to (0.1

and 0.5 mg.l^{-1}) compared to control fish (table 2).

Histopathological were found in the gills examined, there were some evidence of gills damage in the exposures fish. The microscopic examination revealed hypertrophy of gill filaments and hyperplasia of epithelium surface of respiratory lamellae and inter lamellar filament epithelium.

Exposure time (Days)	Treatments groups		
	control	0.1 mg^{-1}	0.5 mg^{-1}
7	34.5 $\pm$ 2.1 a	30.6 $\pm$ 2.3 a	28.9 $\pm$ 0.85 a
15	33.7 $\pm$ 1.8 a	27.3 $\pm$ 2.8 b	20.8 $\pm$ 2.3 c
30	30.8 $\pm$ 1.5 a	29.4 $\pm$ 1.9 a	25.6 $\pm$ 3.1 b

Table (1):Effect of cadmium on Na^+ regulation in *L. abu* fish

Exposure time (Days)	Treatments groups		
	control	0.1mg ⁻¹	0.5 mg ⁻¹
7	80.4 ±4.5a	70.3 ±2.3 b	68.9 ±7.8 b
15	78.6±6.4 a	74.5 ±6.4 a	64.3 ±8 b
30	81.4±3.2 a	65.9 ±7.1 b	60.7 ±4.9 b

Table (2):Effect of cadmium on Ca⁺² regulation in *L. abu* fish

Exposure time (Days)	Treatments groups		
	control	0.1mg ⁻¹	0.5 mg ⁻¹
7	68.9 ±2.9a	70.1 ±5.5 b	68.9 ±7.8 b
15	64.7±4.1 a	64.9 ±6.2 a	61.7 ±4.7 b
30	67.8±5.0 a	65.8 ±5.0 b	68.9 ±5.8 b

Table (3):Effect of cadmium on K⁺ regulation in *L. abu* fish

Exposure time (Days)	Treatments groups by cadmium		
	control	0.1mg ⁻¹	0.5 mg ⁻¹
7	2.5 ± 0.12 a	3.4 ± 0.41 b	4.5 ± 0.76 c
15	2.1 ± 0.22 a	5.0 ± 0.79 b	4.8± 0.45 b
30	2.4 ± 0.13 a	4.6 ± 0.44 b	5.6 ± 0.87 c

Table (4):Effect of cadmium on number of chloride cells in *L. abu* fish

DISCUSSION :

The result of this study demonstrates that Sublethal Cd exposure of fish *Liza abu* affects several physiological parameters, such as the Na^+ , K^+ , and Ca^{+2} concentrations and number of chloride cells of gills in addition to Histopathological of this organ.

The exposure of Cd included challenge of physiological homeostasis (13). The above mentioned aspects are well illustrated by Na^+ , K^+ , and Ca^{+2} concentrations of gills. The regulation and maintenance of ion-homeostasis is very sensitive to exposure metals in freshwater fish the gills are target during exposure to heavy metals and other stressors, because they play a predominant role in maintenance of ion-homeostasis and the uptake of waterborne heavy metals during increased concentrations in water (21). In this study we observed an increase in number of chloride cells, in freshwater fish chloride cells functions in ions uptake, and therefore, an increase in chloride cells numbers has been considered a compensating response to pollutants included ions-loss (14).

However, environment of endocrine system can not be excluded. Hormone control of bronchial function enable fish to modify the ion's exchanges capacity in the gills in response to metals (21). The release of cortisol, produced by internal tissue located in head kidneys of fish is stimulated as part of stress response (11). The hormone is known to affect ionic regulation and to include chloride cells proliferation (16). However, in spite of an increase in chloride cells number in groups exposure to metal, this response was not adequate to oppose these ion's losses (13). Exposure to Cd resulted inhibition of the active Na^+ , K^+ , and Ca^{+2} transport activity (22), and Na^+ enzymes these metals induced inhibition of Ca^{+2} and activity which it's related to the free heavy metals ions available (24). Bronchial Na^+ and Ca^{+2} transport differ from another in many ways including the hormonal control and specific transport sites (11). This also may contribute to the metal and transport enzymes specific effects of Cu and Cd on trans epithelium ion exchange (12). This was in line with

previous observations in *Liza abu* fish where active Na^+ and Ca^{+2} uptake was inhibited in fish exposure to 200 mg-l Cd (13).

The heavy metals have a high affinity for nucleophilic groups, in particular for SH-residues of amino acids and proteins, and heavy metals interaction with ion-ATP ase has mainly been attributed to the affinity of the metals for Sulfhydryl groups on the ion transporting enzyme (6).

Alteration of gills filaments and hyperplasia of epithelial surface of

respiratory lamellae and interlamellar filament epithelium. Some authors suggested that respiratory system of teleost fish maybe damaged by cadmium, since acute concentration of cadmium was found to cauterize the gill lamellae of several freshwater fish (7). What we study has been agreed with the study of(1) and. Which recorded pathological alternations in gill filament and respiratory lamellae in (cyprinoid carpio). However, the morphological effects of metals exposure have been comprehensively reviewed (1%, and fall into two broad categories :Accumulated damage and compensatory responses at moderate level of exposure, the damage consists of separation of epithelial layers, tissue edema and clubbing of lamellae. While at more several levels, tissue necrosis and rupture and fusion of secondary lamellae become more prominent. The later compensatory responses include hypertrophy and hyperplasia of mucous cells and chloride cells which appear to be associated with a pair of gill damage, probably play an important role in recovery and acclimation.

**تأثير التراكيز تحت المميتة لعنصر الكاديوم على بعض التغيرات الفسيولوجية
لأسماك الخشني LIZA ABU**

أ.م.د. مجدي فيصل مجيد	أ.م.د. رائد سامي عاني
قسم التشريح والانسجة	قسم الثروة الحيوانية
كلية الطب البيطري /جامعة البصرة	كلية الزراعة /جامعة ديالى
م.م. طه خاطر خضير	
قسم الثروة الحيوانية	
كلية الزراعة / جامعة ديالى	

المستخلص

اجريت الدراسة الحالية على صغار الاسماك LIZA ABU لتحديد تأثيرات التراكيز تحت المميتة من عنصر الكاديوم (٠.١, ٠.٥ ملغم /لتر) على التغيرات الفسلجية الحاصلة للتبادل الايوني لكل من NA^{+} , K^{+} , CA^{+2} بعد فترات تعريض ٧ و ١٥ و ٣٠ يوم اضافة لدراسة التشوهات الحاصلة لانسجة الغلاصم خلال هذه الفترة من التعريض اظهرت النتائج اخفاضات معنوية في مستوى التراكيز NA^{+} , CA^{+2} لغلاصم الاسماك (٠.٠٥ P) في حين لم يسجل عنصر K^{+} أي تغيرات خلال هذه الفترة فيما أظهرت اسجة هذا العضو حالات الفرط النسيجي و التنخرات الواضحة.

BIBLOGMPHY :

- 1 . Al-sudani, A. (1999) . The effects of sublethal concentrations of Cd on survival rates and some physiological aspects of common carp cyprinoid carnie. A thesis, College of agriculture, Basrah-Iraq, put .
2. Bewers, J.M., Barry, P.J., and McGregor, D.J. (1987). Distribution and cycling of cadmium in the environment. In cadmium in aquatic environment. Eds. Nriagu, J.O. and Sprague, J.B. Wiley Interscience, New York, pap. 1-18 .
3. Chmielnicka, J. and Cherian, M.G. (1986). Environmental exposure to (Cd land Factors affecting trace-element metabolism and metal bolide, Biol. Trace. Elem. Res. 10, 243- 262 .
4. Frieden, E. (1979). Ceruloplasmain : The serum copper transport protein with oxidize activity. In Copper in the environment, Part 11 . Hea1th effects. Eds. Nriagu, J.O. and Sprague, J.B. Wiley Interscience, New York, pap. 241-284 .
5. Hinton (1992). Biomarkers - Bio chemical, physiological and Histological marker of anthropogenic stress (Ed. R-J. Hugett ; R.A. Kimerlr ; P.M. Mehrle ; H.J. Bergman), Lewis publishers, Chelsea, MI, USA, p.115 .
6. Jacobson, K.B. and Turner, J.E. (1980). The introduction and certain other metals ions with protein and nucleic acids. Toxicology 16, 1-37.
7. Karlsson, L., Runn, P. and Hanx, C. and Porlin, L. (1985).

Cadmium induced changes in morphological of Zebra fish, *Brachydanio rerio* and rainbow trout *Salmo gairdneri*. J.

Fish. Biol. 27: 81-95.

8. Lauren, P. (1984). Gills internal morphology, p73-183. In W.S. Hoar and D.J. Randall (ed) Fish physiology. Vol IOA. Academic Press, New York (by.

9. Lyndon, A.R. (1994). A method for measuring oxygen consumption in isolated perfused gills. J. Fish. Biol. 44:707-715. '

10. Mallatt J. (1985). Fish Gill structure changes induced by toxicant and other irritants : a statistical review. Can. J. fish. Aquat. Tech. 5, 1 138-9.

11: McDonald, D.G. and Wood, C.M. (1993). Bronchial mechanisms of acclimation to metals in freshwater fish. In : Fish ecophysiology. Fish and Fisheries series 9. Eds. : Rankin, J.C. and Jensen, F.B., Chapman and Hall, London, pap. 297-321.

12: Niebor, E. and Richardson, D.H.S. (1980). The replacement of the non-descript term heavy metals ions. (Environ . Pollute. IB, 3-26).

13: Spelgrom, S.M.G.J., Lamers, L.P.M., Lock, R.A.C., Balm, P.H.M., and Wendelaar Bonga, S.E. (1995). Interaction between copper and cadmium modify metal organ distribution in Mature tilapia, *Orachromis mossambicus*. Environ. Poll. (in press).

14. Perry, S.F. ; Goss, G.G. and Laurent, P. (1992). The interrelationship between gill chloride cell morphology and ionic uptake in four freshwater teleosts. Can. J. 2001. 70, 1775-1786.

15: Perry, S.F. and Laurent, P. (1993). Environmental effects on fish gill structure. In : Fish ecophysiology . Eds. : Rankin, J.C. and Jensen, F.B. Chapman and hall, London, pap. 231-264.

16. Perry, S.F. and Laurent P. (1993). Environmental effects on fish gill,

structure and function. 1n: Fish ecophysiology. Eds : Rankin, J.C. and F.B. Jensen. Chapman and Hall, London, pap. 23 1-264.

1 7. Perry, S.F. and Walsh, R.J. (1989). Metabolism of isolated fish gill cells : contribution of epithelial chloride cells. J. Exp. Biol. 144:507-520.

1 8. Playle, R.C., Gensmer, R.W. and Dixon, D.G. (1992). Copper accumulation on gills of fathead minnows : influence of water hardness, complication and pl-! of the gill micro-environment. Environm. Toxicol. Chem. l 1, 381-391.

19. Potts, W.T.W. (1977). Fish gills, p. 453-480. In B.L Gupta, R. Moreton, J. Oschman, and B. Wall (ed.) Transport of ions and water in animals. Academic press, New York, NY.

20. R.O.P.M.E. (1983). Manual of Oceanographic Observation and Pollutant analysis Methods.

21. Stagg, R.M., and Shuttleworth, T.J. (1982). The accumulation of copper in platichthys Jesus L. and its effect on plasma electrolyte concentrations. J. Fish. Bill., 209: 491-500.

22. Tokushiga, A. Higashino, H. and Tamura, H.)1984). Cd effect on the Na, K system in cultured vascular smooth muscle cells Hypertension 6, 20-26.

23. Verbost, P.M. (1989). Cadmium toxicity : interaction of cadmium with cellular calcium, sodium and potassium transport mechanisms. Thesis university of Nijmegen. Netherlands.

24. Verbost, P.M. ; Van Rooji, J. ; Flick, G. ; Lock, R.A.C. ; Wendelaar Bonga, S.E. (1989). The Movement of cadmium through freshwater trout bronchial epithelium and its interference with calcium tz|mlspolt J. Exp. Biol. 145, 185-197.