

Synthesis, Characterization of some New 4-Thiazolidinone of Acenaphthoquinone

Muntaha .Y. Hayal
Dept. of Chemistry/ Science College/ University of Thi-Qar
Iraq


Mahmood Shakir Magtoof
Dept. of Chemistry/ College of Science / University of Thi-Qar
Iraq
Mahmood672000@yahoo.com

Abstract— This work summarized with Synthesis of schiff's bases 1(a-c) from reaction of parent compound acenaphthoquinone with primary amines . Treating of imines compounds with thioglycolic acid was produced thiazolidinone derivatives 2 (a-c) .The structures of synthesized compounds were confirmed by using some spectroscopic analysis such as FT.IR , ¹H-NMR and ¹³C-NMR spectral

Keywords— thioglycolic acid ; ¹³C-NMR spectra ; Acenaphthenquinone; thiazolidin

I. INTRODUCTION

Thiazolidined-4-one are the derivatives of thiazolidine which belong to an important group of heterocyclic compounds containing sulfur and nitrogen in a five member ring ⁽¹⁾ 1,3-Thiazolidin-4-ones are heterocyclic at position 1 containing a sulfur atom , position 3 nitrogen and position 4 carbonyl group ⁽²⁾ figure (1) .Thiazolidine and its composites are key components of many natural products and drugs . In addition, Thiazolidine found to uses as antitubercular ⁽³⁾, antibacterial ⁽⁴⁾, anti-inflammatory ⁽⁵⁾, as antiviral agents, especially as anti-HIV agents ⁽⁶⁾, anticancer ^(7,8), anticonvulsant ⁽⁹⁾ and antidiabetic activity ⁽¹⁰⁾.



Figure(1): Thiazolidinone ring

II. EXPERIMENTAL PART

A. General procedure for the preparation of imines 1(a-c)

In general ⁽¹¹⁾, imines prepared by mixed the amine with acenaphthoquinone in (25 ml) of Suitable solvent ethanol and add (10 -15) drops of glacial acetic acid were refluxed in water bath for (24 – 25) h .The reaction was followed up with TLC the eluent using [Hexane :Ethyl acetate (7:3)]. When the reaction completion the solvent removed by evaporation then the product recrystallized by methanol.

B. (1Z,2E)-N,N'-bis(4-methylphenyl)-2a,5-dihydroacenaphthylene-1,2-diimine (1a)

Prepared by reaction Acenaphthenquinone (1g, 5.489 mmol) with 4-methylaniline (1.176 g , 10.978 mmol) m.p = 220-222 R_f = 0.9 IR (KBr disk): (1629. 85) cm⁻¹ (C=N) yield = 78% .

C. (1Z,2E)-N,N'-bis(4-(dimethylamino) phenyl)-2a,5-dihydro acenaphthylene-1,2-diimine (1b)

Prepared by reaction Acenaphthenquinone (1g, 5.489 mmol) with N,N-dimethylbenzene-1,4-diamine (1.495 g , 10.978 mmol) m.p = 246-248 R_f = 0.7 IR (KBr disk): (C=N) 1654. 92 cm⁻¹ yield = 76 % .

D. (2Z)-2-[(4-bromophenyl)imino]-6,8a-dihydroacenaphthylene-1(2H)-one (1c)

Prepared by reaction Acenaphthenquinone (1g, 5.489 mmol) with 4- bromoaniline (0.94g , 5.489 mmol) m.p = 238-240 R_f = 0.8 IR (KBr disk): 1654. 92cm⁻¹ (C=N) yield = 72 % .

E. General Procedure of thiazolidinones 2(a-c)

In general, ⁽¹²⁾ the thiazolidinones were mixed with imines 1(a-c) and thioglycolic acid in (15 ml) chloroform , Then refluxed for (18-24) h with Stirring. The reaction monitored by TLC using eluent [n-Hexane-Ethyl acetate (3:7)] . When The reaction completed the solvent removed to give thiazolidinone. The product was precipitated and recrystallized with the addition of methanol droplets.

F. N-(4-methylphenyl)-4'-spiro[acenaphthylene-2,2'-thiazolidine] (2a)

Prepared by reacting (1g, 2.778 mmole) (2E)-2-[(4-methylphenyl) imino]-1,2-diphenyl ethanone (1a) and (0.674mL, 0.447g, 5. 556 mmole) of thioglycolic acid. R_f = 0.7 , yield = 63 % , m.p. =232-234 °C. IR (KBr disk): 1689.64 cm⁻¹ (–N–C=O).

G. *N*-(4--(dimethylamino) phenyl)-4'-spiro[acenaphthylene-2,2'-thiazolidine] (2b)

Prepared by reacting (1g, 3.049 mmole) (1*Z*,2*E*)-*N*, *N*'-bis (4-(dimethylamino) phenyl)-2a,5-dihydroacenaphthylene-1,2-diimine(1b) and (0.354 mL, 0.469 g, 6.098 mmole) of thioglycolic acid. $R_f = 0.7$, yield = 60%, m.p. = 255-257°C. IR (KBr disk): 1686.51 cm⁻¹ (–N–C=O)

H. *N*-(4-bromophenyl)-4'-spiro[acenaphthylene-2-thiazolidine] (2c)


prepared by reacting (1g, 2.974 mmole) (2*Z*)-2-[(4-bromophenyl)imino]-6,8a-dihydroacenaphthylene-1(2*H*)-one (1c) and (0.361 mL, 0.272 g, 2.974 mmole) of thioglycolic acid $R_f = 0.8$, yield = 62%, m.p. = 236-238 °C. IR (KBr disk): 1698.85 cm⁻¹ (–N–C=O)

III. MEASUREMENTS

Melting points were determined in open capillary tubes using an electro thermal melting point /SMP31 apparatus. FTIR spectra in the range (200-4000) cm⁻¹ were recorded as KBr discs using a Shimadzu FTIR spectrophotometer. The ¹H-NMR were recorded on VARIAN spectrophotometer (300 MHz), the ¹³C-NMR spectra were recorded using VARIAN spectrophotometer (75 MHz) relative to the internal standard tetramethylsilane (TMS), DMSO-d₆ used as solvent.


IV. RESULT AND DISCUSSION

Prepared Compounds 1(a-c) from the reaction of acenaphthoquinone with 4-methylaniline, *N,N*-dimethylbenzene-1,4-diamine and 4-bromoaniline respectively with the presence of glacial acetic acid in absolute ethanol as shown in scheme (1).



scheme (1): Synthesis of the compounds 1(a-c)

The reaction involves a nucleophilic attack of the amine group on the carbon of the carbonyl group of the ketone to form a compound *N*-(substituted hemiaminals) which loses a water molecule to give the stable compound. Mechanism of imines formation⁽¹³⁾ shown in scheme (2)



scheme(2): Mechanism of imines formation

Measured the melting points Compounds 1(a-c) as shown in Table (1) diagnosed by specifying (FT-IR) shown in Table (2). It features ranges corresponding to the expansion vibrations, azomethine band (C=N), aromatic (C=C), bands aromatic (C-H) and aliphatic (C-H). These bands occur (1629.85, 1654.92, 1654.92), (1587.42, 1602.85, 1598.99), (3051.39, 3047.53, 3055.24), (2916.37, ---, 2835.36) cm⁻¹ respectively.

Table (1) shows the physical properties data imines 1(a-c)


Comp. 1 (a-c)	m.p. °C	Colour	Reaction time
1a	220-222	Orange	24h
1b	246-248	Red	25h
1c	238-240	Yellow	24h

Table (2): shows IR spectra of Imines 1(a-c)

Comp. 1(a-c)	Aromatic C-H stretching cm ⁻¹	Aliphatic C-H stretching cm ⁻¹	Azomethine C=N stretching cm ⁻¹	Aromatic C=C stretching cm ⁻¹
1a	3051.39	2916.37	1629.85	1587.42
1b	3047.53	2875.86	1654.92	1602.85
1c	3055.24	2972.31	1654.92	1598.99


V. SYNTHESIS 4-THIAZOLIDINON

Compounds were 2(a-c) prepared from the reaction of imines 1(a-c) with acid thioglycolic acid in absolute chloroform as shown in scheme (3).



Scheme (3): Synthesis of the compounds 2(a-c)

Involved mechanism cycloaddition⁽¹³⁾ formation 4-Thiazolidinone as shown in Scheme (4)



scheme (4) Mechanism of 4-Thiazolidinone formation

The melting point of the prepared compounds 2(a-c) was measured as shown in Table (3) The (FTIR) spectra Table (4) of compound 2(a-c) show featured packages most notably, C-H aromatic, aromatic C=C, aliphatic C-H and carbonyl amide group which occur within; (3051.39, 3054.83, 3069.35), (1608.63, 1606.28, 1691.57), (2920.23, 2984.09, 1600.77), (1689.64, 1686.51, 1600.77)

Table 3: physical properties of thiazolidinones 2(a-c)

Comp. 2 (a-c)	m.p. °C	Colour	Reaction time
2 a	232-234	Grey	24h
2b	255-257	Yellow	25h
2c	236-238	White	24h

Table (4): FTIR spectral data of Thiazolidinones 2(a-c)

Comp. 2 (a-c)	Aromatic C-H stretching g cm^{-1}	Aromatic C=C stretching cm^{-1}	Aliphatic C-H stretching g cm^{-1}	Amide C=O stretching cm^{-1} (thia-)
2 a	3051.39	1608.63	2920.23	1689.64
2b	3054.83	1606.28	2984.09	1686.51
2c	3069.35	1691.57	1600.77	1698.58

The $^1\text{H-NMR}$ of 2(a-c) shows signals at $\delta(2.39)$ ppm for CH_3 Component (2a) and $\delta(2.51)$ ppm for $\text{N}(\text{CH}_3)_2$ Component (2b) also show characteristic chemical shift (CH_2) group of thiazolidinone ring showed doublet of doublet signal at chemical shift δ [dd-(4.10- 4.05ppm, $J=15\text{Hz}$) - (4.08- 4.03ppm, $J=15\text{Hz}$) - (4.12- 4.07ppm, $J=15\text{Hz}$), and doublet of doublet signal at chemical shift δ [dd-(4.34- 4.29ppm, $J=15\text{Hz}$) - (4.23- 4.18ppm, $J=15\text{Hz}$) - (4.27- 4.22ppm, $J=15\text{Hz}$) respectively. a multiplet signal at δ (6.72-8.01), (6.39-8.29), (6.89-8.28) ppm for aromatic protons respectively as show Table (5).

$^{13}\text{C-NMR}$ spectral of 2(a-c) gives signal at $\delta(20.89)$ ppm for carbon $-\text{CH}_3$ Component (2a), at $\delta(40.11)$ ppm for carbon $-\text{NCH}_3$ Component (2b) and characteristic signal of 2(a-c) of thiazolidinone ring show at [(33.17), (32.92), (33.07) ppm] for carbon $-\text{CH}_2-$, at [(72.98), (73.55), (73.29) ppm] for carbon S-C-N , at [(172.67), (172.55), (172.41) ppm] for carbon $-\text{N-C=O}$ as show Table (6).

Table(5): $^1\text{H-NMR}$ spectral data of Thiazolidinones 2(a-c)

Comp. 2 (a-c)	thiazolidin-4-one ring		Aromatic proton	Aliphatic proton
	C-H ring, J Hz	C-H ring, J Hz		
2 a	4.10-4.05 ppm $J=15\text{Hz}$	4.34-4.29ppm $J=15\text{Hz}$	6.72-8.01	2.39
2b	4.08-4.03 ppm $J=15\text{Hz}$	4.23-4.18ppm $J=15\text{Hz}$	6.39-8.29	2.51
2c	4.12-4.07ppm $J=15\text{Hz}$	4.27-4.22ppm $J=15\text{Hz}$	6.89-8.28	----

Table(6) $^{13}\text{C-NMR}$ spectral data of Thiazolidinones 2(a-c)

Chemical shift ppm					
Comp. 2 (a-c)	$-\text{CH}_2-$	C-N-S	N-C=O	C-Ar	Other
2 a	33.17	72.98	172.67	118.08 - 148.24	$-(\text{CH}_3)$ 20.89
2b	32.92	73.55	172.55	112.37- 150.05	-----
2c	33.07	73.29	172.42	121.72- 140.58	$-(\text{NCH}_3)$ 40.11

REFERENCES

1- Krishnakant T. Waghmode, Journal of Chemical and Pharmaceutical Research, 6(5):1101-1105, (2014).

- 2- Altintas H, Ates O, Birteksoz S, Gulten O, Uzun M, et al. Turk J Chem 29: 425-435, (2005) .
- 3 - Zhou, C.F.; Chen, J.; Liu, P.G.; et al.: Design, Bioorg. Med. Chem.; 18(1):314-9, (2010)..
- 4 - Handan, A.; Oznur, A.; and Seher, B.: Turk.Chem.J.; 29: 425 – 435 , (2005) .
- 5 - Chandra, T.; Garg, N. and Kumar, A., World Journal of Chemistry.; 4 (2): 210-218 , (2009) .
- 6- Rawal, R.K.; Tripathi, R.; Kulkarni, S.; et al ., Chem Biol. Drug Des, 72(2):147 –54 , (2008) .
- 7 - Senkiv, J.; Finiuk, N.; Kaminsky, D.; Havrylyuk, D.; Wojtyra, M.; Kril, I.; Lesyk, R. Eur J Med Chem, 117, 33, (2016) ..
- 8- Szychowski, K. A.; Leja, M. L.; Kaminsky, D. V.; Binduga, U. E.; Pinyazhko, R.; Lesyk, R. B.; Gmiński, J. Chem Biol Interact, 262, 46 , (2017) .
- 9 - Shingalapur, R. V.; Hosamani, K. M.; Keri, R. S.; Hugar, M. H. Eur J Med Chem, 45, 1753 , (2010) .
- 10- Deshmukh, A. R.; Bhosle, M. R.; Khillare, L. D.; Dhumal, S. T.; Mishra, A.; Srivastava, A. K.; Mane, R. A. Res Chem Intermed , 43, 1107, (2016) .
- 11 - Krishnaswamy, D.; Tetrahedron , 34, 4567, (2002).
- 12- Divyesh P., Premlata K., and Navin P., Arch. Appl. Sci. Res., 2(6), 6875, (2010).
- 13- Wang L., Feng Y. , Xue J. and Li Y., J, Serb. Chem. Soc., 73, 1, (2008) .
- 14- Shah J., and Desai A., ARKIVOC, 14, 218, (2007).