

Antimicrobial resistance, Virulence profiles of *Salmonella enterica* serovar Typhimurium isolated from clinical samples

Ezat Hussain Mezal

Department of Biology-College of Sciences- University of Thi-Qar

Abstract:

A total of 33 *Salmonella enterica* serovar (ser.) Typhimurium isolates were isolated from clinical samples. These isolates were subjected to testing and analyzed for antibiotic resistance and virulence genes by using simplex PCR. All isolates were sensitive to gentamycin, kanamycin, nalidixic acid, chloramphenicol, and sulfisoxazol. on other hands the isolates showed intermediate resistance to streptomycin while one isolate (# 22) showed resistance to chloramphenicol and tetracycline. For ampicillin, six isolates were resistance to the drug. 33 of the showed either intermediate or full resistance to one or two of the antimicrobials tested. Most isolates were positive for teen of the virulence genes tested (*msgA*, *tolC*, *spaN*, *invA*, *ipfC*, *sitC*, *sopB*, *orgA*, *pagC* and *pefA*) . For *sitC*, three isolated were negative to this virulence gene .While two isolate were negative to *lpfC* . One isolate # 33 was negative to *orgA* and *spaN*. These results suggest that *S. Typhimurium* from clinical is virulent, and that capable of causing salmonellosis in humans and it may contribute to pathogenesis

Key words: Virulence genes, Antibiotic resistance, *Salmonella Typhimurium*

مقاومة الجراثيم للأدوية وملاحضات الضراوة لبكتريا السالمونيلا نوع Typhimurium المعزولة من العينات السريرية

عزت حسين مزعل

جامعة ذي قار - كلية العلوم - قسم علوم الحياة

الخلاصة:

تضمنت الدراسة عزل ٣٣ نوع من السالمونيلا (*Typhimurium*) من العينات السريرية. تعرض هذه العزلات لاختبار وتحليل للمقاومة وضراوة الجينات للمضادات الحيوية باستخدام البسيط PCR. كانت جميع العزلات حساسة للجنتاميسين، الكاناميسين، حمض الناليديكسيك، الكلورامفينيكول، *sulfisoxazol* من جهة اخرى العزلات أظهرت مقاومة متوسطة للستربتوميسين عدا عزلة واحدة (# ٢٢) وأظهر مقاومة الكلورامفينيكول والنتراسيكلين. أظهرت ستة عزلات مقاومة للامبيسلين . ٣٣ عزلة أظهرت مقاومة إما متوسطة أو كاملة إلى واحد أو اثنين من *anmicrobials*. وكانت معظم العزلات كانت موجبة لعشرة من جينات الضراوة (*msgA*, *tolC*, *spaN*, *InvA*, *ipfC*, *sopB*, *OrgA*, *pagC*, *sitC* و *pefA*). فيما يخص *sitC* , كانت ثلاث عزلات سالبة لهذا الجين. بينما اثنين من العزلات كانت سالبة ل *lpfC*. وعزلة واحدة (٣٣) كانت سالبة ل *orgA* و *spaN* . وتشير هذه النتائج إلى أن السالمونيلا نوع *Typhimurium* هي الأكثر ضراوة والأكثر امراضية في البشر .

1. Introduction

Salmonella are recognized as major food-borne pathogens in humans worldwide usually due to the

consumption of contamination food or water. A variety of foods have been implicated as vehicles transmitting salmonellosis to humans, including poultry, beef, pork,

eggs, fish and vegetables [1,2,3]. Infection by *Salmonella enterica* is significant public health concerns across the globe. *Salmonella* penetrate from the gut lumen into the epithelium of the small intestine, causing acute gastrointestinal illness such as gastroenteritis, organ focal infection, and systemic febrile infection [4]. According to FoodNet which was established in 1996 in collaboration with CDC, USDA, FDA and selected state health departments estimated that 3.6 million (39 %) food borne illness were caused by bacteria in which non-typhoid *Salmonella* has caused about 1, 0267,561 cases of food-borne illnesses, 19, 336 cases of hospitalization [5]. So far more than 2610 serovar of *Salmonella enterica* have been recognized from all over the world, and almost all are able to cause illness in humans and animals [6,3]. Children are prone to an infection caused by *Salmonella*, but infants, elderly, and immunocompromised people are more likely to attract severe infections. *Salmonella enterica* serovar Typhimurium (*S. Typhimurium*) is one of the main serovars causes of human gastroenteritis according to Center for Disease Control and Prevention. Furthermore, the two most frequent serovars *S. Typhimurium* and *S. Enteritidis*. *S. Typhimurium* is one of the main serovars of *Salmonella enterica*. *S. Typhimurium* is predominately found in the intestinal lumen. This serovars gains its toxicity because of the large amount of lipopolysaccharides (LPS) that make up the outer membrane of the bacteria. *S. Typhimurium* is not the most dangerous type of *Salmonella*. Some symptoms caused by *S. Typhimurium* include diarrhea, abdominal cramps, vomiting, and nausea. Strain identification is important for effective investigation of source outbreaks on other hands, molecular tools are necessary for monitoring and prevention diseases. Among molecular -based techniques used recently Polymerase cycle reaction. In this study, *S. Typhimurium* isolates from clinical samples were examined for Antimicrobial resistance and PCR for virulence genes.

2. Materials and methods

2.1 Bacterial strains

Thirty three isolates of *Salmonella enterica* serovar Typhimurium were selected for this study. These strains were of clinical origin and were obtained from the Arkansas Department of Health (USA).

2.2. Antimicrobial susceptibility testing by disk diffusion

All of the *S. Typhimurium* isolates used in this study were tested for resistance to eight antibiotics on Mueller-Hinton agar (Difco Laboratories, Detroit, MI) by a disk diffusion method [7]. The antibiotics that were used: kanamycin (30 µg), streptomycin (10 µg), tetracycline (30 µg), nalidixic acid (30 µg), sulfisoxazol (25 µg), ampicillin (10 µg), chloramphenicol (30µg) and gentamycin (10 µg). Susceptibility and resistance were determined in accordance with the criteria of the Clinical and Laboratory Standards Institute [8]. *Escherichia coli* ATCC 25922 was used for quality control, because it is susceptible to all of these antibiotics.

2.3 PCR detection of virulence genes

S. Typhimurium isolates were screened for 10 virulence genes by the simplex PCR method using single set primers [9]. Primers used for this study are listed in Table 1. The template DNA from the isolates was extracted from overnight cultures by using the DNeasy ® Blood and Tissue kit (Qiagen, Valencia, CA, USA). The PCR reaction mixture with a final volume of 10 µl, contained 2 µl of template DNA, 5 µl of GoTaq Green Master Mix (Promega), 1 µl of each needed forward and reverse primers, and 1 µL of distilled water. PCR cycle conditions were as follows: 5 min for the initial denaturation at 95 °C; 30 cycles of 40 s at 94 °C, 60 s at 66.5 °C, and 90 s at 72 °C, and a final extension of 10 min at 72 °C. The PCR products were visualised by electrophoresis on 1.2 % agarose gels in 1 × TAE buffer at 50 V for 85 min.

Table 1: Primers used in PCR for detection of virulence genes in *S. Typhimurium*

Gene	Sequence of Nucleotides	size (bp)	Function of gene
<i>pagC</i>	F- CGCCTTTCCGTGGGGTATGC R- GAAGCCGTTTATTTTGTAGAGGAGATGTT	454	survival within macrophage
<i>msgA</i>	F- GCCAGGCGCACGCGAAATCATCC R- GCGACCGCCACATATCAGCCTCTTCAAC	189	survival within macrophage
<i>invA</i>	F- CTGGCGGTGGGTTTGTGCTCTCTATT R- AGTTTCTCCCTCTTCATGCGTTACCC	1070	Host recognition /invasion
<i>spaV</i>	F- AAAAGCCGTGGAATCCGTTAGTGAAGT R- CAGCGCTGGGGATTACCGTTTTC	504	Entry into nonphagocytic cells
<i>orgA</i>	F- TTTTGGCAATGCATCAGGGAACA R- GGCGAAAGCGGGGACGGTATT	255	Host recognition /invasion
<i>sitC</i>	F- CAGTATATGCTCAACGCGATGTGGGTCTCC R- CGGGGCGAAAATAAAGGCTGTGATGAAC	768	Iron acquisition
<i>lpgC</i>	F- GCCCGCCTGAAGCCTGTGTTC R- AGGTGCGCCGCTGTTGAGGTGGATA	641	Host recognition /invasion
<i>sopB</i>	F- CGGACCGGCCAGCAACAAACAAGAAG R- TAGTGATGCCGTTATGCGTGAAGTATT	220	Host recognition /invasion
<i>pefA</i>	F- GCGCCGCTCAGCCGAACCAAG R- GCAGCAGAAAGCCAGGAACAGTG	157	Host recognition /invasion
<i>tolC</i>	F- TACCCAGCGCAAAAAGAGGCTATC R- CCGGTTATCCAGTTGTTC	161	Host recognition /invasion

3. Results

Antimicrobial susceptibility testing of study isolates showed that all 33 isolates were sensitive to gentamycin, kanamycin, nalidixic acid, chloramphenicol, and sulfisoxazol. Most of the isolates showed intermediate resistance to streptomycin while one isolate (# 22) showed resistance to chloramphenicol and tetracycline Table 2. For ampicillin, six isolates were resistance to the drug. 33 of the showed either intermediate or full resistance to one or two of the antimicrobials tested. (Table 2). In this study, the size of zone of inhibition of every antibiotic disc was measured in millimeter and while those zones of inhibition compared with zone diameter interpretive standards from CLSI [8]. (Table 2).

Table 2: Antibiotic resistance patterns of isolates

ID	AM-10	C-30	GM-10	K-30	NA-30	SXT-25	S-10	TE-30
1	S	S	S	S	S	S	I	S
2	S	S	S	S	S	S	I	S
3	S	S	S	S	S	S	I	S
4	R	S	S	S	S	S	I	S
5	R	S	S	S	S	S	I	S
6	S	S	S	S	S	S	I	S
7	S	S	S	S	S	S	I	S
8	R	S	S	S	S	S	I	S
9	S	S	S	S	S	S	I	S
10	S	S	S	S	S	S	I	S
11	S	S	S	S	S	S	I	S
12	S	S	S	S	S	S	I	S
13	S	S	S	S	S	S	I	S
14	R	S	S	S	S	S	I	S
15	S	S	S	S	S	S	I	S
16	S	S	S	S	S	S	I	S
17	S	S	S	S	S	S	S	S
18	S	S	S	S	S	S	I	S
19	S	S	S	S	S	S	I	S
20	S	S	S	S	S	S	I	S
21	S	S	S	S	S	S	I	S
22	S	R	S	S	S	S	I	R
23	S	S	S	S	S	S	I	S
24	S	S	S	S	S	S	S	S
25	S	S	S	S	S	S	S	S
26	S	S	S	S	S	S	I	S
27	S	S	S	S	S	S	S	S
28	S	S	S	S	S	S	I	S
29	S	S	S	S	S	S	I	S
30	S	S	S	S	S	S	S	S
31	R	S	S	S	S	S	S	S
32	S	S	S	S	S	S	S	S
33	R	S	S	S	S	S	I	S

S. Typhimurium were screened for teen virulence genes (*msgA*, *tolC*, *spaN*, *invA*, *ipfC*, *sitC*, *sopB*, *orgA*, *pagC* and *pefA*) by using simplex PCR, most thirty three isolates were positive for teen of the virulent genes tested. For *sitC*, three isolated were negative to this virulence gene (Fig.1). While two isolate were negative to *lpfC* (Fig. 2).




Figure 1. Agarose gel electrophoresis of *sitC* PCR products amplified from *S. Typhimurium*. M: MW marker (1 kb ladder); 1 to 33 Lanes: *S. Typhimurium* isolates.




Figure 2. Agarose gel electrophoresis of *lpfC* PCR products amplified from *S. Typhimurium*. M: MW marker (1 kb ladder); 1 to 33 Lanes: *S. Typhimurium* isolates.

One isolate # 33 was negative to *orgA* and *spaN* (Fig.3 and Fig. 4). These results suggest that *S. Typhimurium* from clinical is virulent, and that capable of causing salmonellosis in humans and it may contribute to pathogenesis.




Figure 3. Agarose gel electrophoresis of *orgA* PCR products amplified from *S. Typhimurium*. M: MW marker (1 kb ladder); 1 to 33 Lanes: *S. Typhimurium* isolates.




Figure 4. Agrose gel electrophoresis of *spaN* PCR products amplified from *S. Typhimurium*. M: MW marker (1 kb ladder); 1 to 33 Lanes: *S. Typhimurium* isolates.

4. Discussion

Salmonellosis is one of the most common food borne illnesses. FDA estimated that non-typhoidal *Salmonella* caused total 1,412,498 cases of illness. Every year, there have been approximately 42,000 cases of *Salmonella* infections reported. Though this number is large itself, this is only the reported amount. There are many milder cases of Salmonellosis that are not reported, because they are not that severe. *Salmonella* infections are usually caused by the consumption of fecal contaminated water and food [5]. Our study shows that *S. Typhimurium* isolated from clinical carried the teen virulence genes, which might play an important role in invasion and survival in the host [9]. Recently, Akiyama *et al.* (2011) and Mezal *et al.* (2013) [10,3] indicated that the same virulence genes were percent in *S. Enteritidis* and *S. Saintpaul* isolated from clinical isolates, which are capable of causing human infections. In this study *S. Typhimurium* isolates examined for antimicrobials resistance, all 33 isolates were sensitive to gentamycin, kanamycin, nalidixic acid, chloramphenicol, and sulfisoxazol. Seven isolates were resistant to ampicillin, chloramphenicol and tetracycline. Lower rates of resistance in this study are in agreement with Yang (2002) [11] that have reported a low prevalence of antimicrobial resistance among *S. Typhimurium* from sources in South Korea. Further investigations with bigger samples size are needed to identify the source and cause of drug resistance. Conclusion: *Salmonella Typhimurium* can present virulence genes (*msgA*, *tolC*, *spaN*, *invA*, *ipfC*, *sitC*, *sopB*, *orgA*, *pagC* and *pefA*) related to invasion and survival within macrophage, but that have low prevalence of antimicrobial resistance among.

References:

1. Angulo FJ, Nargund VN & Chiller TC. Evidence of an association between use of anti-microbial agents in food animals and anti-microbial resistance among bacteria isolated from humans and the human health consequences of such resistance. *J Vet Med B Infect Dis Vet Public Health* **51**: 374-379. 2004.
2. CDC .National enteric disease surveillance: *Salmonella* annual summary. 2008. <http://www.cdc.gov/ncezid/dfwed/PDFs/salmonella-annual-summary-2009-508c.pdf>. ed.^eds.), p.^pp.
3. Mezal, E.H., Stefanova, R., Khan, A.A. Isolation and molecular characterization of *Salmonella enterica* serovar Javiana from food, environmental and clinical samples. *Int. J. FoodMicrobiol.* **164**, 113-118. 2013
4. Swamy SC, Barnhart HM, Lee MD & Dreesen DW .Virulence determinants *invA* and *spvC* in *Salmonella* isolated from poultry products, wastewater, and human sources. *Appl Environ Microbiol* **62**: 3768-3771. 1996
5. CDC. Estimates of Food borne Illness. http://www.cdc.gov/food_borne_burden/2011-foodborne-estimates.html. 2011.
6. Guibourdenche, M., Roggentin, P., Mikoletit, M., Fields, P.I., Bockemuhl, J., Grimont, P.A.D., Weill, F.X., 2010. Supplement 2003-2007(No.47) to the White -Kauffmann-Le Minor scheme. *Res. Microbiol.* **161**, 26-29.
7. Khan, A.A., Cheng, C.M., Khanh, T.V., Summage-West, C., Nawaz, M.S., Khan, S.A., Characterization of class 1 integron resistance gene cassettes in *Salmonella enterica* serovars Oslo and Bareilly from imported seafood. *Journal of Antimicrobial Chemotherapy* **58**, 1308-1310. 2006.
8. Clinical and Laboratory Standards Institute, Performance Standards for Antimicrobial Susceptibility Testing: Sixteenth Informational

Supplement.M100-S16. CLSI, Wayne, PA, USA. 2006.

9. Skyberg, J.A., Logue, C.M., Nolan, L.K., Virulence genotyping of *Salmonella* spp.with multiplex PCR. *Avian Diseases* 50, 77–81. 2006.
10. Akiyama, T., Khan, A.A., Cheng, C.M., Stefanova, R., Molecular characterization of *Salmonella enterica* serovar Saintpaul isolated from imported seafood, pepper, environmental and clinical samples. *Food Microbiol.* 28, 1124-1128. 2011
11. Yang, S., Park, K.Y., Kim, S.H., Min No, K., Besser, T.E., Yoo, H.S., Kim, S.H., Lee, B.K., Park, Y.H., Antimicrobial resistance in *Salmonella enterica* serovars Enteritidis and Typhimurium isolated from animals in Korea: comparison of phenotypic and genotypic resistance characterization. *Vet. Microbiol.* 112, 1-10. 2002.