

EFFECT OF POTASSIUM FERTILIZERS APPLICATION ON GROWTH, YIELD AND WATER USE EFFICIENCY OF POTATO UNDER REGULATED IRRIGATION TREATMENTS ⁺

تأثير معدلات التسميد بالبوتاسيوم في نمو وحاصل البطاطا وكفاءة استعمال المياه تحت معاملات ري مختلفة

A.S. Ati ^{*}

S. M. Nafaou ^{**}

Abstract:

Studies on the effect of K fertilizers application in increasing water use efficiency and decreasing the negative effect of water stress, quite limited especially under field conditions. A field experiment was conducted in silty clay loam in Agricultural Research Station of Abu-Graib- Baghdad, Iraq to study the effect of different levels of K fertilizers application on potato (*Solanum tuberosum* L.) and their role in decreasing negative effect of water stress.

Treatments include four levels of K fertilizer (0, 200, 400 and 600 kg K₂SO₄ ha⁻¹) and two levels irrigation (after depletion of 50 and 75 % of available water). Results revealed the following:-

1. The negative effects of water stress on potato growth and yield were decreased by increased levels application of K fertilizers.
2. Negative effects of water stress on plant quality and water use efficiency were decreased by application of increase level of K fertilizer in depletion 75% of available water, and saving in irrigation water with irrigation scheduling by 9%.
3. Tuber yield, plant height, leaf area and carbohydrates increased with increased K application. Higher K rates did improve yield reached to 30.26 and 25.66 T.ha⁻¹ in irrigation with 50 and 75% depletion of available water, respectively.
4. As a conclusion nutrients application may have a good impact on plant performance under water stress, and decreasing amount of water applied through alterative irrigation improved yield of fruit. Therefore this experiment signifies the importance of potassium application together with conserving water application.

المستخلص :

ان الدراسات ذات العلاقة بالمستويات المختلفة للبوتاسيوم ودورها في تقليل التأثير السلبي للشد الرطوبي تحت الظروف الحقلية، قليلة لاسيما في ظروف المياه الحالية وضرورة ايجاد السبل الكفيلة للتعايش مع العجز المائي القادم.

⁺ Received on 4/9/2011 , Accepted on 12/9/2012 .

^{*} prof ./ Collage of Agric. /University of Baghdad

^{**} Assit .prof ./ Collage of Agric. /University of Baghdad

نفذت هذه الدراسة في احد حقول كلية الزراعة-جامعة بغداد في محافظة بغداد -العراق في تربة مزيج طينية غرينية، لدراسة مستويات مختلفة من البوتاسيوم في نمو وحاصل البطاطا (*Solanum tuberosum* L.) لتقليل التأثير السلبي للشد الرطوبي.

تضمنت معاملات التجربة على اضافة أربعة مستويات من البوتاسيوم وهي ٠، ٢٠٠، ٤٠٠ و ٦٠٠ كغم K_2SO_4 . هـ^١ ومستويين من الري بعد استنفاد ٥٠ و ٧٥% من الماء الجاهز. وبينت النتائج الآتي:-

١. اختزال التأثيرات السلبية للشد الرطوبي (عجز المياه) في كمية حاصل النبات ومعدل نموه بزيادة مستويات اضافة البوتاسيوم.

٢. اختزال التأثيرات السلبية للشد الرطوبي في الصفات النوعية للنبات وكفاءة استخدام المياه بزيادة مستويات اضافة البوتاسيوم وعند استنفاد ٧٥% من الماء الجاهز، ليصل معدل تقليل كمية المياه المضافة الى ٩%.

٣. ازدياد حاصل البطاطا، ارتفاع النبات والمساحة الورقية ومحتوى الدرنات من الكربوهيدرات بزيادة مستويات اضافة البوتاسيوم. اذ سجل اعلى معدل لانتاج البطاطا ٣٠.٢٦ و ٢٥.٦٦ طن هكتار^١ عند الري باستنفاد ٥٠ و ٧٥% من الماء الجاهز.

٤. بشكل عام ان اضافة المغذيات (الاسمدة) ممكن ان يحسن من أداء النبات تحت اجهاد رطوبي معتدل ممكن ان يستخدم لزيادة كفاءة استخدام المياه وكمية المياه المضافة.

Introduction:

The potato (*Solanum tuberosum* L.) is the fourth most important world crop, after wheat, maize and rice [1]. It is a major source of inexpensive energy. It contains high levels of carbohydrates and significant amounts of vitamins B and C and other minerals. In Iraq, potato is one of the most important crops, average yields 15.84 t/ha [2]. The crop is judiciously fertilized with N, P and K and other elements based mainly on practical experience as there is a lack of recommendations based on correlation/calibration research. Potato plants require much more potassium than many other vegetable crops.

Although most soils in Iraq are rich in K, potassium fertilizer should be applied to sustain high yields [3]. An adequate supply of potassium strengthens stems to prevent lodging, increases yield and improves tuber quality. Potatoes require high levels of potassium in concentrations which are comparable to or greater than nitrogen [4, 5]. In a similar study [5, 6]. reported that dry matter of potatoes decreased with increasing K level.

In arid and semiarid regions, potato is sensitive to water stress and irrigation has become an essential component of potato production in comparison with the other crops. Potato may be quite sensitive to drought [7] as it needs frequent irrigation for suitable growth and optimum yield [8, 9]. [10] and [11] found that all growing stages of potato, especially tuber formation stage, are very sensitive to water deficit stress. Water supply and scheduling have important impacts on tuber quality – frequent irrigation reduces the occurrence of tuber malformation. Water deficit in the early phase of yield formation increases the occurrence of spindle tubers. Using good agricultural practices, including irrigation when necessary, a crop of about 120 days in temperate and subtropical climates can yield 25 to 40 tones of fresh tubers per hectare [12].

The potato is extensively cultivated in large area of Iraq; but the effect of water stress on yield of this cultivar has not been investigated. Therefore, the objective of this study was to evaluate the effect of different rates of potassium sulfate (K_2SO_4) on plant vegetative performance, total yield and water use efficiency of potatoes grown under water stress.

Materials and Methods:

The experiment was carried out during spring seasons of 2009 at the Agricultural Research Station of Abu-Graib- Baghdad, Iraq (33° 20' N, 44° 12' E; elev. 34.1 m). Potato (*Solanum tuberosum* L.) was planted on soil of silt clay loam texture (Sand=150 g kg⁻¹, Silt=570 g kg⁻¹ and Clay=284g kg⁻¹) with average bulk density of 1.41 Mg.m⁻³ and soil content moisture 0.31 cm³ cm⁻³ at field capacity and wilting point equal 0.16 cm³ cm⁻³. The content of nutrition in soil (available of K, P and K are 267, 285 and 10.21 mg. kg⁻¹, respectively). During the cultivation seasons; the mean relative humidity was 45% and mean rainfall were 18.17 mm during potato growing season. The total soil water, calculated between field capacity and wilting point for an assumed potato root extracting depth from 0.15 to 0.45 m.

Planting took place on 7 February using seeding rate of 2000 kg ha⁻¹ in 75 cm spaced rows with net plot size of 12 m × 4.5 m, in a randomized complete block design with three replicates. Each experimental unit consisted of 6 rows. All plots were irrigated with river water an EC_i = 1.1 dS.m⁻¹. Irrigation were scheduled when soil water content in the root zone was depleted by the crop to specific fraction of available water (irrigation was imposed at 50% and 75% depletion of available water). The soil depth of the effective root zone is increased from 0.15 m at planting to 0.45 m in bulking and tuber enlargement stages. Measured amount of water were delivered to the furrows using water meter. Soil water content was measured gravimetrically. The sum of differences in soil water and applied irrigation water plus rainfall were calculated as ET_a using water balance equation, assuming negligible deep percolation, groundwater contribution and runoff.

All plots received basic application of 300 kg N and 250 kg P₂O₅ ha⁻¹. Four levels 0, 200, 400, and 600 kg K₂SO₄ ha⁻¹ were therefore studied. Granular fertilizers were hand spread before planting. Phosphorus source was di-ammonium phosphate, applied at once before planting. Nitrogen source was urea 46-0-0, split tow times during growing season.

During the growing season, plant height and leaf area were determined. Harvesting was carried out on 26 May, 2009. Potato tubers were then graded visually into marketable (>3.5 cm in diameter) and cull (<3.5 cm, bruised, green or sprouted tubers). Marketable and total tuber yield were determined by weight, and only marketable tuber data is reported in this manuscript. Potato tuber samples were then collected from all treatments for specific gravity, percent carbohydrates and K content determinations.

Reference evapotranspiration ET₀ was calculated using Penman-Montieth modified equation [13]. Water – Use Efficiency (WUE) calculated as fallows:

$$W.U.E(kg.m^{-3}) = \frac{Yield(kg \quad ha^{-1})}{total \quad water \quad applied \quad m^{-3}}$$

Statistical analysis: analysis of variance was performed to evaluate the statistical effect of irrigation treatment and K fertilizer application on potato yield and yield components using [14]. LSD test was used to find any significant difference between treatment means.

Results and discussion:

The results of Table 1 showed that the addition of potassium fertilizer (K₂SO₄) to soil has achieved an increase vegetative growth of plant potato, lengths and plant leaf area index were increased, and the increase was significant with the levels of irrigation in the study (after depletion of 50 and 75% of available water). Plant height and leaf area reached to 85.76 cm, 41.22 cm². Plant⁻¹ and 74.23 cm, 34.78 cm².plant⁻¹ in irrigation with 50% and 75% depletion of available water, respectively in 600 kg K₂SO₄ ha⁻¹ level. The reason for this

increase with increase level of fertilizer added to soil, to the root composition fundamentally strong, active and efficient in the absorption of nutrients from the soil [4] and increasing concentration in the shoots, leading to improved vegetative growth and increase the efficiency of photosynthesis.

The results were similar to the attributes of the content of carbohydrates and potassium in the tuber with increase the level of fertilizer potassium sulfate with two levels irrigation. Tuber potassium content increased significantly with K rates to reach a statistical maximum of 3.34% and 2.87% in irrigation with 50% and 75% depletion of available water, respectively at K rate of 600 kg K₂SO₄ha⁻¹ (Table 1). Carbohydrate percent was increased significantly with increasing K rates. The highest level was reached with application of 600 kg K₂SO₄ha⁻¹ to 12.69 and 11.74 % in irrigation with 50% and 75% depletion of available water, respectively. The increase potassium fertilizer caused a significance increasing the efficiency of the plant in the absorption of nitrogen, which plays a role in protein synthesis, transforms to the amino acids that move later places of manufacturing in the leaves to the storage places in the tubers to form protein, as well as the role of potassium in the process of protein synthesis [4].

Table 1. Effect of different levels of potassium sulfate and irrigation level on plant height, leaf area, tuber specific gravity, carbohydrates, potassium concentration and marketable yield.

Irrigation level (depletion of AW)	Treatment K ₂ SO ₄ kg. ha ⁻¹	Plant height (cm)	Leaf area (cm ² plan ⁻¹)	Specific gravity	Carbohydr ates (%)	K (%)	Marketable tuber yield (Ton ha ⁻¹)
50%	0	65.63	28.18	1.051	11.59	2.01	19.45
	200	74.21	35.42	1.062	11.92	2.65	23.89
	400	80.22	39.34	1.069	12.45	2.97	28.34
	600	85.76	41.22	1.071	12.69	3.34	30.26
75%	0	50.68	24.06	1.050	10.76	1.89	16.22
	200	66.65	26.21	1.059	10.99	1.97	20.29
	400	71.59	31.23	1.061	11.28	2.43	23.12
	600	74.23	34.78	1.065	11.74	2.87	25.66
LSD (0.05)		4.85	2.04	N.S	0.23	0.18	1.75

The availability of potassium in the phase composition of the tubers is important and essential for the transfer of materials manufactured in the leaves, especially carbohydrates to the tubers that are stored in the form originated. It should be noted that the plants potato are characterized by (Luxury consumption) in absorption of potassium to the point of recreational consumption [15]. Our results did agree with those of [16] and [17] who found that specific gravity did not respond to K application.

Marketable tuber yield increased significantly ($p < 0.05$) with increasing K rates up to 600 kg K₂SO₄ ha⁻¹. Higher K rates did improve yield reached to 30.26 and 25.66 T.ha⁻¹ in irrigation with 50 and 75% depletion of available water, respectively.

Negative effect of water stress on potato growth, quality and yield by increased application rates of K showed the role of K in the composition of cells Alsklarnkami most thicker and hardness and are resistant to conditions of water scarcity (drought) its role in the process of closing and opening gaps Securities [18], as well as the effect of K in proline acid concentration has special importance in regulation water uptake by plants.

Table (2) shows the amounts of applied irrigation water under water stress during the growth of potato. Total water supply were 410, 404, 400, 400; 375, 370, 365, 365 mm for 0.0, 200, 400 and 600 kg K₂SO₄ ha⁻¹ treatments in 50% and 75% depletion, respectively. The

amounts of irrigation water in different treatment were similar to those reported by [19] and [20] where experiment carried out near the area of this study.

Table 2. Actual evapotranspiration (ETa) (mm), water use efficiency (WUE) (kg m⁻³) from water stress in cropping.

Irrigation level (depletion of AW)	Treatment kg K ₂ SO ₄ ha ⁻¹	Irrigation (I) mm	Rain fall (R) mm	(ETa) Actual evapotranspiration mm	WUE kg m ⁻³	Irrigation water saving (%)
50%	0	410	18.71	429	5.0	-
	200	404	18.71	423	6.0	1.0
	400	400	18.71	419	7.0	2.0
	600	400	18.71	419	8.0	2.0
mean		404	18.71	423		1.67
75%	0	375	18.71	394	4.0	-
	200	370	18.71	389	5.0	1.0
	400	365	18.71	384	6.0	3.0
	600	365	18.71	384	7.0	3.0
mean		369	18.71	388		2.33

The cumulative ETa under different water treatments are also presented in Table 2. The mean ET_a measured during the season was 404 mm for 50% depletion of AW and 369 mm 75% depletion of AW. Water use efficiency (WUE) and rainfall expressed as the ratio of potato yield to water supply from planting to harvest varied typically comparable to those obtained in other field studies [21] who reported that range of WUE was from 5 to 14 kg m⁻³.

The net saving in irrigation water with irrigation scheduling under K fertilizers application reached 8% (Table 2). Cumulative ETo calculated from Penman-Monteith modified equation totaled 472 mm which are close to the ETa (404, 369 mm) proved the validity of this equation for estimating the water requirements of potato within the context of the region.

Conclusion and Recommendation:

Both plant nutritional status and tuber quality parameters responded positively to increasing K application rates and water stress. Marketable tuber yield was significantly affected by K fertilizer levels. Highest yield and other quality parameters were achieved when a rate of 600 kg K₂SO₄ ha⁻¹ was used. We recommend that, for similar regions in Iraq, farmers apply potassium sulfate at the rate 600 kg K₂SO₄ ha⁻¹ for optimum yield. More researchers are needed to investigate the benefits of further splitting of K fertilizer for potatoes grown in the different soils texture under arid conditions of Iraq.

References:

1. Spooner, D. M. and J. B. Bamberg.. Potato genetic resources: sources of resistance and systematic. Amer. Potato J.71:325-338.1994.
2. Arab organization for agricultural development.2006. (Cited by PhD Thesis, Omar Hasham. 2009. University of Baghdad. Collage of Agriculture).

3. Mahmood, J. T. Effect of NPK addition and foliar application on growth, yield and components of potato plants tubers. MSC Thesis. University of Baghdad. Collage of Agriculture.2006.
4. Tisdale, S. L., W. N. Nelson and J. D. Havlin. Soil Fertility and Fertilizers. Prentice-Hall New Delhi.1997.
5. Westermann, D. T., D. W. James, T.A. Tindall and R. L. Hurst. Nitrogen and potassium fertilization of potatoes: Yield and specific gravity. Amer. Potato J.71:417-432.1994.
6. Hart, T. G. and O. Smith Effect of levels sources of potassium on absorption of phosphorus by potatoes. Lants. Amer. Potato J. 43:217-235.1966.
7. Van Loon, C.D. The effect of water stress on potato growth, development and yield. American Potato J., 58: 51–69.1981.
8. Yuan, B.Z.; S. Nishiyama and Y. Kang. Effects of different irrigation regimes on the growth and yield of drip-irrigated potato. Agric. Water Manag., 63: 153–167.2003.
9. Kiziloglu, F.M.; U. Sahin, T. Tune and S. Diler. The effect of deficit irrigation on potato evapotranspiration and tuber yield under cool season and semiarid climatic conditions. J. Agron., 5: 284–288.2006.
10. Thornton, M.K. Effects of heat and water stress on the physiology of potatoes. Idaho Potato Conference, Idaho.2002.
11. Shock, C.C. Efficient Irrigation Scheduling. Malheur Experiment Station, Oregon State University, Oregon, USA.2004.
12. International Year of the Potato: **Potato and Water Resources (2008)**. Secretariat Food and Agriculture Organization of the United Nations. E-mail: potato2008@fao.org.
13. Allen, R.G.; L.S. Pereira; D. Raes and M. Smith. Crop Evapotranspiration. FAO Irrigation and Drainage paper 56, Rome.1998.
14. SAS. SAS Users guide, Statistics SAS, Inst.Gary,N.C.,U.S.A. 2002.
15. Chapman, K. S. R., L. A. Sparrow, P.R. Hardman, D. N. Wright and J.R. A. Thorp. Potassium nutrition of Kennebec and Russet Burbank potatoes in Tasmania: Effect of soil and fertilizer potassium on yield, petiole and tuber potassium concentrations and tuber quality. Australian J. Exp. Agric. 32:521-527. 1992.
16. Davenport, J. R., and E. M. Bentley. Does potassium fertilizer form, source, and time of application influence potato yield and quality in Columbia basin. Amer. J. Potato Res. 78:311-318. 2001.
17. Abdeldagir, A. H., M. A. Errebhi, H. M. Al Sarhan, and M. Ibrahim. The effect of different levels of additional potassium on yield and industrial qualities of potato (*Solanum tuberosum* L.) in an irrigated arid region. Amer. J. Potato Res. 80:219-222.2003.
18. Pissarek, H. P. The development of potassium deficiency symptoms in spring rape. 2-Pfanzmerahr. Bodenk. 136:1-96.1993.
19. Onder, S.; M. Caliskan; D. Onder and S. Caliskan (2005). Different irrigation methods and water stress effects on potato yield and yield components. Agric. Water Manage. 73:73-86. 2005.
20. Erdem, T., Y. Erdem, H. Orta and H. Okursoy. Water – yield relationships of potato under different irrigation methods and regimes. Sci. Agric. 63(3):226-231.2006.
21. Bowen, W.T. Water Productivity and Potato Cultivation. pp: 229-238. In Water Productivity in Agriculture: Limits and Opportunities for improvement. Ed. J. W. Kinje, R. Baker and D. Molden. CAB International. 2003.