



Review Article

Organic Synthesis under Solvent-free Condition (Green Chemistry): A Mini Literature Review

Sahar Balkat Aljuboori¹ , Nedaa Abdulhameed Abdulrahim^{2*}, Shuhad Yassen¹ , Heba Hashim Khaleel²¹ Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Al-Rafidain University College, Baghdad, Iraq;² Department of Pharmaceutical Chemistry, College of Pharmacy, University of Baghdad, Baghdad, Iraq

Received: 19 November 2022; Revised: 13 December 2022; Accepted: 15 December 2022

Abstract

Solvents are important components in the pharmaceutical and chemical industries, and they are increasingly being used in catalytic reactions. Solvents have a significant influence on the kinetics and thermodynamics of reactions, and they can significantly change product selectivity. Solvents can influence product selectivity, conversion rates, and reaction rates. However, solvents have received a lot of attention in the field of green chemistry. This is due to the large amount of solvent that is frequently used in a process or formulation, particularly during the purification steps. However, neither the solvent nor the active ingredient in a formulation is directly responsible for the reaction product's composition. Because these characteristics have little bearing on how well or quickly the system in which the solvent is applied works, it appears unnecessary to use toxic, combustible, or environmentally hazardous solvents. However, the beneficial properties of the solvent required for the application are strongly linked to these unfortunate side effects of solvent use. Distillation can be used to recover and purify solvents because they are volatile; however, this process can produce unwanted air pollutants and be hazardous to workers during exposure.

Keywords: Organic synthesis, Green chemistry, Purification, Solvent-free reaction.

التخليق العضوي في أوساط خالية من المذيبات (الكيمياء الخضراء): مراجعة مصغرة للنشریات

الخلاصة

المذيبات هي مكونات مهمة في الصناعات الدوائية والكيميائية، ويتم استخدامها بشكل متزايد في التفاعلات الحفازة. المذيبات لها تأثير كبير على الحركية والديناميكا الحرارية للتفاعلات، ويمكن أن تغير بشكل كبير انتقائية المنتج. ويمكن أن تؤثر على انتقائية المنتج ومعدلات التحويل ومعدلات التفاعل. ومع ذلك، فقد حظيت المذيبات بالكثير من الاهتمام في مجال الكيمياء الخضراء. ويرجع ذلك إلى الكمية الكبيرة من المذيبات التي يتم استخدامها بشكل متكرر في التفاعلات الكيميائية، خاصة أثناء خطوات التنقية. ومع ذلك، لا يكون المذيب ولا المكون النشط في التركيبة مسؤولاً بشكل مباشر عن تكوين منتج التفاعل. نظراً لأن هذه الخصائص لها تأثير ضئيل على مدى جودة أو سرعة عمل النظام الذي يتم فيه تطبيق المذيب، يبدو أنه من غير الضروري استخدام مذيبات سامة أو قابلة للاحتراق أو خطيرة بيئياً. ومع ذلك، فإن الخصائص المفيدة للمذيب المطلوب للتطبيق ترتبط ارتباطاً وثيقاً بهذه الآثار الجانبية المؤسفة لاستخدام المذيبات. يمكن استخدام التقطير لاستعادة وتنقية المذيبات لأنها متطايرة. ومع ذلك، يمكن أن تنتج هذه العملية ملوثات هواء غير مرغوب فيها وتكون خطرة على العمال أثناء التعرض.

* Corresponding author: Nedaa A. Abdulrahim, Department of Pharmaceutical Chemistry, College of Pharmacy, University of Baghdad, Baghdad, Iraq; Email: nedaarahim@yahoo.com

Article citation: Abdulrahim NA, Aljuboori SB, Yaseen S, Khaleel HH. Organic synthesis under solvent-free condition (Green chemistry): A mini literature review. *Al-Rafidain J Med Sci.* 2022;3:109-115. doi: 10.54133/ajms.v3i.94.



INTRODUCTION

The solvent, which accounts for approximately 90% of the total mass of the reaction component, is essential in a solution-based chemical reaction. Solubility, equilibrium position, and rate of reactions are all affected by solvents [1]. Because all biochemical and chemical reactions occur in moist environments, solvent selection is one of the most difficult problems in modern synthetic chemistry [2]. Because it operates on the premise that "like dissolves like," the one solvent method is the best choice for chemical synthesis. However, most, if not all, reactions can occur in mixed-solvent systems. The mixed solvent system has been widely used for a long time. Despite the fact that mixed solvents have been used in synthesis chemistry, there has been little research into how we can choose mixed solvents. Because water and ethanol are both effective solvents, the ethanol-water system has been widely used. According to numerous studies in this area over the last decade and recently published monographs, the effects of solvents on chemical processes and the physical and chemical properties of compounds continue to pique the interest of researchers. The effect of the solvent on the protonation-deprotonation equilibrium or the change in basicity/acidity constants when switching solvents is one of the most important aspects of this issue [3,5]. Professors frequently discuss and rationalize the types of solvents used for various transformations when introducing new chemistry students to the concepts of organic reactivity. For operations involving Grignard reactions and the SN2 mechanism, examples include the commonly used solvents acetonitrile and dry ether. Students quickly wonder, "Which solvent should I select?" when developing conditions for their own chemistry experiments, having quickly realized that solvents are an important part of organic processes [6,7].

The Role of Solvents in Chemical Processes

Organic reactions are thought to take place in a solvent solution medium in general. The rationale for this concept is straightforward. That is, if the reactant compounds are in a homogeneous solution that can be shaken, stirred, or agitated in any way, the molecules of the reactant can interact efficiently and quickly. Furthermore, if necessary, it is extremely simple to uniformly heat or cool the combination in a solution. However, a solvent's role in an organic reaction entails much more than simply providing a homogeneous environment for numerous interactions between the reactants. A solvent can sometimes significantly increase or decrease the rate of a reaction [8]. A change in the solvent of a reaction can affect how quickly it proceeds and even change its direction. This could manifest as varying product yields and ratios. As a result, a solvent can be intimately and irreversibly linked to the processes of many organic reactions by preserving reactants, products, transition states, or other intervening species. Because of a variety of factors, including

electrostatic, static, and conformational effects, the solvent and reaction partners have such direct contacts. Any liquid can be used as a solvent, but only a few are commonly used solvents—hydrocarbons, chlorinated hydrocarbons, alcohols, esters, ethers, ammonia, H₂O, and so on. With the exception of solvolysis reactions, the reactants can be used alone or in combination with clays, zeolites, silica, alumina, or other matrices to conduct a solid reaction that produces a solvent-free product and is collected without change at the end of the reaction process. Even so, without a solvent, it may be impossible to imagine or plan a reaction [9].

Toxicity of Common Organic Solvents

Any product containing solvents, including fruits, meats, and vegetables, spoils quickly when left out at room temperature. This means that the use of solvents facilitates a variety of processes, such as chemical reactions that change the form of an object. Dry fish, meat, grains, vegetables, and so on will remain stable and in normal condition if the dryness is properly maintained. This fundamental understanding explains how chemical reactions occur in the liquid state. The general classifications for common organic solvents are aliphatic hydrocarbons, aromatic hydrocarbons, cyclic hydrocarbons, halogenated hydrocarbons, amines, ketones, esters, ethers, aldehydes, alcohols, and so on (Table 1) [10].

Table 1: The general classes of the organic solvents [10]

Hydrocarbons	Alcohols	Ethers	Chlorinated solvents
n-Pentane	Methanol	Diethyl ether	Methylene chloride
n-Hexane	Ethanol	Diisopropyl ether	Chloroform
n-Heptane	n-Propanol	Dibutyl ether	Carbon tetrachloride
n-Octane	i-Propanol	Methyl tert butyl ether	1,2-dichloroethane
n-Nonane	n-Butanol	1,4-Dioxane	1,1,1-Trichloroethane
n-Decane	i-Butanol	Tetrahydrofuran	Trichloroethylene
Benzene	2-Butanol	Esters	Perchloroethylene
Toluene	n-Amyl alcohol	Methyl acetate	Monochlorobenzene
2,2,4-Trimethyl pentane	i-Amyl alcohol	Ethyl acetate	Miscellaneous solvents
Cyclohexane	Cyclohexanol	Isopropyl acetate	Dimethylformamide
2,2,4-Trimethylpentane	n-Octanol	n-Butyl acetate	Dimethylacetamide
Cyclohexane	Ethandiol	Cellosolve acetate	Dimethylsulphoxide
Ethylbenzene	Diethylene glycol	Glycol Ethers	Sulfolane
Ketones	1,2-Propanediol	Propylene glycol methyl ether	Carbon disulphide
Acetone		Ethylene glycol methyl ether	Acetic acid
Methyl ethyl ketone		Ethylene glycol ethyl ether	Aniline
Methyl isobutyl ketone		Ethylene glycol monobutyl ether	Nitrobenzene
Cyclohexanone			Morpholine
n-Methyl-2-pyrrolidone			Pyridine
Acetophenone			2-Nitropropane
			Acetonitrile
			Furfuraldehyde
			Phenol

Organic synthesis is difficult to perform without organic solvents. Although many organic solvents are chemical compounds used to dissolve other chemical compounds, understanding the reactivity of each individual solvent under specific reaction conditions is critical. As a result, it is critical to always consider the appropriate solvents for a reaction. Solvents are typically chosen based on prior experience with specific solvents and their chemical structures, or on similar patterns discovered in the literature and demonstrated in a laboratory setting [10]. However, due to some strict guidelines and regulations that we must follow, such as solvent power and volatility,

which can be toxic to the researcher as well as our society and the environment as a whole, the situation has recently changed. This drives the search for ever-better alternatives, which is why solvent-free organic synthesis is booming right now [11,12] (Figure 2).




Figure 1: Common organic solvents' structures [11].

Numerous studies on the toxic effects of solvents have revealed the molecular mechanism as well as the likely main organs that may be affected [13-15]. Because researchers are becoming more aware of the dangers of organic solvents faster than the general public, there is a call for safety, strict rules, and regulation, all of which reduce the danger and associated health effects. There are more laboratories with well-equipped setups, good ventilation to draw reagent and solvent fumes into very diluted atmospheres, and proper disposal systems; however, the scale of synthesis and the number of researchers are rapidly growing, resulting in a higher volume of organic solvent consumption that still poses a risk. Many organizations are attempting to address this issue. Organic solvents have high lipophilicity and volatility because their skeletons are primarily composed of carbon and hydrogen atoms with the occasional hetero atom. Lipophilicity influences the distribution of solvents to various body parts from the standpoint of toxicity. To improve renal elimination, the lipophilic compounds must be converted into a water-soluble form via a series of metabolic processes. The metabolite that forms is sometimes even more toxic than the original compound [12]. Organic solvents can enter the brain and have very negative effects because of their high lipophilicity. Some chemicals have anesthetic (Trichloroethylene, as an example, [16]), anxiolytic (Toluene, as an example, [17]), convulsive (Fluorothyl, for example, [18]), anticonvulsant (Toluene, as an example, [19]), narcotic

(Trichloroethylene), and antidepressant (Benzyl Chloride) effects at high doses. Organic solvents affect a large portion of the population as a result of the rapidly expanding plastic and chemical industries. Organic solvents are easily exposed to air due to their high volatility. Because the majority of these solvents are inhaled via respiration, the lungs are the first to be affected, alerting employers to the need for adequate ventilation in the workplace. As a synthetic chemist, the researcher must spend a significant amount of time in a lab on a daily and ongoing basis handling a variety of toxic reagents and organic solvents [14].

Guidelines for using Chemicals and Solvents

Before handling any chemicals, always wear appropriate protective clothing and read the Material Safety Data Sheet (MSDS) to learn more about specific chemicals, which will provide information about potential hazards, safety precautions, other handling techniques, and so on. When working with chemicals or solvents, always use fully operational chemical fume hoods, and remove any contact lenses you may be wearing and immediately wash your eyes with running water for at least 15-20 minutes. If you suspect chemical eye contact, keep the eyelid open to allow solvents to quickly evaporate from the eye and reduce your risk of further injury. Similarly, if skin comes into contact with chemicals, wash the area thoroughly with water, apply an emollient to soothe the area, remove any contaminated clothing and shoes, and seek medical attention if necessary. If at all possible, avoid working alone in the lab [20].

Organic Reactions in the Solvent-free Conditions

Although any chemical reaction can occur in a gaseous, solid, or liquid state, many chemical changes in solutions typically occur in the liquid phase. It is the ideal medium for transferring heat between endothermic and exothermic chemical reactions. Solvents dissolve liquid or gaseous reactants, break the crystal lattice of solid reactants, and significantly affect reaction rates and chemical equilibrium positions. Furthermore, the reactant molecules can combine quickly if the reactants are in a homogeneous solution that allows for stirring, shaking, or other types of mixing [21,22]. Furthermore, it is simple to uniformly heat or cool the solution mixture as needed. However, the function of a solvent in an organic reaction entails much more than simply creating a homogeneous environment in which many reactants can collide. A solvent can sometimes significantly increase or decrease the rate of a reaction. The process of changing the solvent can affect the rate of the reaction and even change its overall course. This could manifest as different product yields and ratios. As a result of reactant solvation, a solvent can become inextricably and irreversibly linked to the operation of an organic reaction, products, a transition state, or another species that came before. Although the

solvent remains unchanged and is recovered after the reaction, electrostatic, steric, and conformational effects are the main causes of such close interactions between the solvent and the reaction partners. A reaction without a solvent has not yet been imagined or planned. However, another adage states that the best solvent is none at all [23]. Despite the strength of this assertion, the use and comprehension of solvent-free conditions has not advanced as much as solvent-based techniques. Because different names are frequently used, it can be difficult to define the concept of solvent-free synthesis. Scott *et al.* [24] make the following distinctions to make their points clear: i) solid-phase reaction, which is the reaction of molecules from the liquid phase with a solid substrate, such as polymer-supported synthetic peptide; ii) solvent-free reaction, which is any system in which reagents react with one another without the use of a solvent; and iii) solid-state reaction or solid-solid reaction, which is when two macroscopic solids interact directly without the help of a liquid or vaporous phase to create a third, solid product. Despite the presence of macroscopic organic solids in many solvent-free reactions, the majority of them occur in a liquid or melt phase. This fascinating fact has served as the basis for extensive research on a variety of organic reactions [25]. These organic reactions reported as "in the solid phase" involve the formation of a liquid phase. This melting implies the presence of a eutectic mixture with a melting point lower than the melting points of the reactants. When heating is required, it is sometimes obvious that a phase change (from solid to liquid) occurs. Solvent-free reactions may provide access to many simple and complex reactions, as well as all classes of compounds. Among the most common and significant reactions are carbonyl compound condensation, cycloadditions, alkylation, aromatic substitution, amine, water, and alcohol additions, cyclization, eliminations, rearrangements, carbon-carbon coupling, cascade reactions, and catalyzed reactions [26]. Compared to reactions in molecular solvents, solvent-free reactions have the following advantages: i) No reaction medium exists for collection, purification, or recycling; ii) The chemicals produced are frequently sufficiently pure to avoid thorough chromatographic purification, and in certain situations, recrystallization is not even necessary; iii) In high-yielding systems, successive solvent-free reactions are feasible; iv) when compared with organic solvents, the reactions are quick, sometimes reaching a significant completion in a few minutes; v) specialized equipment is frequently not required; vi) energy consumption may be considerably lower; vii) preformed salts and metal-metalloid complexes are frequently unnecessary; viii) it is possible to prevent functional group protection-deprotection; ix) while establishing industrial processes, there can be a smaller capital expenditure for equipment; and x) solvent-free methods may be made more economically viable and ecologically friendly by significant reduction of batch sizes and processing costs [27]. There are several disadvantages to

solvent-free reactions, but they can be overcome with advances in reactor design technology [28]. There are some disadvantages to using solvent-free reactions, such as the formation of "hot spots" and the possibility of runaway reactions. Rather than continuing to operate under the outdated paradigm in which a reaction medium or solvent is used as a heat sink or agent for heat transfer, it would be preferable to work on designing reactors for continuous flow systems. In solvent-free processes, it is obvious that quantifying reaction heat is just as important as effective heat dissipation. If extremely exothermic processes can be demonstrated to work well in solvent-free environments, the problem may be solved by developing sophisticated reactors. Another argument against it is that working with solid or extremely viscous materials is difficult. This can be overcome through technological advancements and novel reactor designs. If a solvent-free or specific molecular solvent reaction is used, selectivity, waste, stereochemistry, viscosity, yield, ease of recycling, energy use, easy product separation, competing reactions, and reaction heat will all be affected [29]. It can be seen that 58% of the operations used MW irradiation, 29% used conventional thermal heating, 3% used grinding, and 2% used sonication.

Green Technique and its Advantages

Green chemistry is the environmentally friendly synthesis of chemicals. The synthetic schemes are designed to cause the least amount of environmental damage possible (Figure 2). The majority of environmental pollution is currently caused by a variety of chemical industries. The cost of waste disposal is also extremely high. The 12 guiding principles of green chemistry can be applied to almost any aspect of chemistry, such as the design of molecules with desirable properties and structures, process catalysis, the use of less polluting reaction conditions, and so on [30].




Figure 2: Key merits of the green synthesis methods [30].

These guidelines are summarized below:

1. Before they can be carried out, processes must be designed. Chemical products should be made with the least amount of toxicity possible while still performing the required function.

2. Avoiding waste altogether is preferable to cleaning up or treating garbage that has already been produced as a preventive measure.

3. The use of less hazardous solvents and auxiliaries. Auxiliary chemicals and processes (separation, solvents, reagents, etc.) should be rendered harmless and used as little as possible [31].

4. Design for energy efficiency. The energy required to perform chemical processes should be understood in many terms, including their effects on the environment and the economy, and it should be kept to a minimum. Synthetic procedures are to be carried out at room temperature and pressure whenever we can.

5. Less hazardous chemical synthesis. The synthetic pathways should always be designed to utilize and produce chemicals wherever possible that are healthy for both the environment and humans.

6. Atomic economy. The inclusion of all components utilized throughout the process into the finished product should be kept to a minimum when developing synthetic procedures. In this instance, Ibuprofen's synthesis by the BHC firm, which generates less waste and byproducts, is a very good example. In comparison to the 6-step brown synthesis (BOOTS approach), the 3-step catalytic green synthesis (BHC approach) is more economical by 77% and 40%, respectively. The percentage of atom economy increases to 99% from 77% when taking into account the recovered acetic acid produced in step 1 of green synthesis.

7. Utilizing catalysts Catalytic reagents, which should be as selective as possible, are superior to stoichiometric reagents. By utilizing the catalytic reagents, one can increase selectivity, raise yield, and produce reactions that weren't previously achievable.

8. Minimize derivative use. Wherever possible, derivatization that is not necessary (such as the use of blocking groups, protection or deprotection, and temporary change of chemical and physical processes) should be avoided as it might lead to waste and demand more reagents.

9. Planning for deterioration. When a chemical product has served its purpose, it should be designed to decay into harmless degradation products and not pollute the environment [32].

10. Make use of sustainable feedstock. When it is technically and financially viable, renewable raw resources, or "feedstock," are preferable to finite ones.

11. Immediate analysis to prevent contamination The analytical techniques must be strengthened much more. Prior to the production of dangerous compounds, real-time process monitoring and control is made possible.

12. Chemistry that is inherently safer for preventing accidents. To lessen the risk of chemical accidents like explosions, leaks, and fires, compounds and the forms of substances utilized in chemical processes should be carefully chosen [33].

Green chemistry has various benefits, including waste avoidance, non-toxicity, safety, economic viability, and environmental friendliness [34]. Synthesis manufacturing processes have been created to generate little waste, have no adverse effects on the environment, and be simple to discard [35]. To carry out reactions, starting materials, solvents, and catalysts must be carefully chosen. For instance, benzene must never be used as a solvent because it is carcinogenic. If at all possible, it is preferable to carry out reactions in the aqueous phase. With this viewpoint in mind, synthetic methods should be developed such that the starting materials are utilized as fully as is practical in the final product. The reaction must also not result in any negative byproducts [36]. It is no longer true, as once was believed, that no reaction can take place without the presence of a solvent. Such reactions are essential for industry because they are easy to operate [37], emit few pollutants, are relatively inexpensive to run, and limit pollution. The many features of solid-state organic synthesis have been covered in literature. It is believed that solvent-free organic synthesis and transformations have industrial value and are primarily environmentally favorable.

Conclusion

We have examined both historical and present general solvent selection guidelines in order to comprehend the advantages and disadvantages of each. More research is needed in the areas of application-specific techniques and life cycle studies in order to choose solvents in the future with more sophistication based on a sustainable supply chain. To do this, more knowledge will be required, especially about the physical properties and environmental effects of innovative solvents. Even though the synthesis of organic molecules without the use of solvents has grown very widespread, it remains one of the most effective techniques in green chemistry for quickly and sustainably producing a wide range of important substances. Due to the numerous advantages of solvent-free reactions, new solvent-free procedures are being developed for the ecologically friendly synthesis of diverse compounds.

Conflict of interests

The author declares no conflict of interests.

Source of fund

No specific fund received.

Data sharing statement

N/A

REFERENCES

- Jiang X, Wang Y, Li M. Selecting water-alcohol mixed solvent for synthesis of polydopamine nano-spheres using solubility parameter. *Sci Rep*. 2014;4:6070. doi: 10.1038/srep06070.
- Buncel E, Stairs RA, Wilson H, (Eds.), The Role of the Solvent in Chemical Reactions, Oxford University Press, New York, 2003.
- Reichardt C, (Ed.), Solvents and Solvent Effects in Organic Chemistry, Weinheim: Wiley, 2003, (3rd ed.) p. 472.
- Gould NS, Li S, Cho HJ, Landfield H, Caratzoulas S, Vlachos D, et al. Understanding solvent effects on adsorption and protonation in porous catalysts. *Nat Commun*. 2020;11:1060. doi: 10.1038/s41467-020-14860-6.
- Adams DJ, Dyson PJ, Tavener SJ, (Eds.), Chemistry in Alternative Reaction Media, Chichester: Wiley, 2004, p. 24.
- McMurry J, Organic Chemistry, (7th ed.), Thomson Higher Education, Belmont, CA, 2008; pp 613-615.
- Chanda A, Fokin VV. Organic synthesis "on water". *Chem Rev*. 2009;109(2):725-748. doi: 10.1021/cr800448q.
- Nagendrappa G. Organic synthesis under solvent-free condition: An environmentally benign procedure — I. *Reson* 2002;7(10):59-68. doi: 10.1007/BF02835544.
- Zhou KG, Mao NN, Wang HX, Peng Y, Zhang HL. A mixed-solvent strategy for efficient exfoliation of inorganic graphene analogues. *Angew Chem Int Ed Engl*. 2011;50(46):10839-10842. doi: 10.1002/anie.201105364.
- Smallwood I, Handbook of Organic Solvent Properties, Butterworth-Heinemann; 2012.
- Tanaka K, Toda F. Solvent-free organic synthesis. *Chemical Reviews*. 2000;100(3): 1025-1074.
- Varma RS. Solvent-free accelerated organic syntheses using microwaves. *Pure Appl Chem*. 2001;73(1):193-198. doi: 10.1351/pac200173010193.
- Spencer PS, Schaumburg HH. Organic Solvent Neurotoxicity: Facts and Research Needs, Scandinavian Journal of Work, Environment & Health. 1985, p. 53-60.
- Takeuchi Y, Ono Y, Hisanaga N, Kitoh J, Sugiura Y. A comparative study of the toxicity of n-pentane, n-hexane, and n-heptane to the peripheral nerve of the rat. *Clin Toxicol*. 1981;18(12):1395-1402. doi: 10.3109/15563658108990348.
- Baker EL. Organic solvent neurotoxicity. *Annu Rev Public Health*. 1988;9:223-232. doi: 10.1146/annurev.pu.09.050188.001255.
- Dumas O, Despreaux T, Perros F, Lau E, Andujar P, Humbert M, et al. Respiratory effects of trichloroethylene. *Respir Med*. 2018;134:47-53. doi: 10.1016/j.rmed.2017.11.021.
- Bowen SE, Hannigan JH, Davidson CJ, Callan SP. Abstinence following toluene exposure increases anxiety-like behavior in mice. *Neurotoxicol Teratol*. 2018;65:42-50. doi: 10.1016/j.ntt.2017.12.010.
- Fink M. The seizure, not electricity, is essential in convulsive therapy: the flurothyl experience. *J ECT*. 2014;30(2):91-93. doi: 10.1097/YCT.0000000000000110.
- Cruz SL, Gauthereau-Torres MY, Rivera-García MT. Structure-activity relationship for the anticonvulsant effects of organic solvents. *Neurotoxicology*. 2016;57:121-127. doi: 10.1016/j.neuro.2016.09.012.
- Joshi D, Adhikari N. An overview on common organic solvents and their toxicity. *J Pharm Res Int*. 2019;28(3):1-18. doi: 10.9734/jpri/2019/v28i330203.
- Haase S, Tolvanen P, Russo V. Process intensification in chemical reaction engineering. *Processes*. 2022;10(1):99. doi: 10.3390/pr10010099.
- Cave GW, Raston CL, Scott JL. Recent advances in solventless organic reactions: towards benign synthesis with remarkable versatility. *Chem Commun*. 2001;21:2159-2169.
- Walsh P, Li H, Parrodi CA. A green chemistry approach to asymmetric catalysis: Solvent-free and highly concentrated reactions. *Chem Rev*. 2007;107(6):2503-2545. doi: 10.1021/cr0509556.
- Dunk B, Jachuck R. A novel reactor for UV irradiated reactions. *Green Chem*. 2000;2:G13-G14. doi: 10.1039/B000158I.
- Levine RD, Bernstein RB, (Eds.), Molecular Reaction Dynamics and Chemical Reactivity, Oxford University Press: New York, 1987.
- Mukhopadhyay S, Rothenberg G, Wiener H, Sasson Y. Solid-solid palladium-catalyzed water reduction with zinc: mechanisms of hydrogen generation and direct hydrogen transfer reactions. *New J Chem*. 2000;24:305-308. doi: 10.1039/B001655L.
- Chemat F, Abert Vian M, Ravi HK, Khadhraoui B, Hilali S, Perino S, et al. Review of alternative solvents for green extraction of food and natural products: Panorama, principles, applications and prospects. *Molecules*. 2019 Aug;24(16):3007. doi: 10.3390/molecules24163007.
- Beyer MK, Clausen-Schaumann H. Mechanochemistry: The mechanical activation of covalent bonds. *Chem Rev*. 2005;105(8):2921-2948. doi: 10.1021/cr030697h.
- Sineriz F, Thomassigny C, Lou JD. Recent advances in solvent-free oxidation of alcohols. *Curr Org Synth*. 2004;1(2):137-154. doi: 10.2174/1570179043485376.
- Sharma SK, Chaudhary A, Singh RV. Gray chemistry verses green chemistry: challenges and opportunities. *Rasayan J Chem*, 2008;1:68-92.
- Girijesh KV, Keshav R, Rajiv KV, Pratibha D, Singh MS. An efficient one-pot solvent-free synthesis and photophysical properties of 9-aryl/alkyl-octahydroxanthene-1,8-diones. *Tetrahedron*. 2011;67:3698-3706. doi: 10.1016/j.tet.2011.03.078.

32. Gomez JL, Bastida J, Maximo MF, Montiel MC, Murcia MD, Ortega S. Solvent-free polyglycerol polyricinoleate synthesis mediated by lipase from *Rhizopus arrhizus*. *Biochem Engineer J*. 2011;54:111-116. doi: 10.1016/j.bej.2011.02.007.
33. Ahmad S, Mozhddeh S, Ali M, Fatemeh H. Diketene as an alternative substrate for a new Biginelli-like multicomponent reaction: one-pot synthesis of 5-carboxamide substituted 3,4-dihydropyrimidine-2-(1H)-ones. *Tetrahedron*. 2010;66:4040-4042. doi: 10.1016/j.tet.2010.04.028.
34. Ryabukhin SV, Andrey SP, Bondarenko SS, Ostapchuk EN, Oleksandr OG, Oleg VS, et al. Acyl pyruvates as synthons in the Biginelli reaction. *Tetrahedron Lett*. 2010;51(32):4229-4232. doi: 10.1016/j.tetlet.2010.06.032.
35. Weili H, Shiyan C, Qiushu X, Huaping W. Solvent-free acetylation of bacterial cellulose under moderate conditions. *Carbohydrate Polymers*. 2011; 83(4):1575-1581. doi: 10.1016/j.carbpol.2010.10.016.
36. Shunxi L, Jin-Xian W, Xiaoliu W, Xiaofang M. Mild and efficient barbier allylation reaction mediated by magnesium powder under solvent-free conditions. *Tetrahedron*. 2011;67(4):849-855. doi: 10.1016/j.tet.2010.12.035.
37. Maasoumeh J, Abdolreza R, Maryam D. Easy access to quinoxalone derivatives using alumina as an effective and reusable catalyst under solvent-free condition. *Appl Catalysis A: General*. 2011;394(1-2):48-51. doi: 10.1016/j.apcata.2010.12.022.