

Genotyping of *Listeria Monocytogenes* in Iraqi Women with Spontaneous Abortion Using Pulse Field Gel Electrophoresis

Khalid W. Qassim¹ MSc, Azhar A.F. AL Attraqchi² PhD, Yaarub I. Khatab FIBMS (Path)

¹Department of Pathology and Forensic Medicine, ²Dept. of Medical Microbiology, College of Medicine, Al-Nahrain University, Baghdad, Iraq

Abstract

Background	<i>Listeria monocytogenes</i> (<i>L. monocytogenes</i>) is a Gram-positive, facultative intracellular bacterial pathogen that can cause a severe invasive disease (listeriosis) mainly in pregnant women. Based on genetic content, <i>L. monocytogenes</i> can be divided into 3 lineages I, II and III. Several molecular methods have been developed to assist in the characterization of <i>L. monocytogenes</i> , macrorestriction analysis by pulse-field gel electrophoresis (PFGE) is one of the most used methods for the genotyping of <i>L. monocytogenes</i> .
Objective	To determine the predominant genotype of <i>L. monocytogenes</i> isolated from clinical cases in a group of aborted Iraqi women.
Methods	A study was designed and included 15 clinical isolates of <i>L. monocytogenes</i> and one isolate from locally made cheese. The PFGE protocol was performed as described by Graves and Swaminathan. Briefly, bacterial suspensions adjusted to optical density of 1.3 at 610 nm were embedded in 1.2% Sea Kem Gold agarose plugs. The lysed, washed five times and a 2 mm thick piece was cut, equilibrated and digested with 200 µl of Ascl enzyme master mix at 37 °C for 2 h. The macrorestriction fragments were separated by electrophoresis on a CHEF-DRIIBIO-RAD. Images were analyzed with the software Bio Numerics Gel Compar II version 6.6.11 (Applied Maths).
Results	All <i>L. monocytogenes</i> isolates displayed Ascl digestion profiles after incubation with the enzyme, about 6-12 major fragments of 30 to 675 kb were obtained, sixteen isolates have been used in this study, however, 11 different Ascl profiles were encountered, the distribution of the isolates according to the restriction profiles showed that two profiles were the most predominant A, A2 with a percentage of similarity of 100%.
Conclusion	<i>L. monocytogenes</i> genotyping showed profiles that can be well used in tracing down infection source and outbreak of this bacterium. PFGE represents a great discrimination and reproducibility method for molecular sub-typing of <i>L. monocytogenes</i> and it is considered the gold standard.
Keywords	<i>Listeria monocytogenes</i> , genotyping, placenta, aborted women, pulse field gel electrophoresis
Citation	Qassim KW, AL Attraqchi AAF, Khatab YI. Genotyping of <i>Listeria Monocytogenes</i> in Iraqi women with spontaneous abortion using pulse field gel electrophoresis. Iraqi JMS. 2018; Vol. 16(1): 8-13. doi: 10.22578/IJMS.16.1.3

List of abbreviations: CHEF = Contour-clamp homogeneous electric field, PFGE = Pulse field gel electrophoresis, RFLP = Restriction Fragment Length Polymorphism, UPGMA = Unweighted-pair group method with arithmetic averages

Introduction

Listeria monocytogenes (*L. monocytogenes*) is a Gram-positive intracellular, rod shaped, nonspore-

forming, motile, facultative anaerobic, bacterium. It causes a severe invasive disease (1).

Listeriosis during pregnancy may lead to intrauterine infection, which may result in severe complications like preterm labor, spontaneous abortion, and stillbirth and or infection of the neonate which may result in high morbidity and mortality rate (2). Of several molecular methods currently available like

polymerase chain reaction (PCR), amplified fragment length polymorphism detection, DNA sequencing and others, macrorestriction analysis by pulse-field gel electrophoresis (PFGE) is one of the most used methods for the subtyping of *L. monocytogenes*. The using of restriction endonuclease *Ascl*, as advised by Pulse Net USA, has shown excellent discrimination for *L. monocytogenes* and the technique is shown to be reproducible. PFGE is considered to be the international standard for subtyping ⁽³⁾.

PFGE is a form of RFLP (Restriction Fragment Length Polymorphism) typing in which, bacterial genome is digested with a rare cutting restriction enzymes that cut the bacterial genomic DNA infrequently and therefore generate a small number of DNA fragments between (10-20 bands); these fragments are of a large sizes, from 20 kb to 10,000 kb, and are separated using specific electrophoresis techniques ⁽⁴⁾ for *L. monocytogenes*, the enzymes are *Ascl* or *Apal*, these enzymes may generate between 6 to 12 and 14 to 17 fragments respectively separated through the PFGE. Combinations of the two profiles generated by the enzymes are used to characterize *L. monocytogenes* strains ⁽⁵⁾.

Genetic comparisons among isolates carry out by using the differences in the restriction profiles. Computer-based analysis is very simple and enables rapid comparison on strains. Currently, PFGE represent the “gold standard” of the molecular typing techniques for foodborne pathogenic bacteria such as *Salmonella*, *E. coli*, *Yersinia*, *Vibrio* and *Listeria* ⁽⁶⁾.

The aim of this study and the benefit of it is to assess the extent of *L. monocytogenes* in causation of human spontaneous abortion in a sample of Iraqi women, and to determine the genetic subtypes of listeria monocytogenes associated with abortion.

Methods

A cross-sectional study was designed that included 250 placental tissues obtained from

aborted women (which their acceptance was taken before sample collection) attended Al-Imamein AL-Kadhimein Teaching Hospital in Baghdad during the period from June 2014 to November 2015. Out of total number of the placentas samples, only fifteen isolates of *L. monocytogenes* were identified. The isolation and identification of *L. monocytogenes* were performed according to Collee et al. 1996 and McFaddin, 2000 ^(7,8).

PFGE of *L. monocytogenes*

The laboratory Protocol for molecular subtyping of *L. monocytogenes* by PFGE includes the following:

The Culture growth: An isolated colony from a pure culture of *L. monocytogenes* was streaked onto nutrient agar (NA) plates and the plates were incubated at 37 °C for 14-18 h.

Pulse field gel electrophoresis: The PFGE protocol was performed as described by Graves and Swaminathan (2001) ⁽⁶⁾. Briefly, bacterial suspensions adjusted to optical density of 1.3 at 610 nm were embedded in 1.2% SeaKem Gold agarose plugs. The lysed, washed five times and a 2 mm thick piece was cut, equilibrated and digested with 200 µl of *Ascl* enzyme master mix at 37 °C for 2 h. The macrorestriction fragments were separated by electrophoresis on a CHEF-DR II BIO-RAD (USA) in a 1% Pulse Field Certified Agarose gels (Ultrapure DNA grade agarose) at Initial switch time: 4.0 s; Final switch time: 40.0 s; Voltage: 6 V; Included Angle: 120° Run time: 18-19 hours.

Staining and Documentation of an Agarose gel: After electrophoresis run was over, gel was removed and stained with ethidium bromide for 20-30 min in covered container. The use of restriction endonucleases *Ascl*, is advised by Pulse Net USA, and has shown excellent discrimination for *L. monocytogenes*. Images were analyzed using Gel Compare II evaluation software version 6.6.11 from Applied Maths.

The similarity between fingerprints was determined using number of different bands coefficient with a 1% tolerance between band positions. The cluster analysis and generation of Dendrogram was performed using

Unweighted-pair group method with arithmetic averages (UPGMA) with a similarity score value of 100% as the cutoff.

Results

All isolates of *Listeria monocytogenes* were digested with *Ascl* restriction endonuclease...GG▼CGCGCC...
...CCGCGC▲GG...

Only one was isolated from local made cheese, which presented in lane number 1 (Figure 1). Isolates of *L. monocytogenes* were typed using PFGE. Analysis of the *Ascl* restriction profiles by automated cluster analysis was done using number of different bands as a similarity coefficient and tolerance: 1 %.

All the isolates displayed *Ascl* digestion profiles after incubation with the enzyme (Figure 1), about 6-12 major fragments of 30 to 675 kb were obtained following the digestion using *Ascl*.

Sixteen isolates have been used in this study, fifteen represent the clinical isolates and the

Eleven PFGE types and two subtypes were distinguished with the restriction endonuclease *Ascl* through visual inspection, using Unweighted-pair group method with arithmetic averages (UPGMA) with a similarity score value of 100% as the cutoff.




Figure 1. Genomic DNA restriction patterns of *Listeria monocytogenes* isolates, performed by PFGE after digestion with *Ascl*

Cluster analysis of the *Ascl* restriction digest profiles showed that PFGE subtype A and subtype A2 were predominant with a percentage of similarity of 100% for both followed by type A3 with a percentage of similarity of 99% then A5 and B with a percentage of similarity of 98%, D and A of

97.5%, A4 and A5 of 97%, A1 of 94.5%, F and E of 93.5%, I was of 93%, and G was presented with a percentage of similarity of 90%. Cluster analysis of *L. monocytogenes* isolates and the PFGE types assigned by visual inspection and by automated cluster analysis are shown in (Figure 2).




Figure 2. Dendrogram representing PFGE types assigned by Automated Cluster Analysis (ACA) of the Ascl restriction digest profiles of *Listeria monocytogenes* isolates

Discussion

The large number of different PFGE types (sixteen isolates yield eleven PFGE types) may indicate that a variety of genetically distinct strains cause listeriosis in Iraq, however, this high number of PFGE types is in line with several published data for clinical cases ^(9,10), this variation in the number of PFGE types obtained in different studies may be due to that the clinical strains used in these studies not corresponded to sporadic cases; the choice of the rare cut endonucleases; and the analysis protocol used in the study. However, the current work was done using the Pulse Net method described by Graves and Swaminathan ⁽⁶⁾.

The cluster analysis results showed that there was two predominant subtypes A and A2, which include three isolates and two

respectively, those isolates were genetically indistinguishable from each other and also genetically related to other subtypes and or PFGE types with a percentage of similarity range from 99% to 89%, however those with high similarity may differ in a single band only while the lower similarity percentage was associated with isolates differ in more than one band.

Although PFGE profiles give a considerable degree of strain diversity in *L. monocytogenes* populations, the number of bands differences required for discrimination between strains is controversial ⁽¹¹⁾. Strains differing by more than one band are clearly distinct but some strains from geographically distinct sources may exhibit few or no differences in DNA profile and by the Turnover criteria could be considered epidemiologically related ⁽¹²⁾. According to

Autio et al. (2002) ⁽¹³⁾, the isolation of indistinguishable PFGE types may mislead the investigation towards establishing the vehicle or the source of infection. However, many food products may carry genetically identical strains without any cross-contamination or common infection sources. Therefore, a suspected epidemic infection may consist of two or more independent sporadic cases.

In conclusion, human listeriosis in Iraq involves a great variety of genetically different isolates. However, in this study, indistinguishable isolates were isolated from more than one patient, suggesting the existence of more than one episodes of listeriosis. In all of the listeriosis episodes, the food vehicle was not determined, because of a lack of epidemiological evidence.

Acknowledgments

The authors are deeply thankful to the staff of Laboratory of Medical Microbiology Department and the staff of Laboratory of Pathology Department in the Medical College of Al-Nahrain University for their valuable support and advice. The authors also would like to thank Dr. Saad Sabah and Mr. Farked Farhan in Ministry of Science and Technology, for their continuous support in providing a suitable environment to finish my research.

Authors Contribution:

Qassim collected the cases, performed the bacterial tests and analyzed the results. Dr. AL Attraqchi helped in the study design and supervising the work. Dr. Khatab participated collection of cases and performed the histopathological examination.

Conflict of interest

The authors declare no conflict of interest.

Funding

This research was funded by College of Medicine/Al-Nahrain University.

References

1. Nes FD, Riboldi GP, Frazzon AP, et al. Antimicrobial resistance and investigation of the molecular

epidemiology of *Listeria monocytogenes* in dairy products. *Rev Soc Bras Med Trop*. 2010; 43(4): 382-5. doi: 10.1590/S0037-86822010000400009.

2. Jahangirsakht A, Kargar, M, Mirzaee, A, et al. Assessing *Listeria monocytogenes* hly A gene in pregnant women with spontaneous abortion using PCR method in Yasuj, south west of Iran. *Afr J Microbiol. Res*. 2013; 7(33): 4257-60. doi: 10.5897/AJMR12.1484.
3. Roussel S, Félix B, Grant K, et al. Fluorescence amplified fragment length polymorphism compared to pulsed field gel electrophoresis for *Listeria monocytogenes* subtyping. *BMC Microbiol*. 2013; 13: 14. doi: 10.1186/1471-2180-13-14.
4. Herschleb J, Ananiev G, Schwartz DC. Pulsed-field gel electrophoresis. (2007). *Nat Protoc*. 2007; 2(3): 677-84. doi:10.1038/nprot.2007.94.
5. Carriere C, Allardet-Servent A, Bourg G, et al. DNA polymorphism in strains of *Listeria monocytogenes*. *J Clin Microbiol*. 1991; 29(7): 1351-5.
6. Graves LM, Swaminathan B. PulseNet standardized protocol for subtyping *Listeria monocytogenes* by macrorestriction and pulsed-field gel electrophoresis. *Int J Food Microbiol*. 2001; 65(1-2): 55-62. [https://doi.org/10.1016/S0168-1605\(00\)00501-8](https://doi.org/10.1016/S0168-1605(00)00501-8).
7. Collee JG, Fraser AG, Marmion BP, et al. Mackie and McCartney Medical microbiology. 14th ed. USA: Churchill Living Stone. Inc; 1996.
8. McFaddin JF. Biochemical tests for identification of medical bacteria. 1st ed. Baltimore USA: The Williams and Wilkins; 2000.
9. Gianfranceschi M, Pourshaban M, Gattuso A, et al. Characterization of *Listeria monocytogenes* strains isolated from food and humans in Italy by pulsed-field gel electrophoresis. *Food Microbiol*. 2002; 19(1): 47-55. doi: <https://doi.org/10.1006/fmic.2001.0463>.
10. Lozniewski A, Humbert A, Corsaro D, et al. Comparison of sludge and clinical isolates of *Listeria monocytogenes*. *Lett Appl Microbiol*. 2001; 32(5): 336-9. doi: 10.1046/j.1472-765X.2001.00918.x.
11. Brosch R, Brett M, Catimel B, et al. Genomic fingerprinting of 80 strains from the WHO multicentre international typing study of *Listeria monocytogenes* via pulsed-field electrophoresis (PFGE). *Int J Food Microbiol*. 1996; 32(3): 343-55. doi: [https://doi.org/10.1016/S0168-1605\(96\)01147-6](https://doi.org/10.1016/S0168-1605(96)01147-6).
12. Tenover FC, Arbeit RD, Goering RV, et al. Interpreting chromosomal DNA restriction pattern produced by Pulsed-field gel electrophoresis: Criteria for bacterial strain typing. *J Clin Microbiol*. 1995; 33(9): 2233-9.
13. Autio T, Lundén J, Frederiksson-Ahomaa M, et al. Similar *Listeria monocytogenes* pulse types detected in several foods originating from different sources. *Int J Food Microbiol*. 2002; 77, 83–90. doi: [http://dx.doi.org/10.1016/S0168-1605\(02\)00055-7](http://dx.doi.org/10.1016/S0168-1605(02)00055-7).

Correspondence to Dr. Azhar A.F. AL Attraqchi
E-mail: tariq_963@yahoo.com
dr.azhar.ibrahim@colmed-alnahrain.edu.iq

Received Jan.11th 2017
Accepted May.24th 2017