



IRAQI
Academic Scientific Journals



العراقية
المجلات الأكاديمية العلمية

TJAS

Tikrit Journal for
Agricultural
Sciences

ISSN:1813-1646 (Print); 2664-0597 (Online)

Tikrit Journal for Agricultural Sciences

Journal Homepage: <http://tjas.tu.edu.iq>

E-mail: tjas@tu.edu.iq

**Khaldon Aziz Hammed
Al-Azawie ***
Taher Abdullateef Shujaa

Department of Animal
Production

College of Agriculture -
University of Tikrit

KEY WORDS:

Corn gluten meal, Soybean
meal, Awassi lambs.

ARTICLE HISTORY:

Received: 12/03/2021

Accepted: 28/05/2021

Available online: 30/06/2021

Tikrit Journal for Agricultural Sciences (TJAS)

EFFECT OF USING THREE SOURCE OF NITROGEN ON PRODUCTIVE PERFORMANCE and SOME BLOOD PARAMETERS OF AWASSI LAMBS

ABSTRACT

The study was carried out at the farm located at the Animal Production Farm, College of Veterinary Medicine - University of Tikrit, For the period from 1st Desember 2019 to 9th February 2020 (70 days) excluding 14 days as an adaptation period. The aim of this study find source of nitrogen alternative soybean meal for fattening Awassi lambs. Twelve Awassi lambs were used, aged from 4-5 months, with an average starting body weight of 23.5 ± 1.12 kg. Three formulation diets were randomly allocated to three groups of animals. Group one (T1 control) 14.7% soybean meal, Group two(T2) 7.5% soybean meal with 5.5% corn gluten meal and the third group 11% corn gluten meal.). All the experiment diets were supplemented with 1% salts and 1% vitamins and minerals with different levels of barley and wheat bran. Lambs were fed concentrate feed at 3% of live body weight on a dry matter basis with 200 grams of roughage (straw) day/lamb. The results of this study indicated that the addition of 11% corn gluten meal (T3) significantly ($p \leq 0.05$) higher in final weight, average daily gain, total weight and feed conversion efficiency compared to the addition of soybean alone or mixture soybeans with corn gluten meal. The results also showed that there were no significant differences in total protein concentration, albumin and globulin concentration, triglyceride concentration, cholesterol and urea concentrations in serum, and creatinine concentration.

© 2021 TJAS. College of Agriculture, Tikrit University

INTRODUCTION

Sheep considered one of the main sources of meat production in Iraq, the lamb's meat is palatable and favorable among Iraqi people more than other red meat-products. (Amen, 2014). As a result of the high prices of Soybean meal researchers were searched for alternatives to Soybean meal, corn gluten meal was founded to be the alternatives feed staff which recently used in Iraq especially in poultry diets. As these byproducts are characterized by being good sources of protein and energy, as well as vitamins dissolved in water in addition to mineral elements (Wang et al., 2007). It is also considered a good source of crude protein which contains 60-65% leads to reduce the addition of other protein materials to the diet and thus reduces the costs of feed (Shingoethe et al., 2009; Ji et al., 2012). Protein in the diets were the important sources for increasing the production of meat and milk. In this study, part or all of the corn gluten meal replaced the soybean meal in order to study the performance of the lamb's final weight, daily weight gain and feed conversion ratio, as well as the effect on some blood parameter.

Material and Methods

This study was conducted in the animal house of the College Veterinary Medicine - Tikrit University, for the period from 1st Desember 2019 to 9th February 2020 (70 days) excluding 14 days as an adaptation period to study the effect of using three sources of nitrogen on the productive

* Corresponding author: E-mail: Kaldunazez@gmail.com

performance of the Awassi lambs. This experiment was carried out using 12 Awassi lambs, aged between 4-5 months, with an average weight of 23.59 ± 1.12 Kg. They were distributed into three groups according to lamb's weight and each group included 4 lambs and the treatments were distributed randomly among the groups. The lambs were numbered with plastic numbers by ear and animals were housed in semi-open barns divided into individual cages of 1.5 x 1.5 square meters. They were equipped with feeders and drinkers, and salt blocks have been added in the cages. the feed provided to the animal was 3% of the live body weight on the basis of dry matter and served on two meals, the first at eight o'clock in the morning and the second at one o'clock in the afternoon with 200 grams of roughage (straw) day/lamb, the animals were weighed weekly and the amount of feed provided to them was adjusted with the change in the weights of the animals. As for the remaining feed, it was collected every morning, if present, to calculate the amount of feed intake. Clean water was created continuously, all lambs were fed experimental diets gradually for a period of 14 days for the purpose of acclimatization, after which the amount of feed intake was fixed, three experimental diets were used in this experiment, the first nitrogen source were 14.7% soybean meal, the second diet a mixture of 7.5% soybean meal and 5.5% corn gluten meal and the third diet 11% corn gluten meal.

Table (1) Percentages of the components of experimental deats (%)

Treatments compounds	T1	T2	T3
Black crushed barley	50	50	50
Wheat bran	33.3	35	37
Soybean meal	14.7	7.5	/
Corn gluten meal *	0	5.5	11
Salts	1	1	1
Vitamins and minerals	1	1	1

* P.N: Crude protein for Corn gluten meal= 60%, metabolizable energy 15.69 MJ/kg

Experimental Diets Chemical Composition

It was calculated according to the chemical analysis tables of Iraqi fodder materials according to Al-Khawaja, et al (1978). As for the corn gluten meal, its nutrient contents were calculated by the manufacturer.

Table (2) the chemical of the experimental diets composition

N.E treatments	Dry matter g / kg	Organic matter g / kg	Crude protein g / kg	Ether extract g / kg	Crude fiber g / kg	Ash g / kg	Nitrogen- Free Extract* g / kg	metabolizable energy ** MJ / Kg
T3	914.4	907.5	159.2	46.0	97	6.9	605.3	12.29
T2	903.6	896.67	160.4	42.2	89.7	6.93	604.37	12.14
T3	910.4	904.27	160.6	41.6	91.7	6.13	610.37	12.22
CGM	904	893	600	30.9	17	11	245.1	15.69

*Nitrogen-Free Extract (NFE)= OM –(CP+CF+EE)

** metabolizable energy was calculated according to (Maaf, 1975)

ME (MJ/Kg DM) = $0.012 \cdot CP + 0.031 \cdot EE + 0.005 \cdot CF + 0.014 \cdot NFE$

BLOOD TESTING

In the last week of the experiment, blood samples were taken after the animals were fasted for 12 hours, according to Jain et al., (1986). Then the samples were placed in a test tube (10 ml). Then the blood serum Samples was separated using a centrifuge at a speed of 4000 rpm for 10 minutes the blood serum was kept in a capacity of 6 cm in sealed packages and kept at- 20 ° C until

the analyzes were carried out. Blood samples were analyzed using ready-made (kit) fitted from a company Biolabo French and reading the samples with a spectrophotometer to estimate the total Protein, Urea, Cholesterol and Triglyceride.

Statistical analysis

The statistical analysis was performed using Complete Randomize Design (CRD) in one direction. As for the significance of differences between the parameters, the Duncan's multiple range test (Duncan, 1955) was used, and the ready-made statistical analysis program SAS (2001) was used. To analyze the data according to the following mathematical model: -

$$Y_{ij} = \mu + T_i + e_{ij}$$

As:

Y_{ij} = The view value of j for treated i.

μ = The general mean of the studied trait.

T_i = The effect of treatment i, as i = 1 (first), 2 (second), and 3 (third).

e_{ij} = Experimental error which assumes a normal and independent distribution with mean of zero and equal variance of σ^2 .

Results and discussion

Productive performance:

The results of the statistical analysis in Table (3) indicated that there were no significant differences ($P < 0.05$) between the different treatments in the initial weight, which was limited between (22.66 - 24.83 kg). Data presented in table (3) showed that average final body weight, total weight and average daily weight significantly ($P < 0.05$) improved for animal fed treatment contained 11% corn gluten meal as nitrogen source compared to the treatment soybean meal 14.7% and soybean meal 5.5% with 7.5% corn gluten meal.

Table (3) The effect of different nitrogen sources on the productive performance of the Awassi lambs (mean $\pm$ standard error)

Treated	Initial weight (Kg)	Final weight (Kg)	total weight (Kg)	Average daily weight gain (g / day)	Food intake (g / day)	FCR kg DM feed /kg weight gain
T1 SBM 14.7%	22.66 1.31 $\pm$	35.3 b 1.27 $\pm$	12.68b 0.46 $\pm$	181b 6.57 $\pm$	949.53b 41.70 $\pm$	5.26ab 0.32 $\pm$
T2 SBM 7.5%+ CGM 5.5%	23.28 ± 0.47	34.25 b ± 0.49	10.96 c ± 0.45	156 c 6.70 $\pm$	942.01b 14.11 $\pm$	6.40a 0.27 $\pm$
T3 CGM 11%	24.83 ± 0.50	39.89 a ± 0.34	15.06 a ± 0.35	215 a 5.06 $\pm$	1059.72a ± 12.82	4.93b 0.15 $\pm$
	NS	NS	*	*	*	*

The vertically different letters indicate significant differences with a probability level ($0.05 \geq P$), NS= no significant difference. - soybean meal =SBM, corn gluten meal = CGM

The reason for the improvement in the daily weight gain and the final weight rate for the third treatment (11% corn gluten meal) were may be due to a significant improvement in the amount of digested nutrients in digestibility trial and consumption of a greater amount of crude protein, ether extract, and crude fiber (Al-Azawie and Shujaa, 2021), Also due to a the resistance of corn gluten protein to digestion in the rumen and may be the reason for the improved in the third treatment (NRC, 1985). Also table (3) showed that the rates of feed consumed for lambs fed corn gluten meal (1059.73 g / day) increased significantly ($P < 0.05$) compared to the first treatment (949.55g/day) and second treatment (942.06 g /day). The reason may be due to the third treatment contained corn gluten substance (CGM), which led to an increase in the lamb's appetite for that diet. The third treatment also outperformed significantly ($P \geq 0.05$) in the Feed conversion ratio, as it reached 4.93 kg feed / kg weight gain compared to the first and second treatments, 5.26 - 6.40 kg feed / kg

weight gain respectively. The results of this study disagree with Capote (2013) and Al-Baldwin (2018), as they emphasized that the use of different protein sources was not lead to significant difference in Feed conversion ratio, Al-Baldawi (2018 was explained the reason that the animals had reached maturity). Al-Jubouri (2018), his results agreed with the results of this study, as well as with Al-Nasiri (2018). It is noticed through the results in Table (3) that the improvement in the performance of the lambs in the third treatment may be due to the diet containing the corn gluten meal, which is characterized by the high amount of protein crossing into the small intestine, reaching 55%, while the proportion of protein crossing into the intestine in the Soybean meal reached 22%, and this led to an increase in the efficiency of protein utilization, such as corn gluten, compared to soybean NRC (1985).

Blood chemical parameters

The results in Table (4) indicated that there were no significant differences ($P>0.05$) between the treatments in Total protein, Albumin concentration, Globulin, Triglyceride, Blood Urea and Creatinine but a mathematical increase in the protein concentration in general, especially the second and third treatment (1.38 ± 10.00 , 0.30 ± 9.05 g / 100 dl), which contained soybean meal and corn gluten meal compared to the first treatment (0.13 ± 8.57 g / dl blood) and that the increase in total protein concentration is due to the improvement in nitrogen absorption and the improvement and increase of the crude protein digestibility factor (Yousef and Zaki, 2001). The study agreed with Al-Nasiri (2018); Hussein (2018). The results in Table (4) also showed lowered in blood urea concentration with added 11% corn gluten meal though no significant difference ($P>0.05$) but may by the reason for the decrease due to an improvement in nitrogen absorption and an increase in the digestibility of crude protein (CP) (Yousef and Zaki, 2001), where the results agreed with Al-Mallah (2012); Nasser et al. (2014); Al-Nasiri (2018) and Al-Baldawi (2018) and disagree with both Hussein (2018) and Shahwaz (2019).

Table (4) the effect of different nitrogen sources on blood biochemical characteristics (mean $\pm$ standard error)

Treated	Total protein g / dl	Albumin g / dl	Globulin g / dl	Triglyceride mg/ dl	Cholesterol mg/ dl	Urea mg/ dl	Creatinine μ mol / L
T1 SBM 14.7%	8.57 $0.13 \pm$	3.57 $0.22 \pm$	4.99 $0.32 \pm$	16.70 $1.09 \pm$	85.37 $4.6 \pm$	52.25 $1.79 \pm$	0.439 $0.02 \pm$
T2 SBM 7.5% + CGM 5.5%	10.00 $1.38 \pm$	3.34 $0.05 \pm$	6.57 $1.40 \pm$	19.72 $3.50 \pm$	70.50 $9.91 \pm$	56.17 $3.17 \pm$	0.473 $0.11 \pm$
T3 CGM 11%	9.05 $0.30 \pm$	3.08 $0.16 \pm$	5.96 $0.41 \pm$	19.60 $0.42 \pm$	87.82 $8.42 \pm$	48.00 $3.29 \pm$	0.400 $0.02 \pm$

SBM= soybean meal, CGM = Corn gluten meal

CONCLUSION

It was concluded that fattening lamb given diet with 11% corn gluten meal improved performance of Awassi lamb (live body weight, Feed conversion ratio).

References

- A.O.A.C. (2002). Official Methods of Analysis. 17th Ed. Association of Official Analytic Chemists, Washington, DC.
- Al-Baldawi, H. A. J, (2018). The effect of using different protein sources on the production performance and some blood characteristics of the Awassi lambs. Master Thesis, College of Agriculture - Tikrit University.
- AI-Azawie, K.A. and T.A. Shujaa, (2021). effect of using three source of nitrogen on the productive performance and digestibility nutrients of Awassi lambs.

- Al-Jubouri, S.N. and T.A. Shujaa, (2018). The effect of feeding different proportions of crude protein sources on production performance and some measurements of Awassi lambs. Tikrit University Journal of Agricultural Sciences, Volume (18) Issue of Proceedings of the Seventh and First International Scientific Conference on Agricultural Research 10-11 April 2018.
- Al-Khawaja, A.K., A.A. Elham, and S. Al-Ahad, (1978). The chemical composition and nutritional value of Iraqi fodder materials. Ministry of Agriculture and Agrarian Reform Department of General Livestock. Baghdad .
- Al-Mallah, O.D.M, (2012). The effect of adding vitamin E and heat-treated soybean meal to diets on milk production, its components and some blood measurements in domestic ewes. Al-Rafidain Agricultural Journal, 40 (4): 154--163.
- Al-Nassiri, E.A.K.N, (2018). The effect of replacing soybean meal with yellow corn gluten on production performance and some characteristics of Awassi lambs and blood. Master Thesis, College of Agriculture - Tikrit University.
- Amin, M.A.M, (2014). The effect of using different percentages of hydrolyzed to non-degradable protein in fattening characteristics of Awassi lambs. Master Thesis, College of Agriculture - Tikrit University.
- Capote, K.L, (2013). The effect of corn gluten feed on performance and carcass characteristics of feeder lambs (Doctoral dissertation).
- Duncan, C.B, (1955). Multiple rang and Multiple “ F ” test. Biometric 11 : 1-12.
- Hussein, S.N, 2018. The effect of using different nitrogen sources on the productive performance of the Awassi lambs. PhD thesis, College of Agriculture - Tikrit University.
- Jain, N.C, (1986). Schalm veterinary hematology , 4th . Ed. Philadelphia : Lea and febiger.
- Ji, Y., L. Zuo., F. Wang, D. Li., and C. Lai, (2012). Nutritional value of 15 corn gluten meals for growing pigs: Chemical composition, energy content and amino acid digestibility. *Archives of animal nutrition*, 66(4): 283-302.
- MAFF, (1975). Ministry of Agric .and Fisheries and food dept. of Agric. energy allowances and feed systems for ruminants. Technical Bulletin,33.First published .
- Nasser, A.K., Q.Z.S. El Din., and N.Y. Abboud., (2014). The effect of using diets of different nitrogen sources on the growth and blood characteristics of local shrew calves grown under the local conditions of Nineveh governorate. Diyala Journal of Agricultural Sciences, 6 (2): 18-26.
- National Research Council (NRC), (1985). Nutrient requirements of sheep (Vol. 5). National Academies Press.
- SAS, (2001). SAS / STAT Users Guide for Personal Computers . Release . 6:12 . SAS Institute Inc ., Cary , N.C., U.S.A .
- Shahwaz, H.K, (2019). The effect of the use of different nitrogen sources on the growth and haematological parameters of the Awsid lambs. Master Thesis, College of Agriculture - Tikrit University.
- Shingoethe D.J., K.F. Kalscheur., A.R. Hippen., and A.D. Garcia, (2009). Invited review: The use of distillers’ products in dairy cattle diets. J. Dairy Sci. 92: 5802- 5813.
- Wang, Z., S. Cerrate., C. Coto., F. Yan., and P.W. Waldroup, (2007). Effect of rapid and multiple changes in level of distillers dried grains with solubles (DDGS) in broiler diets on performance and carcass characteristics Inter. J. Poult. Sci. : 6725-731.
- Yousef, H.M and A.A. Zaki, (2001). Effect of barley radical feeding on body weight gain and some physiological parameters of growing Friesian crossbred calves. Egy. J. Nutr. Feeds. 6: 465-471.

تأثير استخدام ثلاثة مصادر من النايتروجين في الاداء الانتاجي وبعض معايير الدم للحملان العواسية

طاهر عبد اللطيف شجاع

tshujaa@yahoo.com

خلدون عزيز حميد العزاوي

kaldunazez@gmail.com

قسم الإنتاج الحيواني كلية الزراعة – جامعة تكريت

الخلاصة

أجريت هذه الدراسة في البيت الحيواني التابع لكلية الطب البيطري - جامعة تكريت ، لمدة 70 يوماً سبقتها 14 يوم فترة تمهيديه اعتباراً من تاريخ 01/ 12/ 2019 ولغاية 9/ 2/ 2020/ سبقتها فترة تمهيديه 14 يوم , استعمل في هذه التجربة اثنا عشرة حملاً محلياً بعمر 4 – 5 أشهر وبوزن ابتدائي 23.5 ± 1.12 كغم. كونت ثلاثة علائق تجريبية ووزعت عشوائياً الى ثلاثة مجاميع من الحيوانات , المجموعة الاولى (سيطرة) غذيت فول الصويا بنسبة تضمنت 14.7% والمجموعة الثانية غذيت 7.5% كسبة فول الصويا مع 5.5% كلوتين الذرة والمجموعة الثالثة كسبة كلوتين الذرة بنسبة 11% كل العلائق التجريبية اضيف لها نسب مختلفة من الشعير ونخالة الحنطة وملح الطعام وفيتامينات ومعادن. غذيت الحملان علف مركز بنسبة 3% من الوزن الحي على اساس المادة الجافة بالإضافة الى 200 غرام تبين الشعير كعلف خشن. أشارت نتائج هذه الدراسة ارتفاعاً معنوياً ($P \leq 0.05$) في الوزن النهائي ومعدل الزيادة الكلية ومعدل الزيادة الوزنية اليومية وكفاءة التحويل الغذائي للمجموعة الثالثة (اضافة 11% كلوتين الذرة) مقارنة مع اضافة فول الصويا منفرداً وفول الصويا وكلوتين الذرة مجتمعا. وأشارت النتائج كذلك الى عدم وجود فروقات معنوية في كل من تركيز البروتين الكلي , الألبومين , الكلوبيولين , الدهون الثلاثية , كولسترول , يوريا في مصل الدم وتركيز الكرياتينين .

الكلمات المفتاحية:
كسبة كلوتين الذرة ، كسبة فول الصويا ، حملان عواسية .