



Mesopotamia Environmental Journal

ISSN: 2410-2598

Journal Home Page: <https://mej.uobabylon.edu.iq/index.php/mej/index>



DOI: <http://dx.doi.org/10.31759/mej.2022.6.1.0004>

Environmental and habitats diversity in Iraq, Part 1: Central region

Rihab Ghaleb Mohammed^{1*}, Fadhil Abbas Rhadi² & Nasrullah Rastegar-Pouyani³

¹Animal Production Department-College of Agriculture-University of Al-Qasim Green, Babylon, Iraq.

²Animal Production Department-College of Agriculture-University of Al-Qasim Green, Babylon, Iraq.

³Biology Department- College of Science- Razi University, Kermanshah, Iran.

*Corresponding Author: Rih1972@agre.uoqasim.edu.iq

To cite this article: Rihab Ghaleb Mohammed, Fadhil Abbas Rhadi & Nasrullah Rastegar-Pouyani, "Environmental and habitats diversity in Iraq, Part 1: Central region", Mesop. environ. j., 2022, Vol. 6, No.1, pp.25-36

Received Date: 15 / 11 / 2021, **Accepted Date:** 20 / 4 / 2022, **Publishing Date:** 6/5/2022



ROAD

This work is licensed under a [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Abstract

The area of Iraq is about 437.065 km², its territory encompasses the Mesopotamian alluvial plain, the northwestern end of the Zagros mountain range, and the eastern part of the Syrian desert, includes nine to eleven main terrestrial ecoregions, some of which can be classified as key ecoregions that covers larger areas of the country. Presence of these ecoregions resulted in a considerable environmental and habitat diversity. The present study includes 23 locations throughout 5 central governorates in Iraq: Babylon, Holy Karbala, Al-Najaf alashraf, Qadisiya, Wasit, these locations extended between the geographical coordinates 30° 48' 19.36" -32° 41' N and 43° 29' 25.89" -45° 37' 38" E, with the altitude (minimum: 10; maximum: 344 and mean: 50.25± 73.062 m) above sea level. The study was carried out from 20th Sep.2013 to 3rd Nov. 2015.

Key words: Habitat; Environment; Ecoregions; Central; Iraq.

1- Introduction

Eight ecozones according to (1, 2) Nearctic, Palearctic, Afrotropic, Indomalaya, Australasia, Neotropic, Oceania, and Antractic. The Palearctic, is the largest of the eight ecozones that constituting the Earth's surface, it includes the terrestrial ecoregion of Europe, Asia, North of the Himalaya foothills, northern Africa, and the northern and central parts of the Arabian Peninsula. The Palearctic ecozone consists of five smaller ecozones or subregions: the European Siberian; the Mediterranean Basin; the Sahara and Arabian Deserts; Western and Central Asian; and China and Japan. Two major rivers in the subregion western Asia of the Palaeartic are the Tigris, rises in the Taurus mountains of eastern Turkey and the Euphrates, rises in the mountains of Anatolia. Between these rivers is an ancient area called Mesopotamia, which was also known as the fertile crescent. The two rivers join together near Al Qurna in southern Iraq and flow to the Arabian gulf (3 and 4). The macro-ecosystems of the Iraq have been defined by the World Wildlife Fund & Nature Conservancy in their development of terrestrial, Freshwater, and marine "ecoregions" of the world, with their associated species and conservation status (867 terrestrial ecoregions are categorized within 14 biomes and eight biogeographic realms) (5). There are five terrestrial biomes found in Palearctic realm of Iraq: Temperate broadleaf and mixed forests; Temperate grasslands, Savanas and shrublands; Flooded grasslands and Savannas; Mediterranean forests, woodlands and scrub; Deserts and Xeric shrublands. Iraq includes nine to eleven main terrestrial ecoregions, some of which can be classified as a key ecoregion that covers larger areas of the country, we excluded the ecoregions that are of very weak influence in the Iraqi habitats, and there is no large representation of the elements characterizing these ecoregions in Iraq and focusing only on that kinds of ecoregions, and this is based on the existing information (6,7, 8). The Freshwater and marine ecoregions of the world are also represented in Iraq including three different freshwater ecoregions (Arabian Interior, Lower Tigris and Euphrates, Upper Tigris and Euphrates, and one marine ecoregion (Arabian Gulf, which is part of the Western Indo-Pacific Realm. The presence of these ecozones resulted in a considerable biodiversity of both flora and fauna of Iraq. (9, 10, 11, 12)

2- Materials and Methods

• Study Area

The present study surveyed an area included 23 locations distributed in five governorates throughout central region in Iraq depending on an administrative maps: Babylon, Holy Karbala, Holy Najaf, Qadisiya, Wasit. The locations lied within the geographical coordinates 30° 48' 19.36" -32° 41' N and 43° 29' 25.89" -45° 37' 38" E, and the altitude (10-344 m) above sea level which detected by using Garmin GPS map 60CSx device (Fig. 1). The study was carried out during the period from 20th Sep. 2013 to 3rd Nov. 2015. Photographs have been taken for all habitats (Plates 1-4), and all the ecological information for the visited sites have been provided according to literatures and environmental office related to each governorate, as below:

Babylon Governorate: Seven locations were surveyed (Plate 1: A-F), include:-

- Khagan village, Al-Showmali subdistrict \Hashimiya district: 32° 22' 00.3" N, 044° 46' 07.4" E.
- Al-Ubaid village, Al-Qasim subdistrict\Hashimiya district: 32° 13.729' N, 044° 33.913' E.
- Al-Hilla city\Hilla district: 32° 28' 43.2" N, 044° 24' 59.2" E.
- Abi Gharaq subdistrict \Hilla district: 32° 32.016' N, 044° 20.782' E.
- Al-Naby Ayub village\ Hilla district: 32° 20.767' N, 044° 23.863' E.
- Al-Nikhealah village\ Hilla district: 32° 25.776' N, 044° 29.901' E.
- Ibraheem Al-Khaleel village\ Hilla district: 32° 23.624' N, 044° 20.793' E.

AL-Najaf alashraf Governorate: Eight locations were surveyed (Plate 2: A-F), include:-

- Bahr Al-Najaf \AL-Najaf district: 31° 52' 30.5" N, 044° 15' 47.2" E.

- Madhlum village, Bahr Al-Najaf\Al-Najaf district: 31° 53' 37.6" N, 044° 16' 25.4" E.
- Um Groon called area \ Al-Najaf district: 31° 46' 17.3"N, 044° 13' 37.8"E.
- AL-Shabaka(Shbicha)subdistrict\Al-Najaf district: 30° 48' 19.36" N, 43° 40' 5.87" E.
- Al-Jaiyar area called\ AL-Mushkhab district: 31° 50' 28.4" N, 044° 30' 13.7" E.
- Al-Kufa city \ Kufa district: 32° 8' 05.7" N, , 44° 22' 17.3"E.
- Al-Qizwiniya,Al-Abbasiya subdistrict\Kufa district: 32° 04' 50.0"N, 044° 26' 46.3"E.
- Al-Abbasiya, Al-Abbasiya subdistrict\ Kufa district: 32° 04' 50" N, 044° 26' 46.3" E.

Holy Karbala Governorate: Four locations were surveyed (Plate 3: A-C), include:-

- Ain- Tamor district: 32° 33' 56.52" N, 43° 29' 25.89" E.
- Al-Wadi (Cemetery)\Kerbala district: 32° 33.074' N, 044° 04.402'E.
- Al- Kafeel village\ Kerbala district: 32° 32' 53.6" N, 044° 05' 39.4".
- Al-Razzaza lake\ Ain- Tamor district: 32° 41' N, 43° 40' E.

Wasit Governorate: One locality was surveyed (Plate 4: A), include:-

- Haur Al-Dalmaj\Kut district: 32° 07' 55.3" N, 45° 37' 38" E.

AL-Qadisiya Governorate: Three locations were surveyed (Plate 4: B-D), include:-

- Diwaniya district: 31° 56' 28.8" N, 044° 54' 12.6" E.
- Al-Shamiya district: 31° 58' 17.1"N, 044° 36' 36.4" E.
- Hamza district: 31° 43' 42.1"N, 044° 57' 35.6"E.
- **Statistical Analysis**

Minimum, maximum, mean and standard deviation values for altitude data, have been calculated for all the visited locations by using Excel program (13).



**A) Khagan village , Al-Showmali subdistrict
Hashimiya district \ Babylon Province**



**B) Abi-Gharaq subdistrict, Hilla district
Babylon Province**



**C) Al-Ubaid village, Al-Qasim subdistrict
Hashimiya district\ Babylon Province**



**D) Al-Nikheala village, Hilla district
Babylon Province**



**E) Al-Nabi Ayub village, Hilla district
Babylon Province**



**F) Ibraheem Al-Khaleel village
Hilla district \ Babylon Province**

Plate 1. (A-F) Habitats of the visited locations in Babylon governorate.



**A) BahrAl-Najaf, Al-Najaf district
Najaf Province**



**B) Madhlum village, BahrAl-Najaf
Al-Najaf district\ Najaf Province**



**C) Um-Groon area, Al-Najaf district
Najaf Province**



**D) Al Jaiyar village, Al-Mushkhab district
Najaf Province**



**E) Al Kufa city, Al-Kufa district
Najaf Province**



**E) Al Abbasiya subdistrict, Al-Kufa district
Najaf Province**

Plate 2. (A-F) Habitats of the visited locations in Al- Najaf alashraf governorate.



**A) Razzaza Lake, Ain Tamor district
Karbala Province**



**B) Al-Kafeel village, Kerbalaa district
Karbala Province**



**C) Al-Wadi (Cemetery), Kerbalaa district
Karbala Province**

Plate 3. (A-C): Habitats of the visited locations in Holy Karbalaa governorate



Plate 4. Habitats of the visited locations in governorates: (A) Wasit, (B-D) Qadisiya.




Fig. (1) Map of Iraq showing the 23 studied locations according to their geographical coordinates:- **Karbala'a**:- 1: Ain- Tamor district; 2: Al-Razzaza lake; 3: Al-Kafeel village-Kerbala district E; 4: Al-Wadi (Cemetery)-Karbala'a district; **Babylon**:- 5:Al-Ubaid village-Al-Qasim subdistrict-Hashimiya district; 6: Al-Showmali subdistrict - Khagan village- Hashimiya district; 7: Al-Hilla city, Hilla district; 8: Al-Naby Ayub village - Hilla district; 9: Al-Nikhealah village-Hilla; 10: Abi-Ghraq subdistrict, Hilla district; 11: Ibraheem AL-Khaleel village Hilla district; **Wasit**:- 12: Al-Dalmaj; **Qadisiya**:- 13: Diwaniya district; 14: Hamza district; 15: Al-Shamiya district. **Holy Najaf**:- 16: Bahr Al-Najaf, Najaf district; 17: Madhlum village, Bahr Al-Najaf; 18: Al-Jayir village, Al-Mushkhab district; 19: Al-Kufa district; 20: Al-Abbasiya county, Al-Kufa district; 21: Al-Qizwiniya, Al-Abbasiya county, Al-Kufa district; 22: Um Groon called area Al-Najaf district; 23: Al-Shabaka, Al-Najaf district.

3- Results and Discussion

Twenty three locations distributed in five governorates throughout central region in Iraq were visited: Babylon, Holy Karbala, Al-Najaf alashraf, Qadisiya, Wasit. The locations lied within the geographical coordinates 30° 48' 19.36" -32° 41' N and 43° 29' 25.89" -45° 37' 38" E, and the altitude (minimum: 10; maximum: 344 and mean: 50.25± 73.062 m) above sea level, according to the figure (1) and the plates an environmental description for each governorate can be introduced depending on visited locations:-

Babylon: Seven locations were visited (30° 37' 56.5" - 32° 32.016' N and 044° 20.793' - 46° 41' 16.1" E) with elevation (minimum: 10, maximum: 148 and mean: 59 ±57.066 m) from sea level with their varied habitats from deserted areas with sand dunes and scattered oil fields civilized area consists of many houses with gardens, village with agricultural areas and some sub arid areas. Babylon is a governorate in central Iraq, it has an area of 5.603 km² with geographical coordinates: 32° 37' N and 44° 33' E, the provincial capital is

the city of Hillah on the branch of Euphrates river. It is situated in a predominantly agricultural region, which is extensively irrigated with water provided by the Hillah canal, producing a wide range of crops, fruit and textiles. The river runs exactly in the middle of the town, and it is surrounded by date palm trees and other forms of vegetation enhancing the weather and reducing the harmful effect of dust and the desert winds. (14). Also it located in both Tigris-Euphrates alluvial salt marsh (PA0906), Although bounded by deserts and dry shrubland, this region is characterized by marshlands and seasonally inundated plains in a basin covered by recent (Pleistocene and Holocene) alluvial deposits and forming a vast inland delta of the Euphrates, Tigris, and Karun rivers. The climate is subtropical, hot and arid (15) and the ecoregion Arabian Desert & East Sahero-Arabian Xeric Shrublands (PA1303) is a region of little rain fall (annual precipitation is 125 mm) with occasional oases. Daily temperature varies considerably reaching over 45 C° or higher in July and Aug. and dropping to the teens in Jan. and Feb. Occasional brackish salt flats also exist in some areas (16 and 17).

Holy Karbala'a: Four locations were visited (32° 31' - 32° 31' N and 43° 26'- 044° 05' 39.4" E) with an elevation (minimum 18, maximum 45 and mean 28.25± 57.066 m) and their habitats diverted from large wetland and the vast area of the lake vicinity represented by gypcrete, inland sabkha, depression fill, flood plain and aeolian sediments, civilized area consists of many houses with gardens; some sub arid and arid areas, arid and sub arid areas used for burying dead people, and village with cultivated agricultural fields and some sub arid areas. Karbala'a located about 105 km southwest of the capital Baghdad at 32° 27' N, 43° 48' E, it lies on the edge of the desert west of the Euphrates river and on the left side of Hussainiyya canal, Karbala'a experiences a hot desert climate with extremely hot, dry summers and cool winters (18-20). Almost all of the yearly precipitation received between Nov. and Apr. though no month is truly wet. The climate is damp, very hot in summer, and mild in autumn; its land is watered by the canals of the Euphrates river, it has many orchards, flat lands suitable for various forms of agriculture, and all sides of the city are surrounded by fruit trees and palm groves. Karbala'a resembles Babylon, it locates in both same ecoregions with occasional brackish salt flats exist in some areas, a few of which in Iraq have been utilized for water storage (e.g. Razaza lake) (9). The Bedouins (nomads) cross the region with seasons seeking pasture for their herds of goat, sheep and camels.

Al-Najaf alashraf: Eight locations were visited extended between 30° 48' 19.36"-32° 8' 05.7"N and 43° 40' 5.87"-044° 30' 13.7"E with elevation ranged from (minimum:15, maximum:334 and mean:59.66 ± 113.076 m) from sea level, and with habitats varied from wetland depression area, vast desert or semi desert areas, marsh-like area with limited cultivated orchards beyond and surrounded by vast desert or semi desert areas, sandy dunes and flat sands plains with scattered vegetation, and agricultural areas with livestock grazing, and civilized areas consists of many houses with gardens. Al-Najaf alashraf located in the southwestern portion of Iraq at 29° 50' latitude and 32° 21' longitude to the North and 42° 50' and 44° 44' longitude to the East, with total area of 28824 km², 5% of the region lies in the sedimentary plateau while the rest is located in the western hill. It lies on the shortest path between the abundantly producing fertile hill and the desert dune, known for its rich raw materials (21). The weather is dry and hot (42°C) in summer; cold (18°C) and less rainy in winter, because of its location in the ecoregions: Arabian Desert & East Sahero-Arabian Xeric Shrublands, Red Sea Nubo-Sindian Tropical Desert and Semi-Desert (PA1328) that occurs along the Saudi border and extend slightly into Iraq (22 and 23).

AL-Qadisiya: Three locations were visited extended between 31° 43' 42.1"- 31° 58' 17.1" N and 044° 36 36.4"- 044° 57' 35.6" E , with an elevation ranged (minimum:35, maximum:55 and mean: 43.33±10.408m)

from sea level and their habitats are diversified from civilized and cultivated areas consists of many houses with gardens and some sub arid areas to arid and sub arid areas. Qadisiya locates between the lines along the 48° 24'- 48° 45' and horizontal lines 31° 7'-32° 24', it lies within easy sedimentary areas, and the part within the plains of the delta, also it resembles Babylon governorate where it locates in both same ecoregions, and considers a rural area with many plants are growing there such as rice, barley and dates (24), also, there is Shammiya river which is a branch of Euphrates splits the city. At 30 km to the south there are a lots of date palms and it splits by Hamza river (25, 26, 16).

Wasit: Is located in the central part of Iraq in the ecoregions: Mesopotamian Shrub Desert (PA1320) desert ecosystem, has little biological diversity; Tigris-Euphrates alluvial salt marsh; Arabian Desert & East Sahero-Arabian Xeric Shrublands; and South Iran Nubo- Sindian Desert and Semi-desert. It lies between the line along the 45° 45' and horizontal line 32° 40' and its area measures 17.153 km², most of it fit for agriculture, has irrigation sources as Tigris river and the most important crops harvested include wheat, barley, maize, rice, cotton, sunflower as well as other food products, namely vegetables and fruit, which cover large areas (27). Most farmlands use irrigation, whether directly or through implements. Average annual rainfall is 385 mm. Moreover, the region is flat and its climate alternates between that of the Mediterranean and that of the desert, which is hot and dry. Haur Delmaj the only site was visited with a geographical coordinates 32° 07' 55.3" N and 45° 37' 38" E with an elevation 13 m from sea level, this site listed as a wetland of international importance by (28), and has been identified as an "Important Bird Area" by Bird Life International. Haur Delmaj, is famous marsh that have a several types of yearly migrate birds came from Asia and Europe and was listed by (29, 30) as an important area for the Iraqi biodiversity since it lies within a semi-desert area and comprises terrestrial and aquatic habitats allowing a wide range of biodiversity components. The vertebrates are the most prominent group in the marsh especially birds and fishes and for a lesser extent reptiles and mammals. However, it is still poorly studied and further scientific work is required to understand the biodiversity and the relationships among the biotic and abiotic factors in the area (11, 31).

4- Conclusion

Current study is pointed that in an environment which varies in terms of topographical and vegetation structure it would create a mosaic of temperature and humidity within a relatively small area throughout the day. Anthropogenic activity (unmanaged grazing, plowing of lands, infrastructural developments, and clearance of forests) resulted in the decrease in the size of some populations and/or their diversity. In general, there are several abiotic and biotic parameters such as temperature, moisture (through rainfall and/or dew), wind speed and vegetation cover are affecting the diversity and ecology of both flora and fauna in Iraq

5- Acknowledgement

We are grateful to Iraqi ministry of higher education and scientific research. Also, we would like to thank the authorities of Al-Qasim Green University/ college of Agriculture /Animal production department.

6- References

- (1) WWF. World Wide Fund for Nature. (2014). Ecoregions. Retrieved from worldwildlife.org/terrestrial-ecoregions. (Accessed: Jan. 22, 2014).
- (2) WWF. World Wildlife Fund. (2006). Ecoregions, ver. Jan-06. (Online). Available: <http://gis.wwf.org/wildfinder>. (Accessed: Sept. 14, 2006).

- (3) Coad, B. W. (2010). Lower Tigris and Euphrates, ecoregion description of the freshwater ecoregions of the world, Website, (Online). Available: <http://www.feow.org/ecoregion.details>. (Accessed: Jun. 3, 2010).
- (4) Schultz, J. (2002). The ecozones of the world: the ecological divisions of the geosphere. 2nd ed., Translated into English by Bridget Ahnert. Eugen Ulmer GmbH & Co. Stuttgart, Germany. pp.430.
- (5) Eric, D.; Olson, D. and Graham, D. J. A. (1995). Conservation assessment of the terrestrial ecoregions of Latin America and the Caribbean. World Bank, Washington DC. pp. 129.
- (6) Al-Lami, A. (2012). Action plan for implementing the programme of work on protected areas of the convention on biological diversity: Iraq. Ministry of Environment of Iraq, pp 47.
- (7) Malinowski, J. (2002). Iraq: A geography. United States Military Academy West Point, NY., pp 125.
- (8) Guest, E. (1968). Flora of Iraq. The introduction, Vol. 1. Baghdad: Ministry of Agriculture of the Republic of Iraq. pp.349.
- (9) Scott, D. A. A. (1995). Directory of Wetlands in the Middle East. IUCN, Gland, Switzerland, Slimbridge. pp560.
- (10) Nicholson, E. and Clark, P. (2002). The Iraqi marshlands: A human and environmental study. 2nd ed., London, United Kingdom, The AMAR International Charitable Foundation. pp.332.
- (11) Evans, M. I. (2004). saving mesopotamian marshes of Iraq”, environmental training programme for Iraqi biologists- Waterloo. Canada. pp.229.
- (12) Garstecki, T. and Amr, Z. (2011). Biodiversity and ecosystem management in the iraqi marshlands screening study on potential world heritage nomination. IUCN ROWA, Jordan. Pp.189.
- (13) Fowler, J. ; Cohen, L. and Jarvis, P. (1998). Practical Statistics for Field Biology. 2nd Ed. John Wiley & Sons. Baffins Lane, Chichesters. England, pp.259
- (14) Hassan, N. O. (2010). Iraqi fourth national report to the convention on biological biodiversity. Ministry of Environment. 153 pp.
- (15) Bozkurt, D. and Sen, O. L. (2013). Climate change impacts in the Euphrates–tigris basin based on different model and scenario simulations. Journal of Hydrology, vol.480, pp. 149–161.
- (16) IMoE, Iraqi Ministry of Environment. (2010). National report on biodiversity in Iraq. Fourth national report to the convention on biological diversity (CBD). pp.153.
- (17) Bachmann, A; Chappell, N. and Matti, N. (2011). Key Biodiversity Survey of Iraq 2010. Site Review. Nature Iraq & the Iraq Ministry of Environment, Kurdistan, Iraq. pp.102.
- (18) Al-Dabbas, M. A; Al- Zubaidi, A. A. and Al- Khafaji, R. (2015). Impact of climate changes on the hydrochemistry of Razaza lake and Rahaliya – Shithatha springs – central Iraq. Journal of Environment and Earth Science, vol.5, no. 6, pp.106-116.
- (19) Jassim, R. and Al-Zubaidi, A. (2013). Hydrochemical assessment of water resources southern part of Al-Razzazah lake and surroundings, Karbala area, central Iraq, Iraqi Bulletin of Geology & Mining, vol.9, no.2. pp. 81-95.
- (20) IMoWR. Iraqi Ministry of Water Resources. (2005). Iraq Reconstruction – Water Resources Sector. Speech by the Minister of Water Resources in USA.
- (21) Benni, T. J. and Al-Tawash, B. S. (2011). palynological evidences on paleoclimate and paleo-environmental changes during late quaternary of bahr Al-Najaf depression, central Iraq. Iraqi Bulletin of Geology and Mining, vol.7, no.2, pp. 1-28.

- (22) Al-Atia, M. J. (2006). The Najaf Land: History, geological heritage and natural wealth. 1st ed, Al-Nibras foundation for printing, publication and distribution. Al-Najaf. pp, 9993.
- (23) Al-Hakeem, H. E. (2004). Bahr Al-Najaf and its role in joining Iraq with the world. Journal of Kufa Studies Center, vol.1, no.2, pp. 7-22.
- (24) IMoE. Iraq Ministry of Environment & Nature Iraq. (2014 a). Inventory of key biodiversity areas of Iraq. Baghdad, Iraq: Iraqi Ministry of Environment & Nature Iraq. In preparation. pp.48.
- (25) IMoE Iraq Ministry of Environment. (2012). Program of work for protected areas. (POWPA). pp.48.
- (26) Zohary, M. (1973). Geobotanical foundation of the middle east. Vol 1-2. Gutav Fische Verlag. Germany.
- (27) Carp, E. A. (1980). Directory of Western Palearctic Wetlands. Nairobi/Kenya and Gland/Switzerland, UNEP and IUCN. pp.517.
- (28) George, P. V. and Viellard, J. (1970). Midwinter observations on birds of central and south Iraq". Baghdad, Iraq. Iraq Natural History Museum Bull., Vol. IV, no. 4.
- (29) Nature Iraq. (2013). (Online) Available: <http://www.natureiraq.org>.
- (30) Hassan, F. M. Taylor; W. D.; Al-Tae, M. S. and Al-Fatlawi, H. J. J. (2010). Phytoplankton composition of Euphrates river between Al-hindiya barrage and Kifil city, Iraq. J. Environ. Biol., 31: 343-350.
- (31) Scott, D. A. and Carp, E. A. (1982). Midwinter survey of wetlands in Mesopotamia, Iraq. Sandgrouse, vol. 4, pp. 60-76.