

Prevalence and Antibiotic Resistance in *Aeromonas* species Isolated from Common Carp (*Cyprinus carpio* L.) Cultivated in Floating Cages at Al-Hilla River

Sadeq M. Al-Haider¹, Khalidah S. Al-Niaeem^{2*} & Amjed K. Resen²

¹ College of Veterinary Medicine, University of Al-Qasim Green, Iraq;

² Department of Fisheries and Marine Resources, College of Agriculture, University of Basrah, Basrah, Iraq

Received 15 March 2019; Accepted 24 May 2019; Available online 30 December 2019

*Corresponding author: kalidah_alsalim@yahoo.com

Abstract: The aim of this study is describe the case of ulcer and hemorrhaging (red-sore disease) in a common carp (*C. carpio*) that is cultivated in floating cages at Al-Hilla river and to finding efficient antibiotic treatments. A total of 144 specimens of the common carp were examined for ulcer with hemorrhaging during the period from December 2017 till the end of November 2018. In this study, red-sore disease was found in 45 specimens. This disease is caused by the infection of Motile *Aeromonas* Septicemia (MAS). *Aeromonas hydrophila*, *A. sobria* and *A. veronii* were isolated from the skin, gills and intestine of common carp. *Aeromonas* species were identified with a different percentage in common carp as the follows: *A. hydrophila* (58.76%), *A. sobria* (31.83%) and *A. veronii* (43.3%). Antibiotic susceptibility test of 12 antibiotics (Piperacillin/ Tazobactam, Amikacin, Cefazolin, Gentamicin, Cefoxitin, Ciprofloxacin, Ceftazidime, Levofloxacin, Cefepime, Tigecycline, Imipenem and Trimethoprim/ Sulfamethoxazoleby) showed variable resistances for *Aeromonas* species. *Aeromonas* species were identified and examined for antibiotic susceptibility using the VITEK 2 system.

Keywords: Fish, *Aeromonas* spp., antibiotic susceptibility, Iraq.

Introduction

Aeromonas species cause a wide spectrum of disease syndromes among warm and cold blooded animals, including fish, reptiles, amphibians, mammals and humans (Janda & Abbott, 2010). The members of *Aeromonas* spp. belong to the family Aeromonadaceae comprise Gram negative, nonspore forming and motile bacilli or coccobacilli rods with rounded ends (Skwor *et al.*, 2014). It is cause

most serious infectious disease known as Motile *Aeromonas* Septicemia (MAS), associated with high mortality rates without symptoms in per acute phase but in acute phase, skin and fin ulcers appear on the external surface of fish with ascites in the abdomen and exophthalmia (Bondad *et al.*, 2005; Austin & Austin, 2012; Skwor *et al.*, 2014; Al-Niaeem *et al.*, 2015). The disease

can produce significant losses in aquaculture industry because of reduced growth in survived cases (Pachanawan *et al.*, 2008; Buján *et al.*, 2010). The *Aeromonas* spp. consist of 27 species, but some such as *A. hydrophila*, *A. allosaccharophila*, *A. jandaei*, *A. piscicola*, *A. salmonicida*, *A. schubertii*, *A. bestiarum*, *A. sobria*, *A. veronii*, *A. dhakensis* and *A. caviae* are considered pathogenic to animals and humans (Beazhidalgo & Figueras, 2013; Aravenaromán *et al.*, 2014; Li *et al.*, 2019).

Outbreaks of the disease and its influence on the survival, persistence and ability of *Aeromonas* to cause infection are usually related with the change of environmental state, such as transfer the fish to poor water quality, stress, mishandling, changing of temperature, dioxide levels, overcrowding, high nitrite and carbon (Dixon, 1993; Aoki, 1999; Al-Taei *et al.*, 2017).

Because, there is a little information pertains to *Aeromonas* in the fish farms particularly in Iraq. The aim of the present to describe a case of hemorrhaging and skin ulcer in common carp, *C. carpio* cultivated in floating cages in Al-Hilla river and the susceptibility pattern of bacteria to 12 antimicrobial drugs.

Materials & Methods

A total of 144 fish of common carp were sampled from four farms of floating cages at Al-Hilla, the first and second stations, before the city center and the third and fourth stations, after the city center. During the period December 2017- November 2018. The length of fish ranging between 23.8- 37.16 cm and the weight was 261.66- 693.3 gm. The live fishes were transported to oxygenated pond water. Before transferred to the laboratory in College of Veterinary Medicine, University of Al-Qasim Green.

The collected fishes were dissected and bacterial swabs were taken aseptically using a sterile loop from skin, gill and intestine. For isolation of bacteria, MacConkey Agar medium was used. The inoculated plate was incubated at 37 °C for 24 h. Bacteria were identified and antibiotic susceptibility (Piperacillin/ Tazobactam, Amikacin, Cefazolin, Gentamicin, Cefoxitin, Ciprofloxacin, Ceftazidime, Levofloxacin, Cefepime, Tigecycline, Imipenem and Trimethoprim/ Sulfamethoxazole) by using the VITEK 2 system.

Results & Discussion

The results of this study showed that *A. hydrophila*, *A. sobria* and *A. veronii* isolated and identified from the four floating cages in Al-Hilla river are a dangerous and may be unhealthy for public health when consumed. *Aeromonas* species were identified with different percentages for common carp, including *A. hydrophila* (58.76%), *A. sobria* (31.83%) and *A. veronii* (43.3%) (Table 1). The Motile *Aeromonas* Septicemia is one that causes main disease problems in the carp farm (Cipriano, 2011). Outbreaks of the infectious disease are usually caused the changing of environmental state and stress. Fluctuation of temperature (sudden), crowding, poor water quality, low dissolved oxygen, high ammonia levels are the common factors related with MAS (Ko *et al.*, 1998).

The study was conducted from December 2017 till the end of November 2018 in which there are fluctuations in water quality parameters in the aquaculture in the studied stations (Table 2). The mean of temperature fluctuated from 10.7 °C to 32.9°C. the mean of salinity was recorded ranged from 460 to 580 PSU, ammonia was recorded ranged from 1.3 to 2.8 PSU. The pH value was relatively from 6.4 to 8.1. In culture system of fish that

always exposed an assortment of stresses become over sensitive to disease infection (Rijnsdorp *et al.*, 2009; Albert & Ransangan, 2013).

The growing of bacteria in water increased by rising levels of water temperature from 25 °C to 32°C, organic matters, salinity and pH

5-9 (Kiriratnikom *et al.*, 2000). These levels of growing bacteria were observed in the current study.

Gram negative bacilli, *A. hydrophila*, *A. sobria* and *A. veronii* were isolated from the skin, gills and intestine from common carp (Fig. 1 & table 3).

Table (1): Percentage of isolated *Aeromonas* spp.

Station	<i>A. hydrophila</i> (%)	<i>A. sobria</i> (%)	<i>A. veronii</i> (%)
1	11.76	5.88	9.80
2	15.60	8.26	11.0
3	14.40	7.69	11.5
4	17.0	10.0	11.0
Total	58.76	31.83	43.3

Table (2): The water parameters of studied stations.

Station	Temp (°C)	Salinity (PSU)	Ammonia (mg.l ⁻¹)	pH
	Min-max (mean)	Min-max (mean)	Min-max (mean)	Min-max (mean)
	Range (mean)			
1	11.9 - 32.7 (22.66)	480 - 560 (517.5)	1.3-1.8 (1.62)	6.8 - 8.1 (7.36)
2	10.7 - 32.9 (22.9)	460 - 570 (516.66)	1.4-1.9 (1.69)	6.5 - 8.1 (7.33)
3	13 - 32.6 (23.8)	480 - 580 (530.8)	2.0-2.8 (2.42)	6.4 - 7.6 (7.09)
4	12.5 - 31.2 (22.86)	460 - 560 (513.33)	0.7-1.4 (1.21)	7.1 - 7.9 (7.35)


Biochemical characterization of Motile *Aeromonas* Septicemia using the VITEK 2 system was given in tables (4-6). It has been noticed by some researchers that *Aeromonas* spp. causes skin lesions with hemorrhagic in fresh water fish and cultured fish (Aguilera *et al.*, 2005; Beazhidalgo & Figueras, 2013; Skwor *et al.*, 2014; Gao *et al.*, 2016; Li *et al.*, 2019).

The best antibiotics susceptibility were Tigecycline, Ceftazidime, Cefepime and Gentamicin, when tested in vitro on *A.*

hydrophila while, it was resistant to Cefazolin and Ceftriaxone (Table 7).

In vitro, the more antibiotic susceptibility to *A. sobria* were Ciprofloxacin, Levofloxacin, Tigecycline, Gentamicin, Cefazolin and Cefepime

whereas *A. sobria* had intermediately resistant to Imipenem (Table 8). The antibiotic susceptibility in vitro were more effect on *A. veronii* Ciprofloxacin, Levofloxacin, Tigecycline, Gentamicin, Cefazolin and Cefepime (Table 9).



ce

Table (3): The infected organs of fish by *Aeromonas* spp.

Month	Skin			Gills			Intestine		
	<i>A. hydrophila</i>	<i>A. sobria</i>	<i>A. veronii</i>	<i>A. hydrophila</i>	<i>A. sobria</i>	<i>A. veronii</i>	<i>A. hydrophila</i>	<i>A. sobria</i>	<i>A. veronii</i>
December 2017	-	-	-	-	-	-	+	-	-
January 2018	+	+	-	-	+	-	-	+	-
February	-	+	-	-	+	-	-	+	-
<u>March</u>	+	+	-	+	-	-	+	+	+
<u>April</u>	+	-	+	+	-	+	+	-	+
<u>May</u>	+	-	+	+	-	+	+	-	+
<u>June</u>	+	-	+	+	-	+	+	-	+
<u>July</u>	+	+	+	-	+	+	-	+	-
<u>August</u>	+	-	+	-	-	+	-	-	+
<u>September</u>	+	-	+	+	-	+	+	-	+
<u>October</u>	+	-	+	-	-	+	+	-	+
<u>November</u>	-	-	+	-	+	+	+	-	+

+: Positive, -: Negative

Table (4): Biochemical features of isolated *A. hydrophila*.

Biochemical features			Reaction	Biochemical features			Reaction
2	APPA	-		3	ADO	-	
4	PyrA	-		5	IARL	-	
7	dCEL	+		9	BGAL	+	
10	H ₂ S	-		11	BNAG	+	
12	AGLTp	-		13	dGLU	+	
14	GGT	-		15	OFF	+	
17	BGLU	-		18	dMAL	+	
19	dMAN	+		02	dMINE	+	
01	BXYL	-		00	BAlap	-	
03	ProA	-		26	LIP	+	
07	PLE	-		09	TyrA	+	
31	URE	-		30	dSOR	+	
33	SAC	+		34	dTAG	-	
35	dTRE	+		36	CIT	-	
37	MNT	-		39	5KG	-	
42	ILATk	-		41	AGLU	-	
40	SUCT	+		43	NAGA	-	
44	AGAL	-		45	PHOS	+	
46	GlyA	-		47	ODC	-	
48	LDC	-		53	IHISa	-	
56	CMT	+		57	BGUR	-	
58	O129R	+		59	GGAA	+	
61	IMLTa	-		60	ELLM	+	
64	ILATa	-			-		

+: Positive, -: Negative

Table (5): Biochemical features of isolated *A. sobria*.

Biochemical features			Reaction	Biochemical features			Reaction
2	APPA	-		3	ADO	-	
4	PyrA	-		5	IARL	-	
7	dCEL	-		9	BGAL	+	
10	H ₂ S	-		11	BNAG	+	
12	AGLTp	-		13	dGLU	+	
14	GGT	-		15	OFF	+	
17	BGLU	-		18	dMAL	+	
19	dMAN	+		02	dMINE	+	
01	BXYL	-		00	BAlap	-	
03	ProA	+		26	LIP	-	
07	PLE	-		09	TyrA	+	
31	URE	-		30	dSOR	-	
33	SAC	+		34	dTAG	-	
35	dTRE	+		36	CIT	+	
37	MNT	-		39	5KG	-	
42	ILATk	-		41	AGLU	-	

40	SUCT	+	43	NAGA	+
44	AGAL	-	45	PHOS	-
46	GlyA	-	47	ODC	-
48	LDC	-	53	IHISa	-
56	CMT	+	57	BGUR	-
58	O129R	-	59	GGAA	+
61	IMLTa	-	60	ELLM	+
64	ILAT	-			

+: Positive, -: Negative

Table (6): Biochemical features of isolated *A. veronii*.

Biochemical features	Reaction	Biochemical features	Reaction
2 APPA	-	3 ADO	-
4 PyrA	-	5 IARL	-
7 dCEL	+	9 BGAL	+
10 H ₂ S	-	11 BNAG	+
12 AGLTp	-	13 dGLU	+
14 GGT	-	15 OFF	+
17 BGLU	+	18 dMAL	+
19 dMAN	+	02 dMINE	+
01 BXYL	-	00 BA _{lap}	-
03 ProA	+	26 LIP	-
07 PLE	-	09 TyrA	+
31 URE	-	30 dSOR	-
33 SAC	+	34 dTAG	-
35 dTRE	+	36 CIT	+
37 MNT	-	39 5KG	-
42 ILATk	-	41 AGLU	-
40 SUCT	+	43 NAGA	-
44 AGAL	+	45 PHOS	-
46 GlyA	-	47 ODC	+
48 LDC	-	53 IHISa	-
56 CMT	+	57 BGUR	-
58 O129R	-	59 GGAA	+
61 IMLTa	-	60 ELLM	+
64 ILATa	-		

+: Positive, -: Negative

Table (7): Antibiotic susceptibility of *A. hydrophila*.

Antimicrobial	MIC	Interpretation	Antimicrobial	MIC	Interpretation
Piperacillin/ Tazobactam	<= 4	S	Amikacin	<= 2	S
Cefazolin	>= 64	R	Gentamicin	<= 1	S
Cefoxitin	32	R	Ciprofloxacin	1	S
Ceftazidime	<= 1	S	Levofloxacin	1	S
Cefepime	<= 1	S	Tigecycline	<= 0.5	S
Imipenem	4	S	Trimethoprim/ Sulfamethoxazole	<= 20	S

MIC: Minimum Inhibitory Concentration ($\mu\text{g.ml}^{-1}$), S: Sensitive, R: Resistant.

Table (8): Antibiotic susceptibility of *A. sobria*.

Antimicrobial	MIC	Interpretation	Antimicrobial	MIC	Interpretation
Piperacillin/ Tazobactam	16	S	Amikacin	≤ 2	S
Cefazolin	≤ 4	S	Gentamicin	≤ 1	S
Cefoxitin	≤ 4	S	Ciprofloxacin	≤ 0.25	S
Ceftazidime	≤ 1	S	Levofloxacin	0.5	S
Cefepime	≤ 1	S	Tigecycline	≤ 0.5	S
Imipenem	8	I	Trimethoprim/ Sulfamethoxazole	≤ 20	S

MIC: Minimum Inhibitory Concentration ($\mu\text{g.ml}^{-1}$), S: Sensitive, R: Resistant, I: Intermediate.

Table (9): Antibiotic susceptibility of *A. veronii*.

Antimicrobial	MIC	Interpretation	Antimicrobial	MIC	Interpretation
Piperacillin/ Tazobactam	20	S	Amikacin	≤ 2	S
Cefazolin	≤ 4	S	Gentamicin	≤ 1	S
Cefoxitin	≤ 4	S	Ciprofloxacin	≤ 0.5	S
Ceftazidime	≤ 1	S	Levofloxacin	0.5	S
Cefepime	≤ 1	S	Tigecycline	≤ 0.5	S
Imipenem	8	I	Trimethoprim/ Sulfamethoxazole	≤ 20	S

MIC: Minimum Inhibitory Concentration ($\mu\text{g.ml}^{-1}$), S: Sensitive, R: Resistant, I: Intermediate.

Recently, there are many processes for treating *Aeromonas* spp. (primarily *A. hydrophila*) that infected in cultivated fish, one of them is antimicrobial drugs, it is the common antibiotic for controlling the aquatic bacteria, so antimicrobial sensitivity experiments are crucial for an efficient therapy (Skwor *et al.*, 2014).

High allergy of drug examination in various organisms is feasible technique that can clearly decrease antibiotic dosage (Skwor *et al.*, 2014; Li *et al.*, 2019). In the current study, the Motile *Aeromonas* Septicemia was intensely susceptible to Tigecycline, Ciprofloxacin and Levofloxacin. Therefore, it suggests may be that these drugs are convenient to treat Motile *Aeromonas* Septicemia. On the other hand, there are some risks for using antibiotics, firstly, the medicine can penetrate the

biological membrane and tissues, for example, it can pass through blood vessels in the brain, secondly, using antibiotics for a long time causes the resistance of bacteria, drug accumulation in fish tissues and environmental problems (El-Bouhy *et al.*, 2011; Li *et al.*, 2019).

About 90% of *A. hydrophila*, were susceptible to Ceftazidime, Moxalactam, Aztreonam, Cefepime, Amikacin, Fluoroquinolones and Imipenem, but it was more resistant to Trimethoprim-Sulfamethoxazole, Tetracycline, Aminoglycosides and Cephalosporins (Chopra, & Roberts, 2001; Soltan *et al.*, 2016). Guz & Kozinska (2004) found that about 21 isolates of *A. sobria* and *A. hydrophila* which isolated from common carp were 100% were sensitive to Oxolinic acid, Trimethoprim-Sulphamides,

Cloramphenicol, Flumequine, Norfloxacin, Perfloxacin, lincomycin and resistant to Penicillin.

Nawaz *et al.* (2006); Adanir & Turutoglu (2007) and Hassan *et al.* (2017) confirmed that the resistance to Oxytetracycline, Penicillin, Amoxicillin and other antibiotics may be attributed to genes which recently discovered as responsible for this antibiotic resistance in genus *Aeromonas*.

Conclusion

Aeromonas spp. may cause hemorrhages with ulcers in common carp. Extensive and uncontrolled use of antimicrobial drugs may generate an multiple resistance for antimicrobial. So, the test of antimicrobials susceptibility in genus *Aeromonas* infection and other bacterial infections must be under intensive in cultivated fish. The water should be orderly changed and stress and overcrowding in the fish population should be avoided to prevent the infection.

Acknowledgements

Sincere thanks are due to Dr. Abdul Kareem S. Al-Yassari of the College of Veterinary Medicine, University of Al-Qasim Green, Iraq, for to help with some bacterial analyses.

Conflict of interest: The authors declare that they have no conflict of interest.

Ethical approval: all applicable national and international guidelines for the care and use of animals were followed.

References

Adanir, D.O.R. & Turutoglu, H. (2007). Isolation and antibiotic susceptibility of

Aeromonas hydrophila in a carp (*Cyprinus carpio*) hatchery farm. Bull. Vet. Inst. Pulawy., 51: 361-364.

guilera, A.M.G.; Hernande, C.C.; Zuniga, G.; Figueras, M.J. & Castro, E.G. (2005). *Aeromonas hydrophila* clinical and environmental ecotypes as revealed by genetic diversity and virulence genes. FEMS Microbiol. Lett., 242(2): 231-240.

Albert, V. & Ransangan, J. (2013). Effect of water temperature on susceptibility of culture marine fish species to vibriosis. Inter. J. Res. Pure Appl. Microbiol., 3: 48-52.

Al-Niaeem, K.S.; Shnawa, I.M.S. & Al-Sa'adi, B.A. (2015). The immunological response for spleen as biomarker against of *Aeromonas hydrophila* bacterial in *Cyprinus carpio* fish. Basrah J. Vet. Res., 15(2): 306-315.

Al-Tae, A.M..R; Khamees, N.R. & Al-Shammari, N.A.H. (2017). Vibrio species isolated from farmed fish in Basra City in Iraq. J. Aquac. Res. Dev., 8: 472. DOI: 10.4172/2155-9546.1000472.

Aoki T. (1999). Motile aeromonads (*Aeromonas hydrophila*). Pp: 427-453. In: Fish diseases and disorders. Woo PTK, Bruno D.W. (ed.) CABI Publishing, Wallingford: 944pp.

Aravenaromán, M.; Inglis, T.J.; Riley, T.V. & Chang, B.J. (2014). Distribution of 13 virulence genes among clinical and environmental *Aeromonas* spp in Western Australia. European J. Clin. Microbiol. Inf. Dis., 33(11): 1889-1895.

- Austin, B. & Austin, D.A. (2012). Aeromonadaceae representatives (motile aeromonads): 119-146. In: Austin, B. & Austin, D.A. (Eds.), Bacterial fish pathogens, diseases of farmed and wild fish. 5th ed. Springer, New York, London: 652pp.
- Buján, N.; Romalde, J.L. & Figueras, M.J. (2010). Comparison of phenotypical and genetic identification of *Aeromonas* strains isolated from diseased fish. Syst. Appl. Microbiol., 33: 149-153.
- Beazhidalgo, R. & Figueras, M.J. (2013). *Aeromonas* spp whole genomes and virulence factors implicated in fish disease. J. Fish Dis., 36(4): 371-388.
- Bondad-Reantaso, M.G.; Subasinghe, R.P.; Arthur, J.R.; Ogawa, K.; Chinabut, S.; Adlard, R.; Tan, Z. & Shariff, M. (2005). Disease and health management in Asian aquaculture. Vet. Parasitol., 132(3-4): 249-272.
- Chopra, I. & Roberts, M. (2001). Tetracycline antibiotics: mode of action, applications, molecular biology, and epidemiology of bacterial resistance. Microbiol. Mol. Biol. Rev., 65: 232-260.
- Cipriano, R.C. (2001). *Aeromonas hydrophila* and motile Aeromonad septicemias of fish. Fish Dis Leaflet 68, United States Department of the Interior fish and wildlife service, Div. Fish. Res. Washington, D.C. 25pp.
- Dixon, B.A. & Issvoran G. (1993). Antibacterial drug resistance in *Aeromonas* spp. isolated from domestic goldfish and koi from California. J. World Aquac. Soc., 24: 102-104.
- El-Bouhy, Z.M.; El-Nobi, G.; Reda, R.M. & Ali, S. (2015). Prevalence of septicemia and red mouth disease caused by *Aeromonas sobria* at Sahl El-Housinia fish farm. Zagazig Vet. J., 43(3): 53-63
- Gao, Z.L.; Ji, X.; Guo, X.J.; Chen, P.; Liu, Y.J.; Liu, J. & Sun, Y. (2016). Establishment and application of a duplex real time fluorescence quantitative PCR assay for detection of *Aeromonas hydrophila*. Chinese J. Zoonoses, 32(12): 1126-1130.
- Guz, L. & Kozinska, A. (2004). Antibiotic susceptibility of *Aeromonas hydrophila* and *A. sobria* isolated from farmed carp (*Cyprinus carpio* L.). Bull. Vet. Inst. Pulawy, 48: 391-395.
- Hassan, A.; Noureldin, E.A.; Mahmoud, M. & Fita, N.A. (2017). Molecular identification and epizootiology of *Aeromonas veronii* infection among farmed *Oreochromis niloticus* in Eastern Province, KSA. Egyptian J. Aqu. Res., 43: 161-167.
- Janda, J.M. & Abbott, S.L. (2010). The genus *Aeromonas*: taxonomy, pathogenicity and infection. Clin. Microbiol. Rev., 23: 35-73.
- Ko, W.C.; Wu, H.M.; Chang, T.C.; Yan, J.J. & Wu, J.J. (1998). Inducible β -lactam resistance in *Aeromonas hydrophila*: therapeutic challenge for antimicrobial therapy. J. Clin. Microbiol., 36: 3188-3189.
- Li, F.; Wu, D. & Gu, H.R. (2019). *Aeromonas hydrophila* and *Aeromonas veronii* cause motile *Aeromonas* septicemia in the cultured Chinese sucker, *Myxocyprinus asiaticus*. Aqu.

- Res., 1-12. [https:// doi.org/ 10.1111/are.14028](https://doi.org/10.1111/are.14028)
- Nawaz, M.; Sung, K.; Khan, S.A., Khan, A.A. & Steele, R. (2006). Biochemical and molecular characterization of tetracycline-resistant *Aeromonas veronii* isolates from catfish. Appl. Environ. Microbiol., 72(10): 6461-6466.
- Pachanawan, A.; Phumkhachorn, P. & Rattanachaikunsopon, P. (2008). Potential of *Psidium guajava* supplemented fish diets in controlling *Aeromonas hydrophila* infection in tilapia (*Oreochromis niloticus*). J. Biosci. Bioeng., 106(5): 419-424.
- Rijnsdorp, A.D.; Peck, M.A.; Engelhard, G.H., Mollmann, C. & Pinnegar, J.K. (2009). Resolving the effect of climate change on fish populations. ICES J. Mar. Sci., 66: 1570-1583.
- Skwor, T.; Shinko, J.; Augustyniak, A.; Gee, C. & Andraso, G. (2014). *Aeromonas hydrophila* and *Aeromonas veronii* predominate among potentially pathogenic ciprofloxacin and tetracycline resistant *Aeromonas* isolates from Lake Erie. Appl. Environmen. Microbiol., 80(3): 841-848.
- Soltan, D.M.M.; Yazdi, M.K.S & Avadisians, S. (2012). Study of prevalence and antibiotic resistance in *Aeromonas* spp. isolated from minced meat and chicken samples in Iran. African J. Microbiol. Res., 6(7): 1568-157.