

Evaluation of Mustard oil and temperature in the toxicity reduction of *Aspergillus flavus* in Walnut fruits

Alaa Kaiser Abdul Khaleq Abboud and Sadeq Mohammed Ali

Faculty of Agriculture / University of Kufa / Republic of Iraq

Corresponding author: alaaq.aboud@student.uokufa.iq

DOI: <https://doi.org/10.36077/kjas/2024/v16i1.10507>

Received date: 13/11/2022

Accepted date: 28/12/2022

Abstract

The study aimed to isolate and identify fungi that produce mycotoxins from walnut fruits and the possibility of limiting their presence in the fruits and reducing their toxic effects. The molecular diagnosis in this study of the fungus *Aspergillus flavus* was carried out using the internal transcribed spacer (ITS) genetic marker. As for the results of testing the ability of some *A. flavus* isolates to produce aflatoxins B₁ including the ammonia test, the results of this test showed the ability of four out (of six) isolates of *A. flavus* on the production of aflatoxin B₁ by changing the color of the base of the coconut medium on which the fungus isolates are grown, with a percentage of 66.66 %. The results of the chemical analysis using TLC technology for *Aspergillus flavus* showed the ability of three isolates to produce aflatoxin B₁ out of six isolates of *A. flavus* with a percentage of 50 %. As for the most important results of the effectiveness of mustard oil in inhibiting the growth of the fungus *A. flavus* in walnut fruits, the highest percentage of inhibition was at 15% concentration, which amounted to 57.33%. The most important results of the storage experiment after three months of storage, included the test of the effect of mustard oil and temperature on the growth and density of *A. flavus*. The test was the effect of mustard oil on the reduction ratio of aflatoxin B₁ as it reached the highest percentage of toxin reduction when adding mustard oil only, which amounted to 98.213%, compared to other treatments, the test was the effect of mustard oil and temperature on the amount of aflatoxin B₁ toxin, which gave the highest rate when adding *A. flavus* to walnuts, which amounted to 13.425 micrograms/ g,. At a temperature of 5 °C, it gave the lowest rate of aflatoxin B₁ toxin, which was 0.880 µg.gm⁻¹. The highest rate was 8.620 µg.gm⁻¹, at 25°C.

Keywords: mycotoxins, walnut fruits, *Aspergillus flavus*.



Introduction

The walnut (*Juglans nigra*) is a northern hemisphere fruit tree with a temperate and subtropical climate (30) and is native to Europe and Asia from the Balkans to southwestern China (15). It is widely cultivated all over the world because of its nutritional value (1). Walnuts are high in mono-saturated fatty

acids, vitamins, proteins, and carbohydrates, and as well as it is rich in omega-3. People have been aware of this fruit since the beginning of time. Because its botanical structure is similar to and nutritious to the human brain (29), it is associated with a lower risk of vascular and heart injury (21). During storage with fungal species, including *Asperillus spp*, *Fusarium spp*, and *Pancillium spp*, they produce a wide range of toxic receptors (27). They produce antioxidants against these toxins in addition to their resistance to high temperatures. (34) and many methods have been

used to control fungi in the store, including the use of chemicals such as sterilization using carbon dioxide and others (23, 37). The use of chemicals is not without negatives as a result of environmental pollution and the resistance shown by the pest towards these materials. Chemicals and damage to human and animal health as a result of the accumulation of these substances in the food chain (33), where plant extracts were used because they have antagonistic activity for many fungi and have preferred characteristics such as their rapid decomposition and lack of impact on human health (17). The use of some essential oils derived from plants such as mustard oil because it has anti-fungal, anti-bacterial, anti-microbial, and food pathogen activity (11 and 8). Therefore, the current study aimed to investigate the fungi contaminating

walnut fruits and test the toxins production and study the possible measurements in reducing infection and combined toxicity. It also aimed to confirm the

diagnosis of fungi using the Polymerase chain reaction (PCR) technique and testing the extent of

contamination of walnut fruits with fungi and testing the extent of susceptibility to fungi.

Materials and Methods

Nut sample collection:

Nut samples were obtained from local markets (American, Iranian and Chinese) for the purpose of obtaining isolates of toxin-producing fungi and then transferred to the Laboratory Plant Diseases at the College of Agriculture - University of Kufa to isolate and identify the fungi accompanying those samples.

Isolation and diagnosis of fungi infecting walnuts

Nut samples were taken and isolated from them by dilution method and direct cultivation on PDA medium, where the fruits were cut into small pieces (1 cm) and were superficially sterilized with sodium hypochlorite solution 2% concentration for two minutes after which those pieces were washed with sterile distilled water and then placed on filter papers to get rid of free water. Then the fruit pieces were planted in plastic dishes (9 cm in diameter) containing the medium of PDA by placing four pieces at a distance of 3 cm from the edge of the dish and a fifth piece in the middle of the dish. (20), and after the end of the incubation period, the growing fungi isolates were purified and then identified based on the taxonomic characteristics mentioned by Pitt and Hocking (24).

The percentages of appearance and frequency of fungi were calculated according the following two equations:

$$\text{Percentage of appearance (\%)} = \frac{\text{The number of samples in which the sex or gender appeared}}{\text{Total number of samples}} \times 100$$

$$\text{Percentage Frequency (\%)} = \frac{\text{The number of isolates of one species}}{\text{The total number of isolates of all fungi}} \times 100$$

Molecular diagnosis of isolates of the fungus *A. flavus* The isolates of the fungus *A. flavus* on liquid PD medium by placing three tablets of five mm in diameter, one week old, in a flask of 250 mL volume, at a rate of 100 m. L⁻¹. Thereafter the extract was filtered through Whatman No. 4. Then the filtrate was then placed in a separator funnel with a capacity of 250 mL, then 30 mL of chloroform was added to it, then the funnel was shaken gently for 30 seconds while expelling the accumulated gases whenever needed (at least twice) and then left on the holder for one minute in order to The two layers separate, I neglected the upper layer and took the lower layer and repeated the process for three times. Then the filtrate was taken and placed in a clean, sterile flask and placed in the electric oven at a temperature of 40°C until it dried, then dissolved in one ml of chloroform. The presence of aflatoxin, B1 was detected using thin-sheet chromatography (TLC) technology with dimensions of 20-20 cm. The plates were activated in the electric oven was heated at 105 °C for one hour before use (7).

Aflatoxin B1 producing by *A. flavus* isolates

Detection of Aflatoxins by ammonia treatment method:

The ability of *A. flavus* isolates isolated from the neglected the upper layer and took the lower layer and tested nuts to produce aflatoxin B1 was carried out repeated the process for three times. Then the filtrate using (10) method, using coconut medium. In dishes was taken and placed in a clean, sterile flask and each with a diameter of 9 cm, then 3 dishes were placed in the electric oven at a temperature of 40°C inoculated with tablets of fungi for each isolate, by until it dried, then dissolved in one ml of chloroform. placing a disk with a diameter of 5 mm from the The presence of aflatoxin, B1 was detected using thin-medium of the PDA grown on The Fungi, at the age of sheet chromatography (TLC) technology with one week, in the center of the dish. A week later, the dimensions of 20-20 cm. The plates were activated in the ability of the mushroom isolates to produce aflatoxins the electric oven was heated at 105 °C for one hour was revealed using a 20% ammonia solution by before use (7).

placing filter papers saturated with ammonia solution A light straight line was made on the TLC plate at a in the cover of the dish containing the isolate of the distance of 1.5 cm from the base of the plate, and 15 fungi growing in the middle of the coconut. The color μl was taken by capillary tube of the standard toxin of the colony bases from transparent to pink or red AFB1 and placed on the line at a distance of two cm indicates that this isolate is capable of producing from the left edge of the plate and at a distance of 2 aflatoxins. cm from the spot of the standard poison. *A. flavus* at

Detection of aflatoxins using Thin Layer the same distance and in an amount equal to the Chromatography (TLC): amount of the standard poison, then the spots were left to dry and then placed in a separation tank containing



a separation system consisting of a mixture of 1- Preparation of *A. flavus* inoculum: *A. flavus* chloroform and methanol at a ratio of 2:98 v/v and inoculum was prepared by growing the fungus on monitored until the solution reached a distance of PDA medium for a week under temperature of $30 \pm$ approximately two cm from the upper end For the 2°C . Then the fungal spores were harvested by adding plate, the plates were taken out and dried under 10 mL of sterile distilled water to each of the plates on laboratory conditions for 5 minutes and then which the fungus was grown, and then passed A examined under ultraviolet light at a wavelength of sterile glass rod on the surface of the colonies to 365 nm and the presence of aflatoxin B1 was detected facilitate the process of separating spores from the by matching the migration factor R_f and fluorescence conidia carriers. Then, planktonic spores were color of the standard poison with the color and collected for each fungus separately and their numbers migration factor of samples of extracts of *A. flavus* were calculated using a hemocytometer slide. The isolates (31) fungal spore concentrations were adjusted at (106

Testing the efficacy of Indian mustard oil extract on the growth of *A. flavus* on PDA medium:

Indian mustard oil was obtained from one of the

local markets, so it was added in proportions 5, 10 and 1-Distilled water only 200 gm of nuts

15 mL / 100 mL of PDA medium and poured into the 2-10 mL Indian mustard oil 200 gm walnuts

dishes. A fungus from the tested fungi. Three plates of 3-The spores of *A. flavus* are stuck in only 200 gm of PDA medium (control treatment) were inoculated nuts

with PDA discs only on which the fungus *A. flavus* 4-Suspended *A. flavus* spores + 10 mL of Indian was grown. The dishes were incubated at $25 \pm 2^\circ\text{C}$ mustard oil 200 gm of nuts

for a week. After that, the diameters of *A. flavus*

colonies were calculated, and according to the Each treatment was stored in three bags containing each bag (200 gm) for a period of (30, 60, and 90) days at temperatures (5, 15, 25, and 35) $^\circ\text{C}$, after which the contamination percentage was calculated in each of the implemented treatments. Taking 10 g of each of the repeaters of one treatment and placing it in

extracted according to the Abbott equation reported by Shaaban and Al-Mallah (35) where:

$$\text{Inhibition}\% = \frac{R_1 - R_2}{R_1} \times 100$$

R_1 = maximum radial growth of a fungus an electric mixer with the addition of 100 mL of sterile distilled water, then it was mixed for 5 minutes,

R_2 = maximum radial growth of the studied then a series of dilutions was made until 10^{-5} , then one mL of the last dilution was grown on the PDA medium. And three replications for each treatment,

Evaluation of the efficiency of Indian mustard oil and storage temperature in protecting walnut fruits from infection with the fungus *A. flavus*;

and thus all treatments were dealt with, after which the dishes were incubated at a temperature of $25 \pm 2^\circ\text{C}$. After that, the number of colonies in each dish was

This experiment was carried out to find out there recorded after 3-4 days. Then, the number of active effectiveness of both mustard oil and temperature on units of the fungus was calculated, according to the protecting walnut fruits from infection with *A. flavus*. coefficients, according to the following equation:

The number of active units = the average number of layer, and after three minutes, the upper and lower spores per replicate x the reciprocal of the dilution. layers were separated, keeping the lower layer. The

The concentrations of aflatoxin B1 were measured by a spectrophotometer, after extracting the mycotoxins from the samples of the implemented treatments, the aflatoxin B1 toxin was extracted by taking 30 gm of walnuts from each treatment and then transferring it to an electric mixer containing 100 mL of chloroform and then mixing the mixture for 10 Minutes after that, the mixture was filtered by filter paper, then the filtrate was taken and placed in a clean and sterile flask and placed in an electric oven at a temperature of 40 ° C until dryness, then it was dissolved in 5 mL of chloroform and then the concentration of the poison was estimated. As for the ochratoxin a poison, it was extracted by taking 30 gm of chloroform. The atmosphere from each treatment and Then to an electric mixer containing 100 mL of the extraction solution consisting of acetonitrile-water (90:10) mL, then mix the mixture for 10 minutes, then filter The extract through what man No. 4. Then the filtrate was placed in a separator funnel with a capacity of 250 mL, then 25 mL of hexane was added to it for the purpose of getting rid of fat (Defatting), Then The funnel was shaken gently for 30 seconds while expelling the accumulated gases whenever needed (at least twice). Then it was left on the holder for one minute in order for the two layers to separate, I neglected the top layer and took the bottom layer, and repeated the process Three times to ensure the removal of fat. Then, 25 mL of distilled water, 8 mL of saturated sodium bicarbonate solution (NaHCO₃), and 25 mL of chloroform were added to the lower process was repeated twice. Then the extract was transferred to a separating funnel with a capacity of 100 mL, 15 mL of 1-N hydrochloric acid, 20 mL of chloroform, and 20 mL of chloroform were added to it. Shake well and leave the funnel on the holder for one minute. The lower layer was taken and to the upper layer, 20 mL of chloroform was added and extracted again. The two lower layers were collected and passed through filter paper containing a layer of anhydrous sodium sulfate (Na₂SO₄) in order to get rid of the remaining water. Then the filtrate was taken and placed in an electric oven at a temperature of 40 °C until dryness, after which it was dissolved in 5 mL of chloroform (9) and then the poison concentration was estimated by a Spectrophotometer. This method relied on the spectrophotometric estimation, which depends by its nature, on the property of the compound to absorb light in the ultraviolet or infrared wavelengths, where there is a direct proportion between the absorption period and the concentration of the poison. It is possible by drawing a standard curve between the light absorption and the concentration of the poison and extracting the concentration value corresponding to the reading given by the unknown sample. Then the readings were compared with Standard poison concentrations (Supplements 1 and 2), where the wavelengths were 365 and 360 nm for aflatoxin B1 and ochratoxin A, respectively. The reduction ratios were calculated Using the following equation:

$$\text{Reduction ratio} = \frac{\text{Concentration of control treatment} - \text{concentration of the model}}{\text{Control treatment concentration}} \times 100$$

Statistical analysis:

All experiments were carried out in a completely randomized design, C.R.D (Complete Random Design) as one-factor experiments, and the averages were compared according to L.S.D (Less significant differences) method and under a probability level of 0.05 (4).

Results and Discussion

Isolation and identification of fungi accompanying the nut:

The results of isolating fungi from walnuts showed the appearance of five species of fungi. The most common were *A. tubingensis*, *A. flavus*, *Fusarium* sp, *Penicillium* sp, and *Trichoderma* sp. Its frequency rates reached 55.17, 20.68, 13.79, 6.89 and 3.44%, respectively. (Table 1) While the incidence of (100, 66.66, 33.33, 33.33 and 16.66) %, respectively, explains the reason for the dominance of *Aspergillus* sp. The ability to withstand drought and high humidity (2), and withstand high temperatures up to 50 °C (12).

Table 1. Percentages of the frequency and appearance of fungi isolated from walnut

fungus type	(%) frequency	(%) apperance
<i>A. tubingensis</i>	55.17	100
<i>A. flavus</i>	20.68	66.66
<i>Fusarium</i>	13.79	33.33
<i>Penicillium</i>	6.89	33.33
<i>Trichoderma</i>	3.44	16.66

Molecular diagnosis of *A. flavus* isolates

The results of the molecular diagnosis of the pathogenic fungus *Aspergillus flavus* indicated that this isolate was established in the Neighbor-joining

tree and in the National Center for Biotechnology Information (NCBI) under the entry (Accession number ON394601.1.) Based on the sequences of its nitrogenous bases for the ITS-rDNA region as well as the sequences of global strains of the same pathogenic fungus were obtained from Gen Bank Data



Repository. The genetic distances were calculated strain (16). The biological basis of the ammonia test using the neighbor-joining method.

Testing the ability of some isolates for *A. flavus* to produce aflatoxins B1:

Ammonia test:

The results of this test showed the ability of 4 out of 6 isolates of *A. flavus* to produce aflatoxin B1 by changing the color of the base of the coconut medium on which the fungus isolates were grown, with a percentage of 66.66 %. The isolates that produced the most aflatoxins were Af4, while isolates Af2 and Af5 showed a medium ability to produce aflatoxins, and the rest of the isolates were weak in their production of aflatoxins (Table 2).

The fungal strains vary in their ability to secrete produce aflatoxin quantitatively and qualitatively may aflatoxin, some of them may excrete aflatoxin toxins, be due to the different genetic content of the strains, while there are other strains that secrete more than one and this explains the gradient in the red color. (26).

type of mycotoxin depending on the type of fungal

Table 2. Testing the ability of some isolates of *A. flavus* to produce aflatoxin B1 in medium coconut (CEA)

fungul isolation	The ability to produce aflatoxin B1
Af1	+
Af2	++
Af3	-
Af4	+++
Af5	++
Af6	-

(+): the color of the middle base changed to pink, (-): the color of the middle base did not change.

Detection using thin sheet chromatography (TLC) The results of this test showed the ability of some isolates of *A. flavus* to produce aflatoxin B1, whereas the test showed the ability of 3 isolates to produce

aflatoxin B1 out of 6 isolates of *A. flavus* at a produce aflatoxin B1. The difference in the ability of percentage of 50%, as in (Table 3). The fungus isolates to produce aflatoxins B1 may be attributed to isolates varied in their toxin production, and isolate the presence of genetic differences between the fungal AF4 was the most toxin-producing isolate based on isolates (18). It is noted that the percentage of isolates the intensity of its fluorescence. And Af5 and Af2 are reproducing aflatoxin B1, which were detected by this the least productive of aflatoxin production. This technique, is less than the number of isolates results initially agree with what one study indicated producing aflatoxin B1 using the ammonia reaction about the ability of 75% of *A. flavus* isolates to technique. Accordingly, the thin plate produce aflatoxin B1. And an approach to what was chromatography method is the most accurate method mentioned (14). This indicated that 38.88% of *A. flavus* isolates isolated from walnuts are able to general, including aflatoxin B1.

Table 3. Testing the susceptibility of a number of *A. flavus* isolates to produce aflatoxin B1 isolated from walnuts by thin plate chromatography (TLC) method

fungul isolation	The ability to produce aflatoxin B1
Af1	-
Af2	+
Af3	-
Af4	+++
Af5	++
Af6	-

(+) aflatoxin B1-producing isolate (-) non-aflatoxin B1-producing isolate

Evaluation of the effectiveness of different highest percentage of inhibition was at the concentrations of mustard oil in inhibiting the Qatar concentration 15%, which amounted to 57.33%, and growth of the fungus *A. flavus*: the inhibition percentage in concentrations 5, 10 and

The results of the experiment showed the treatment (table 4) This is similar to what previous studies of nuts contaminated with the fungus *A. flavus* with indicated to the effectiveness of many medicinal mustard oil helped it to inhibit the growth of the plants and their volatile oils in inhibiting toxin-producing fungi (28). The fungus *A. flavus*, as the percentage of inhibition increased by increasing the concentration and the

Table 4. Effect of different concentrations of mustard oil on the growth of *A. Flavus* grown on PDA

Transactions	retarding percentage%	Colonies diameter average (cm)
PDA. Comparison	0.00	7.5
5	48.00	3.9
10	52.00	3.6
15	57.33	3.2
L.S.D.(0.05)	1.631	0.1883




Figure 1. shows the evaluation of the effectiveness of different concentrations of mustard oil in inhibiting the growth of the fungus *A. concentration* of 5% and number (3) *flavus*, which are three different represents a concentration of 10% and number concentrations (5%, 10% and 15%). No. (1) (4) represents a concentration of 15%. represents the comparison, which is *A. Effect of mustard oil and temperature on the growth and density of A. flavus* after three months of storage.

The results of this test showed significant differences in the presence of erucic acid in mustard seed oil, differences in the growth rate of the number of which plays a role in phytoremediation and reproductive units of *A. flavus* spores, as it gave the antioxidants (36). The table also indicates that there is the highest growth rate when *A. flavus* was added to the treatments, where there are significant differences in the effect of temperature on the number of reproductive units of *A. flavus*, so compared to the control treatment (0.25×10^{-5} spores/gm). The storage treatment at a temperature of 35°C gave gm^{-1}). In Table (5), while the mustard oil reduced the lowest rate in the number of reproductive units of numbers of microbial units of the fungus, where the fungus, which amounted to 0.575×10^{-5} spores. treatment of *A. flavus* + mustard oil gave $0.575 \times 10^{-5} \text{gm}^{-1}$ compared to the other treatments (5, 15, 25). mL, spores / gm^{-1} , compared with the treatment of adding which gave ($1.080, 1.140, 1.560 \times 10^{-5}$ spores. g), fungus only, which amounted to 3.200×10^{-5} spores. respectively. The highest rate in the number of Gm^{-1} , while the oil treatment was given Mustard only reproductive units of the fungus was (1.560×10^{-5} had the lowest growth rate in the number of spores. gm^{-1}) at 25°C compared with other reproductive units, which amounted to (0.000×10^{-5} treatments, so crop losses were recorded from 20% to spores. gm^{-1}) compared to the control treatment (0.25550% in developing countries due to inappropriate $\times 10^{-5}$ spores. gm^{-1}). Whereas, the addition of distilled storage practices (22). water gave only 1.425×10^{-5} spores / g, and the reason

Table 5. Effect of mustard oil and temperature on the growth and density of *A. flavus* after three months

Transactions	Number of reproductive units of spores (spore/gm) x 10 ⁵				The average
	Temperatures				
	°5	°15	°25	°35	
Comparison	0.0	0.4	0.2	0.3	0.225
distilled water	0.4	1.6	2.6	1.1	1.425
Mustard oil	0.0	0.0	0.0	0.0	0.000
<i>A. flavus</i>	4.2	2.9	4.6	1.1	3.200
<i>A. flavus</i> + Mustard oil	0.8	0.8	0.4	0.3	0.575
The average	1.080	1.140	1.560	0.575	1.080

L.S.D. 0.05 coefficients = 0.0715, temperatures = 0.0639, coefficients + temperatures = 0.1429

Effect of mustard oil and temperature in reducing the oil, *A. flavus*, mustard oil + *A. flavus*) (98.213, 0.000, toxicity of aflatoxin B1 produced by *A. flavus* after 71.188) % respectively, as indicated in Table (6), three months.

This test showed the effect of mustard oil on the reduction ratio of aflatoxin B1 when it gave (mustard oil), where the highest percentage of toxin reduction when adding mustard oil only, which amounted to 98.213% compared to other treatments, and the lowest percentage of toxin reduction in the treatment of

adding *A. flavus* only, which amounted to (0.0000%) respectively, and the highest rate was 42.953% at a compared to other treatments because it has many The temperature of 25 °C compared to other treatments, characteristics that make mustard a valuable botanical due to the conditions that promote the growth of fungi medicine (19). The table also indicates that there are always may not lead to the production of mycotoxins. significant differences in the effect of temperature on In general, it water activity above 0.78, relative the percentage of toxin reduction produced by the humidity between 88% and 95%, and temperature fungus. % compared to the other treatments (5, 15, 25 between 25 and 30 °C are favorable conditions for the °C) which gave (42.855, 41.927, 42.953) %, growth of fungi and production of mycotoxins (32).

Table 6. Effect of mustard oil and temperature in reducing the toxicity of aflatoxin B1 produced by *A. flavus* after three months

Transactions	% reduction ratio				The average
	Temperatures				
	°5	°15	°25	°35	
Comparison	0.0	0.0	0.0	0.0	0.00
Mustard oil	100.0	92.85	100.0	100.0	98.213
<i>A. flavus</i>	0.0	0.0	0.0	0.0	0.0000
<i>A. flavus + Mustard oil</i>	71.42	74.86	71.81	66.66	71.188
The average	42.855	41.927	42.953	41.665	42.350

L.S.D. 0.05 Transactions = 0.5109, temperature = 0.5109, Transactions + temperature = 1.0219

Effect of mustard oil and temperature on the amount of *flavus* only gave (13.425 µg.g⁻¹). The addition of of aflatoxin B1 microgram/gm produced by *A. flavus* mustard oil only had the lowest rate, which amounted after three months of storage: to (0.075 µg/gm) compared to the control treatment

The results of this test showed significant differences in the amount of aflatoxin B1 toxin produced by *A. flavus* fungus, as it gave the highest rate when adding *A. flavus* to walnuts, which amounted to (13.425 µg.g⁻¹) compared to the control treatment, which amounted to (2.708 µg.g⁻¹) As indicated in (table 7) While mustard oil reduced the amount of aflatoxin B1, the treatment of *A. flavus* + mustard oil gave 3.800 µg.g⁻¹ compared with the treatment of adding *A. flavus* only, which amounted to (13.425 µg.g⁻¹), while the treatment of adding *A.* (2.708 µg.g⁻¹). While the treatment of adding distilled water gave only 8.708 µg/gm. This is similar to what (25) said that mustard oil is used as a treatment and antimicrobial and is of plant origin to inhibit fungal and bacterial microbes. The table also indicates that there were significant differences in the effect of temperature on the amount of aflatoxin B1 toxin produced by *A. flavus*, so the storage treatment at a temperature of 5 °C gave the lowest average in the amount of aflatoxin B1 toxin, which was 0.880 µg.g⁻¹ compared to the other treatments (15, 25, 35) m °



which gave (8.307, 8.620, 4.760) $\mu\text{g.g}^{-1}$ respectively. on the biochemical, genetic and physiological And the highest rate was (8.620 $\mu\text{g. g}^{-1}$), at a characteristics of the fungal strains and also on temperature of 25°C compared with the other external factors such as temperature, humidity and treatments. Because the production of mycotoxin specific environments (12). from *Aspergillus*, *Penicillium* and *Fusarium* depends

Table 7. Effect of mustard oil and temperature on the amount of aflatoxin B1 microgram/gm produced by *A. flavus* after three months of storage

Transactions	The amount of aflatoxin B1 µg/gm				The average
	Temperatures				
	°5	°15	°25	°35	
Comparison	0.0	4.2	5.4	1.2	2.708
distilled water	2.6	14.1	9.0	6.6	8.075
Mustard oil	0.0	0.3	0.0	0.0	0.075
<i>A. flavus</i>	1.4	18.3	22.0	12.0	13.425
<i>A. flavus</i> + Mustard oil	0.4	4.6	6.2	4.0	3.800
The average	0.880	8.307	8.620	4.760	5.617

L.S.D. 0.05 coefficients = 0.3742, temperatures = 0.3347, coefficients + temperatures = 0.7484

Conflict of interest

The authors have no conflict of interest.

References

- 1- A.Jahanban-Esfahlan, A.; M. Ostadrahimi; M. Tabibiazar and Amarowicz, R.2019. A Comprehensive Review on the Chemical Constituents and Functional Uses of Walnut (*Juglans* spp.) Husk. Int. J. Mol. Sci., 20(16):3920.DOI: [10.3390/ijms20163920](https://doi.org/10.3390/ijms20163920).
- 2- Abdullah, S. K. and S. M. AL-Bader.1990. On the Thermophilic and Thermotolerant Mycoflora of Iraqi Soils. Sydowia. 42. Iraq. pp.17.
- 3- AL-Adil, K. M.; A. A. N. Basima; A.Y. Sawsan and Kassim, A. D.1977.Contamination by *Aspergillus flavus* group of some foodstuffs in Baghdad. Bull. Biol. Res. Center., 9:107-115.
- 4- Al-Rawi, K. M. and A. Khalafallah.2000.Design and Analysis of Agricultural Experiments. Second Edition. Dar



- Al-Kutub for Printing and Publishing, University of Mosul. Iraq.
- 5- **Al-Waeli, H. W. 2002.** The effect of the volatile oil of yellow peels of grapefruit *Citrus paradisi* and leaves of lemongrass *Cymbopogon citratus* on the growth of the fungus *Aspergillus flavus* and its production of aflatoxin B1. M. Sc. Thesis. Faculty of Education. University of Baghdad. Iraq.
 - 6- **Adekoya, I.; P. Njobeh; A. Obadina; S. Landschoot; K. Audenaert; S. Okoth; M. De Boevre and De Saeger, S. 2019.** Investigation of the metabolic profile and toxigenic variability of fungal species occurring in fermented foods and beverages from Nigeria and South Africa using UPLC-MS/MS. *Toxins*, 11(2):85.
 - 7- **Al-Jumaili, S. A. 1996.** Integrated resistance against infection with the fungus *Aspergillus flavus* and aflatoxin B1 contamination in the yield of field pistachios. Doctoral thesis. Faculty of Agriculture, University of Baghdad. Iraq.
 - 8- **Andini, S.; C. Araya-Cloutier; B. Lay; G. Vreeke; J. Hageman and Vincken, J. P. 2021.** QSAR-based physicochemical properties of isothiocyanate antimicrobials against gram-negative and gram-positive bacteria. *LWT*, 144, 111222.
 - 9- **Balzer, I.; C. Boddanic and Peplujak, S. 1978.** Rapid thin layer chromatographic method for determining Aflatoxin B1, Ochratoxin A, and Zearalenon in Corn. *J. Assoc. of Anal. Chem.*, 61(3): 584-585.
 - 10- **Dianese, J. C. and M. T. Lin. 1976.** A coconut agar medium for rapid detection of aflatoxin production by *Aspergillus spp.* *Phytopathology*, 66:1416-1418.
 - 11- **Drakopoulos, D.; G. Mica; R. Torrijos; A. Marty; A. Kägi; E. Jenny; H. R. Forrer; J. Six and Vogelgsang, S. 2020.** Control of *Fusarium graminearum* in Wheat with Mustard-Based Botanicals: From *in vitro* to in planta. *Front. Microbiol.*, 11:1595.
 - 12- **Gabler, F.M.; M. F. Mansour; J. L. Smilanick and Mackey, B. E. 2004.** Survival of spores of *Rhizopus stolonifer*, *Aspergillus niger*, *Botrytis cinerea* and *Alternaria alternate* after exposure to ethanol solutions at various temperatures. *Journal of Application Microbiology*. 96(6):1354–1360.
 - 13- **Garcia-Diaz, M.; J. Gil-Serna; C. Vázquez; M. N. Botia and Patiño, B. 2020.** A comprehensive study on the occurrence of mycotoxins and their producing fungi during the maize production cycle in Spain. *Microorganisms* 8(1):141. doi:10.3390/microorganisms8010141.
 - 14- **Gherbawy, A. Y.; H. M. Elhariry and Bahobial, A. S. 2012.** Mycobiota and Mycotoxins (Aflatoxins and Ochratoxins)



- Associated with some Saudi Date Palm Fruits. Foodborne Pathogens and Disease, 9 (6):561-567.
- 15- **Khadivi-Khub, A.; A. Ebrahimi; F. Sheibani and Esmaeli, A.**2015.Phenological and pomological characterization of Persian walnut to select promising trees. Euphytica, 205(2):557–567.
 - 16- **Khatoon, S.**2004.Interactions Between Mycotoxins. Romer Labs. Rawalpindi. Pakistan.
 - 17- **Khalaf, J. M.**2012.Testing the effect of bacteria *Pseudomonas fluorescens* and extracts of some plants to control fungi *F. solani* and *R. solani* that cause pepper rot disease. Master Thesis, Musayyib Technical College. Iraq.
 - 18- **Lee, Y. J. and W. M. J. Hagler.**1991.Aflatoxin and Cyclopiazonic acid produced by *Aspergillus flavus* isolated from contaminated. J. Food Sci., 56:871-872.
 - 19- **Malode, S. N. and P. B. Shelke.**2010.Morphological, phenological and anatomical studies in yellow seeded mutant of *Brassica juncea*. Bionano Frontier, 3 (2):172-177.
 - 20- **Mikhail, S. and T. Baydar**1982.Seed Diseases. House of Books for Printing and Publishing. University of Mosul. Iraq.
 - 21- **Mohammadifard, N.; A. Salehi-Abargouei; J. Salas-Salvadó; M. Guasch- Ferré; K. B. Humphries and Sarrafzadegan, N.**2015.The effect of tree nut, peanut, and soy nut consumption on blood pressure: a systematic review and meta-analysis of randomized controlled clinical trials. Am J Clin Nutr., 101(5):966– 982. doi: 10.3945/ajcn.114.091595.
 - 22- **Neme, K. and A. Mohammed.**2017.Mycotoxin occurrence in grains and the role of postharvest management as mitigation strategies. A Review. Food Control, 78, 412–425.
 - 23- **Nilrattanakhum, W.**2003.Control of Aflatoxin Contamination of z Food and Fertilizer technology center (FFTC). Practical Technology Postharvest Available at http://www.agnet.org/library_pt/2003012 . Accessed February.
 - 24- **Pitt, J. I. and A. D. Hocking.**1997. Fungi and Food Spoilage. 2nd Edition. Blackie Academic and Professional. London. England. pp 593.
 - 25- **Reyes-Jurado, F.; T. Cervantes-Rincon; H. Bach; A. Lopez-Malo and Palou, E.**2019. Antimicrobial activity of Mexican oregano (*Lippia berlandieri*), thyme (*Thymus vulgaris*), and mustard (*Brassica nigra*) essential oils in gaseous phase. Ind. Crop Prod., 131:90–95..
 - 26- **Saito, M. and S. Machida.**1999.A rapid identification method for aflatoxin-producing strains of *Aspergillus flavus* and *A. parasiticus* by ammonia vapor. Mycoscience, 40:205-208.



- 27- **Salau, I. A.; K. Shehu and Kasarawa, A. B.**2017.Morphological characterization of mycotoxigenic fungi contaminating groundnut products in Sokoto State. International Journal of Research in Pharmacy and Biosciences. 4(4):1-7.
- 28- **Saratha, V. and S. P. Subramanian.**2010.Evaluation of antifungal activity of Calotropis gigantea Latex extract: an in vitro study. Int. J. Pharm. sc. & Res., 1(9):88-96.
- 29- **Sen, S. M.**2011.Walnut, Cultivation, Nutritional Value, Folklore. 4th Ed. ICC Publication Ankara, Turkey. pp. 220.
- 30- **Simsek, M.; E. Gulsoy; O. Beyhan; A. Osmanoglu and Turgut, Y.**2017.Determination of some botanical, phenological, physical and chemical characteristics of walnut (*Juglans regia* L.) genotypes grown in Turkey. Applied Ecology and Environmental Research, 15(3):1279–1291.
- 31- **Sobolev, V. S, and Dorner, J. W.**2002.Cleanup procedure for determination of aflatoxins in major agricultural commodities by liquid chromatography Journal of AOAC International, 85(3):642-645.
- 32- **Thanushree, M. P.; D. Sailendri; K. S. Yoha, K.S; M.P. Thanushree; D. Sailendri; K. S. Yaho; J. A. Moses and Chinnaswamy, A.**2019.Mycotoxin contamination in food: An exposition on spices. Trends Food Sci Technol 93: 69–80. doi: 10.1016/j.tifs.2019.08.010.
- 33- **Vasilevski, G.**2003.Perspectives of the application of biophysical methods in sustainable agriculture. Bulgarian Journal of Plant Physiology, 29(3):179-186.
- 34- **Wahba, N. M. and N. A. Al-Nasr.**2010.Mycotoxins in milk and dairy products, danger and prevention. Assiut Journal of Modern Studies. Thirty-fourth issue.
- 35- **Shaaban, A. and N. M. Al-Mallah.**1993.Pesticides. House of Books for Printing and Publishing. Mosul University. Iraq. pp.520.
- 36- **Zhu, Y.; E. A. H. Pilon-Smits; F. Zhao; P. N. Williams and Meharg, A. A.**2009.Selenium in higher plants: understanding mechanisms for biofortification and phytoremediation. Trends Plant Science 14(8):436-442.
- 37- **Al-Hamadani, K. M. H. and Sabit, F. A.** 2023. Effect of *Aspergillus flavus* filtrate on different stages of *Callosobruchus maculatus* (Fabricius.) (Chrysomelidae: Coleoptera) on Broad beans. Kufa Journal for Agricultural Sciences, 15(2): 1-10.

