

HYPOTHALAMIC VASCULARIZATION IN THE GOATS AS REVEALED BY VASCULAR CORROSION CAST

A.A.Sawad

Department of Anatomy, College of Veterinary Medicine, University of Basrah, Basrah, Iraq

(Received 14 May 2006, Accepted 29 June 2006)

Keywords: Cold cure, mammillary body, Circle of willis.

ABSTRACT

The heads of ten adult goats of both sexes were used to study the hypothalamic blood supply. It was found that the hypothalamus is supplied by branches of cranial cerebral, cranial communicating arteries, caudal communicating as well as from tiny branches of ophthalmic artery. The blood supply of mammillary body branches from the caudal cerebral, communicating and last branches of basilar artery. The hypothalamus is rather small in comparison to the rest of brain. Its blood supply is extremely rich, feeding from many branches of cranial cerebral arteries that forms the circle of willis. This indicates that the hypothalamus has very high activity.

INTRODUCTION

The hypothalamus is a region of brain that controls an immense number of body functions. It is located on the middle of the base of the brain and encapsulates the ventral portion of the third ventricle. the lower part of the hypothalamus attached with the pituitary stalk by the tuber cinereum (1).

The blood supply of the hypothalamus and pituitary gland are entirely separated while their functions are linked, thus numerous methods have been used to study the blood supply of two regions together so far, the blood supply to the hypothalamus and hypophysis are well as their connection (2). The injection of Indian ink or Berlin blue into blood vessels supplying this parts was performed and they studies macroscopically by cutting in thick or thin slices. The arterial supply and venous drainage of rat preoptic region, nucleus supraopticus and the hypothalamus, was demonstrated by using Indian ink with double perfusion technique (3,4,5).

The capillaries in the median eminence and neural portion of the neurohypophysis stalk, is in contact with termination for nerve fibers of the hypothalamo-hypophyseal tract. These plexus connect to the sinusoid or secondary plexus in the anterior pituitary via a series of portal trunks (6,7). The hypothalamo-hypophyseal vessels run from the median eminence of the hypothalamus along the sides of pars tuberalis to the anterior pole of the pars distalis (8).

MATERIALS AND METHODS

The heads of fourteen adult goats *Caprus hircus* regardless of their sex, were used in this study. A corrosion cast method was applied to the materials. The vessels were washed with 0.9% saline solution that injected via common carotid arteries. The heads were obtained and followed by the injected by a modified, red pigmented, Polymerized Cold Cure Denture base compounds which prepared by solved 20% powder monomethyl-methacrylate in 80% liquid Methyl methacrylate monomer (Dental Supplies LTD, Surrey GU22, England) were used a 25 ml automatic vaccinating syringe (Hauptner-Muto, Germany). (9) They were kept at a room temperature for 24 hours for polymerization. They were corrosion casted in 30% KOH for 24-48 hours, washed with tap water, and photographed (10).

RESULTS

It was found from present study the hypothalamus of the goats were includes the preoptic area, optic tract, optic chiasma, mammillary bodies, tuber cinereum and infundibulum to which the stalk of hypophysis is attached.

The arteries supplying the hypothalamus in goat are all derived from the circle of willis which is formed by the common carotid artery and the most cranial branches of the basilar artery as well as the anastomotic branches (Fig-1). The preoptic region of hypothalamus (Anterior part) was found to be supplied by branches of the anterior cerebral arteries and anterior communicating artery (Fig-2). These arteries course dorsally and laterally to enter the brain and ramify into many branches to supply the anterior part of the hypothalamus. The optic chiasma was supplied by three main branches. The first one is from prechiasmal branches of the ophthalmic arteries, the second branch from superior chiasmal arteries which are branches of the anterior communicating artery. These arteries pass forward over the dorsal surface of the optic chiasma, they join with tiny prechiasmal branches from the ophthalmic arteries. The third branch is from the superior hypophyseal arteries which are branches of the common carotid artery, they also were give small branches to join with prechiasmal branches from ophthalmic arteries (Fig-3).

The mammillary body is highly vascular of the posterior cerebral arteries connecting between the basilar artery and the posterior communicating arteries, the major parts of mammillary nuclei are supplied by posterior mammillary arteries (Fig-4) which have branches from the distal portion of the basilar artery and posterior cerebral arteries.


The blood supply of the tuber region and infundibulum is observed as circumfundibular anastomosis (Fig-5) deriving from three branches from the circle of willis. They are the superior hypophyseal arteries from the common carotid arteries and the tiny branches of ophthalmic arteries.



Fig (1):Hypothalamic vascular cast in goat shows:
(A) Basilar artery , (B)Common carotid artery , (C) Circle of willis



Fig (2):Hypothalamic vascular cast shows:
(A)Anterior communicating artery , (B)Anterior cerebral artery
(C) Carotid retia , (D) External carotidartery , (E) Ophthalmic artery



Fig(3): Schematic representation of blood vessels distributing in the goat shows: (A)Carotid retia , (B) internal maxillary artery , (C) External carotid artery , (D) Basilar artery , (E) Circle of willis (F) Common carotid artery



Fig (4):Micrograph of vascular cast of blood vessels distributing in the Goat mammary body , (A) posterior cerebral artery , (B) Mammary body. X10



Fig(5): Vascular cast of circulatory flow of hypothalamic region in sheep
(A) Ophthalmic artery , (B) Internal maxillary artery

DISCUSSION

The distribution of the major cerebral arteries in goats are similar to those in man (4,11,12) and rat (4,5,13). The goat cerebral arterial circle is formed by the anterior cerebral arteries, posterior communicating arteries, posterior cerebral arteries and common carotid arteries, that are similar to those in rat but different from man that the arterial circle formed by the anterior cerebral, posterior cerebral and posterior communicating arteries. The anterior communicating artery is present in goat as in man (11,12) but is absent in the rat, Opossum and Armadillo (14), in addition, there is corresponding ophthalmic artery in rat as found in goat and man (12) as well as in dog and cat (12,15).

The source of blood supply as well as the distributions of arterial blood in hypothalamus of goat are principally similar to those reported in human (16), rats (5,13), dog and cat (17) even though different studying methods were used. It has also been shown in this study that the median eminence in the goat was nourished by the internal and external capillaries plexi. These vascular plexi are also illustrated in the rat (4,18) and Rabbit (13). It is demonstrated in this study that the internal capillary plexus of median eminence forms numerous capillary loops protruding into the infundibular recess of the third ventricle. The Subependymal capillary networks located dorsally to these loops are derived from the arterioles which are branches of the hypophyseal arteries.

التغذية الدموية للمنطقة تحت المهادية في الماعز باستخدام تقنية القالب التآكلي

علاء عبد الخالق سواد
فرع التشريح، كلية الطب البيطري، جامعة البصرة، البصرة، العراق.

الخلاصة

استخدمت عشرة رؤوس ماعز بالغة من كلا الجنسين لدراسة التوزيع الدموي للمنطقة تحت المهادية. تجهز منطقة تحت المهاد بفروع من الشريان المخي الامامي وتفرعات الشرايين الموصلة الخلفية، إضافة الى فروع صغيرة من الشريان العيني، تضمنت التغذية الدموية للجسم الحلمي فروع من الشريان المخي الخلفي والشرايين الموصلة والفروع الانتهائي للشريان القاعدي. تعد منطقة تحت المهاد صغيرة الحجم مقارنة بقاعدة الدماغ الا انها تجهز بأوعية دموية غزيرة تتضمن تفرعات الشرايين المخية الامامية التي تكون دائرة ويلز مما يدل على الفعالية العالية لهذه المنطقة.

REFERENCES

- 1-Kingsely R.E. 2000. The hypothalamus and associated systems. In: concis Text of neuroscience, 2nd ed., pp 489-519.
- 2- Danial P.M. 1966. The blood supply of hypothalamus and the pituitary Gland. British medical bulletin, 3, 461-465.
- 3-Wislocki G.B., King L.S. 1936. The permeability of the hypophysis and hypothalamus to vital dyes, with a study of hypophyseal vascular supply. American Journal of Anatomy 58, 421-472.
- 4- Ambach G., Palkovits M. 1974. Blood supply of the rat hypothalamus. I. Nucleus supraopticus. Acta Morphologica, 22, 291-310.
- 5- Ambach G., Palkovits M. 1979. Blood supply of the rat hypothalamus. VI. Posterior region of the hypothalamus. Acta Morphologica, 27, 169-192.

- 6-Murakami T.,Kikuta A.,Taguchi T.,Ohtsuka A.,Ohtani O.1987.Blood vascular Architecture of the rat cerebral hypophysis and hypothalamus dissection – scanning electron microscopy of vascular cast.Archivum Histologicum Japonicum, 50,133-176.
- 7-murakami T.,Miyake T.,Ohtsuka A.,Kikuta A.,Taguchi T.,1993.Microcirculatory patterns in a dult rat cerebral hypophysis:a scanning electron microscope study of replicated specimens. Archives of Histology and Cytology ,56,243-260.
- 8-Negm I.M. 1971.The blood supply of the mouse hypophysis cerebri .Acta Anatomica ,80,377-387.
- 9-Gannon,B. J. 1978.Vascular casting.In: Principles and techniques of scanning electron microscopy. Biological Applications,6,Newyork.pp170-193.
- 10-Anonymous .2005.Nomina Anatomica Veterinaria.5th ed.International Committee on Veterinary Anatomical Nomenclature.Ithaca,New York.
- 11-Gillian L.A.1976.Extra- and intra –cranial blood supply to brains of dogs and cats. American Journal of Anatomy 146,237-253.
- 12-Lee r.m.1995.Morphology of cerebral arteries.Pharmacologyand Therapeutics.66,149-173.
- 13-Page R.B.,Munger B.L.,Bergland R.M.1976.Scanning electron microscopy of pituitary vascular casts. American Journal of Anatomy.146,273-301.
- 14-. Gillian L.A.1972.Blood supply to primitive mammalian brains.Journal of Comparative Neurology.145,209-221.
- 15-Dawson B.H. 1958. The blood vessels of human optic chiasma and their relation to those of the hypophysis and hypothalamus.Brain.81,207-217.
- 16-Popa G.T.,Fielding U.1930.A portal circulation from the pituitary to the hypothalamic region. Journal of Anatomy.65,88-91.
- 17-Wellens D.L.,Wouters L.J.,De Reese R.J.,Beirnaert P.,Reneman R.S. 1975.The cerebral blood distribution in dogs and cats. An anatomical and functional study. Brain Research. 86,429-438.
- 18-Harris G.W. 1972.Humor and hormones. Journal of Endocrinology. 53,i-xxii.