

2024

On $(K^*-N)_n$ -Quasi Normal Operators

Aya Mazin Falih

Ministry of Education, Iraq, aytmazin724@gmail.com

Follow this and additional works at: <https://qjps.researchcommons.org/home>



Part of the [Biology Commons](#), [Chemistry Commons](#), [Computer Sciences Commons](#), [Environmental Sciences Commons](#), [Geology Commons](#), [Mathematics Commons](#), and the [Nanotechnology Commons](#)

Recommended Citation

Falih, Aya Mazin (2024) "On $(K^*-N)_n$ -Quasi Normal Operators," *Al-Qadisiyah Journal of Pure Science*: Vol. 29 : No. 1 , Article 23.

Available at: <https://doi.org/10.29350/2411-3514.1261>

This Original Study is brought to you for free and open access by Al-Qadisiyah Journal of Pure Science. It has been accepted for inclusion in Al-Qadisiyah Journal of Pure Science by an authorized editor of Al-Qadisiyah Journal of Pure Science.

ARTICLE

On $(K^* - N)^n$ - Quasi Normal Operators

Aya M. Falih

Ministry of Education, Iraq

Abstract

The objective is to present a novel variant of a quasi-normal operator, specifically the $(K^* - N)^n$ quasi normal operator, alongside the introduction of related theorems, propositions, and illustrative examples elucidating this concept. Additionally, we present the necessary and sufficient conditions for addition and multiplication of this kind of operators.

Keywords: Operators, Normal operators, Quasi-normal operator, K^* quasi-normal operators

1. Introduction

A .Brow [1] introduced and researched the quasi-normal operator for the first time in 1953. Later Arun B. (1976) [2] provided several properties of the quasi-normal operator. Sh. Lohaj (2010) [3] introduced a novel variation of the quasi-normal operator, namely, N-quasi normal operators, along with presenting some of its fundamental characteristics. Ould A. (2011) submitted the n-power quasi-normal operator along with outlining its characteristics [4]. Valdete R.H. (2013) [5] provided several characteristics of N-quasi normal operators. Saad S. and Laith K (2015) [6] introduced the concept of the quasi-normal operator along with presenting some fundamental properties of this concept. Ahmed M and Salim D. (2017) [7] introduced another type called the $(K - N)^*$ quasi-normal operator, along with presenting some properties.

2. Basics

Definition (2.1). [8]

A bounded linear operator $T : \mathcal{H} \rightarrow \mathcal{H}$, is n operator satisfying $\|Tx\| \leq k\|x\| \quad \forall x \in \mathcal{H}$.

Definition (2.2). [8]

A bounded linear operator $T : \mathcal{H} \rightarrow \mathcal{H}$, is called normal if $TT^* = T^*T$.

Definition (2.3). [1]

A bounded linear operator $T : \mathcal{H} \rightarrow \mathcal{H}$ is quasi-normal if $T(T^*T) = (T^*T)T$.

Theorem (2.4). [2]

Let $T_1, T_2 : \mathcal{H} \rightarrow \mathcal{H}$ be quasi normal operators, such that $T_2T_1 = T_1T_2 = T_1^*T_2 = T_2T_1^* = 0$ then $T_1 + T_2$ is quasi normal.

Definition (2.5). [5]

A bounded linear operator $T : \mathcal{H} \rightarrow \mathcal{H}$ be a bounded linear is N-quasi normal if $T(T^*T) = N(T^*T)T$.

Theorem (2.6). [5]

Let $T_1 : \mathcal{H} \rightarrow \mathcal{H}$ be N-quasi normal and $T_2 : \mathcal{H} \rightarrow \mathcal{H}$ be quasi normal then the product T_1T_2 is N-quasi normal if they satisfy $T_1T_2 = T_2T_1$ and $T_2^*T_1 = T_1T_2^*$.

Definition (2.7). [4]

A bounded linear operator $T : \mathcal{H} \rightarrow \mathcal{H}$ is n-power quasi-normal if $T^n(T^*T) = (T^*T)T^n$.

Theorem (2.7). [9]

If $T : \mathcal{H} \rightarrow \mathcal{H}$ is n-power quasi-normal operator, then λT is also a n-power quasi-normal operator, where $\lambda \in \mathbb{R}$.

Definition (2.8). [10]

Let $T : \mathcal{H} \rightarrow \mathcal{H}$ is a bounded linear operator is said to be $(K-N)$ quasi-normal operator if $T^k(T^*T) = N(T^*T)T^k$.

Theorem (2.9). [10]

Let T_1 be (K-N) quasi normal operator and T_2 is k -quasi normal operator, such that, $(T_1T_2) = (T_2T_1)$ and $T_1T_2^* = T_2^*T_1$, then T_1T_2 is (K-N) quasi normal operator.

Theorem (2.10). [10]

Let $T_1, T_2 : \mathcal{H} \rightarrow \mathcal{H}$ be two (K-N) quasi normal operators such that $T_1^k T_2^* = T_2^* T_1^k = T_1^* T_2 = T_2^* T_1 = 0$ then $T_1 + T_2$ is (k-N) quasi normal operator.

Corollary (2.11). [10]

Let $T_1, T_2 : \mathcal{H} \rightarrow \mathcal{H}$ be two (K-N) quasi normal operators, such that $T_1^k T_2^* = T_2^* T_1^k = T_1^* T_2 = T_2^* T_1 = 0$ then $T_1 - T_2$ is (K-N) quasi normal operator.

3. Main results

Definition (3.1). Let $T : \mathcal{H} \rightarrow \mathcal{H}$ be a bounded linear operator then T is said to be $(K^* - N)^n$ quasi normal operator $((K^* - N)^n$ -QNO) if $(T^*)^K (T^* T)^n = N(T^* T)^n (T^*)^K$.

Example (3.2). $T = \begin{bmatrix} 0 & a \\ 0 & 0 \end{bmatrix}$ is $(K^* - N)^n$ -QNO where $k = 1, n = 1, N = I$

Propositions (3.3). Let $T : \mathcal{H} \rightarrow \mathcal{H}$ is $(K^* - N)^n$ -QNO then λT is $(K^* - N)^n$ -QNO where $\lambda \in \mathbb{R}$.

Proof.

$$((\lambda T)^*)^K ((\lambda T)^* (\lambda T))^n = (\lambda T^*)^K ((\lambda T^*) (\lambda T))^n$$

$$= \lambda^K (T^*)^K (\lambda T^*)^n (\lambda T)^n$$

$$= \lambda^K (T^*)^K \lambda^n (T^*)^n \lambda^n T^n$$

$$= \lambda^{K+2n} (T^*)^K (T^*)^n T^n$$

$$= \lambda^{K+2n} N(T^*)^n T^n (T^*)^K$$

$$= N \lambda^n (T^*)^n \lambda^n T^n \lambda^K (T^*)^K$$

$$= N(\lambda T^*)^n (\lambda T)^n (\lambda T^*)^K$$

$$= N((\lambda T)^* (\lambda T))^n ((\lambda T)^*)^K$$

$((\lambda T)^*)^K (\lambda T)^* (\lambda T)^n = N(\lambda T)^* (\lambda T)^n ((\lambda T)^*)^K$, hence the λT is $(K^* - N)^n$ -QNO ■.

Propositions (3.4). Let $T : \mathcal{H} \rightarrow \mathcal{H}$ is $(K^* - N)^n$ -QNO then T/M is $(K^* - N)^n$ -QNO where M is closed sub space.

$$\begin{aligned} & \left(\left(T/M \right)^* \right)^K \left(\left(T/M \right)^* (T/M) \right)^n \\ &= \left((T^*/M) \right)^K \left((T^*/M) (T/M) \right)^n \\ &= \left((T^*)^K / M \right) \left(T^* T / M \right)^n \\ &= \left((T^*)^K / M \right) \left((T^* T)^n / M \right) \\ &= \left(N(T^* T)^n (T^*)^K / M \right) \\ &= \left(N(T^* T)^n / M \right) \left((T^*)^K / M \right) \\ &= \left(N \left(\left(T^*/M \right) (T/M) \right)^n \right) \left((T^*/M) \right)^K \\ &= N \left(\left(T/M \right)^* (T/M) \right)^n \left(\left(T/M \right)^* \right)^K \end{aligned}$$

Then T/M is $(K^* - N)^n$ -QNO ■

Remark (3.5). Let T_1 and T_2 be two $(K^* - N)^n$ -QNO then $T_1 + T_2$ is not necessary $(K^* - N)^n$ -QNO. To illustrate that consider the following example.

Example (3.6).

$$T_1 = \begin{bmatrix} 2i & 0 & 2 \\ 0 & 0 & 0 \\ 2 & 0 & -2i \end{bmatrix} \text{ and } T_2 = \begin{bmatrix} 1 & 0 & i \\ 0 & 1 & 0 \\ i & 0 & 1 \end{bmatrix}$$

$$T_1 + T_2 = \begin{bmatrix} 1+2i & 0 & 2+i \\ 0 & 1 & 0 \\ 2 & 0 & -2i \end{bmatrix}$$

If $n = 1, K = 2$ and $N = I$.

$$\begin{aligned} & ((T_1 + T_2)^*)^2 ((T_1 + T_2)^* (T_1 + T_2))^1 \\ &= \begin{bmatrix} 32-56i & 0 & -24-52i \\ 0 & 1 & 0 \\ 40-20i & 0 & -40i \end{bmatrix} \end{aligned} \quad (1)$$

$$\begin{aligned}
 & ((T_1 + T_2)^*(T_1 + T_2))^1((T_1 + T_2)^*)^2 \\
 &= \begin{bmatrix} -120i & 0 & 40 - 20i \\ 0 & 1 & 0 \\ 104 - 52i & 0 & 32 + 24i \end{bmatrix} \quad (2)
 \end{aligned}$$

(1) $\neq$ (2)

Then $T_1 + T_2$ is not $(K^* - N)^n$ -QNO

Theorem (3.7). Let $T_1, T_2 : \mathcal{H} \rightarrow \mathcal{H}$ be $(K^* - N)^n$ -QNO on a Hilbert space s.t. $T_1^*T_2^* = T_1T_2 = T_1^*T_2 = 0$ then $T_1 + T_2$ is $(K^* - N)^n$ -QNO.

Proof. -

$$\begin{aligned}
 & (((T_1 + T_2)^*)^K((T_1 + T_2)^*(T_1 + T_2))^n) \\
 &= ((T_1^* + T_2^*)^K((T_1^* + T_2^*)(T_1 + T_2))^n) \\
 &= ((T_1^*)^K + K(T_1^*)^{K-1}T_2^* + \dots + (T_2^*)^K) \left((T_1^* + n(T_1^*)^{n-1}T_2^* \right. \\
 &\quad \left. + \dots + (T_2^*)^n)(T_1^n + nT_1^{n-1}T_2 + \dots + T_2^n) \right) = ((T_1^*)^K \\
 &\quad + (T_2^*)^K) (((T_1^*)^n + (T_2^*)^n)(T_1^n + T_2^n)) \\
 &= ((T_1^*)^K + (T_2^*)^K) \\
 &\quad \times ((T_1^*)^nT_1^n + (T_1^*)^nT_2^n + (T_2^*)^nT_1^n + (T_2^*)^nT_2^n) \\
 &= ((T_1^*)^K + (T_2^*)^K) ((T_1^*)^nT_1^n + (T_2^*)^nT_2^n) \\
 &= ((T_1^*)^K(T_1^*)^nT_1^n + (T_1^*)^K(T_2^*)^nT_2^n + (T_2^*)^K(T_1^*)^nT_1^n \\
 &\quad + (T_2^*)^K(T_2^*)^nT_2^n) \\
 &= ((T_1^*)^KT_1^*T_1^n + (T_2^*)^KT_2^*T_2^n)
 \end{aligned}$$

Since T_1 and T_2 are $(K^* - N)^n$ -QNO

$$= N((T_1^*)^nT_1^n(T_1^*)^K) + N((T_2^*)^nT_2^n(T_2^*)^K)$$

$$= N(((T_1^*)^nT_1^n(T_1^*)^K) + ((T_2^*)^nT_2^n(T_2^*)^K))$$

Therefore; $T_1 + T_2$ is $(K^* - N)^n$ -QNO ■

Remark (3.8). Let T_1 and T_2 be two $(K^* - N)^n$ -QNOs then T_1T_2 is not necessarily a $(K^* - N)^n$ -QNO as shown below.

Example (3.9).

$$T_1 = \begin{bmatrix} 2i & 0 & 2 \\ 0 & 0 & 0 \\ 2 & 0 & -2i \end{bmatrix} \text{ and } T_2 = \begin{bmatrix} 1 & 0 & i \\ 0 & 1 & 0 \\ i & 0 & 1 \end{bmatrix}$$

$$T_1T_2 = \begin{bmatrix} 4i & 0 & 0 \\ 0 & 0 & 0 \\ 4 & 0 & 0 \end{bmatrix}$$

If $n = 1, K = 2$ and $N = I$.

$$((T_1T_2)^*)^2((T_1T_2)^*(T_1T_2))^1 = \begin{bmatrix} -512 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \dots (1)$$

$$((T_1T_2)^*(T_1T_2))^1((T_1T_2)^*)^2 = \begin{bmatrix} -512 & 0 & -512i \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \dots (2)$$

(1) $\neq$ (2)

Then T_1T_2 is not $(K^* - N)^n$ -QNO

Theorem (3.10). Let T_1 and $T_2 : \mathcal{H} \rightarrow \mathcal{H}$ be $(K^* - N)^n$ -QNO and K^* quasi normal operator respectively such that, $T_1T_2 = T_2T_1$ and $T_2^*T_1 = T_1T_2^*$, then T_1T_2 is $(K^* - N)^n$ -QNOs.

Proof.

$$\begin{aligned}
 & ((T_1T_2)^*)^K((T_1T_2)^*(T_1T_2))^n \\
 &= ((T_2T_1)^*)^K((T_2^*T_1^n)(T_1T_2)^n) \\
 &= ((T_1^*T_2^*)^K)((T_1^*T_2^n)(T_1T_2)^n) \\
 &= ((T_1^*)^K(T_2^*)^K)((T_1^*)^n(T_2^*)^n)(T_1^nT_2^n) \\
 &= (T_1^*)^K((T_2^*)^K(T_1^*)^n)((T_2^*)^nT_1^nT_2^n) \\
 &= (T_1^*)^K((T_1^*)^n(T_2^*)^K)(T_1^n(T_2^*)^nT_2^n) \\
 &= ((T_1^*)^K(T_1^*)^n)((T_2^*)^KT_1^n)((T_2^*)^nT_2^n) \\
 &= ((T_1^*)^K(T_1^*)^n)(T_1^n(T_2^*)^K)((T_2^*)^nT_2^n) \\
 &= ((T_1^*)^K(T_1^*)^nT_1^n)((T_2^*)^K(T_2^*)^nT_2^n) \\
 &= N(T_1^*T_1^n(T_1^*)^K)((T_1^*)^nT_2^n(T_2^*)^K) \\
 &= N((T_1^*)^nT_1^n)((T_1^*)^K(T_2^*)^n)(T_2^n(T_2^*)^K) \\
 &= N((T_1^*)^nT_1^n)((T_2^*)^n(T_1^*)^K)(T_2^n(T_2^*)^K) \\
 &= N(T_1^*)^n(T_1^n(T_2^*)^n)((T_1^*)^KT_2^n)(T_2^*)^K
 \end{aligned}$$

$$\begin{aligned}
&= N(T_1^*)^n ((T_2^*)^n T_1^n) (T_2^n (T_1^*)^K) (T_2^*)^K \\
&= N(T_1^* T_2^*)^n (T_1^n T_2^n) ((T_1^*)^K (T_2^*)^K) \\
&= N((T_2 T_1)^*)^n (T_1 T_2)^n ((T_1^*) (T_2^*))^K \\
&= N((T_2 T_1)^*)^n (T_1 T_2)^n ((T_2 T_1)^*)^K \\
&= N((T_1 T_2)^*)^n (T_1 T_2)^n ((T_1 T_2)^*)^K \\
&((T_1 T_2)^*)^K ((T_1 T_2)^*)^n (T_1 T_2)^n \\
&= N((T_1 T_2)^*)^n (T_1 T_2)^n ((T_1 T_2)^*)^K
\end{aligned}$$

Therefore; the product $T_1 T_2$ is $(K^* - N)^n$ -QNO ■

References

- [1] Brown Arlen. On a class of operators. Am Math Soc 1953;4: 723–8.
- [2] Arun BALA. A note on quasi normal operators. Indian J Pure Appl Math 1977;8(4):463–8.
- [3] Lohaj S. Quasi-normal operators. Int J Math Anal 2010;4(47): 2311–20.
- [4] Ahmed OAMS, Ahmed MS. On the class of n-power quasi-normal operators on Hilbert space. Bull Math Anal Appl 2011;3(2):213–28.
- [5] Hamiti VR. Some properties of N-Quasinormal operators. Gen 2013;18(1):94–8.
- [6] Shaakir Laith K, Marai Saad S. quasi-normal Operator of order n. Tikrit J Pure Sci 2015;20(4):167–9.
- [7] Muhsin SD, Khalaf AM. On the class of (KN)* quasi-normal operators on hilbert space. Iraqi J Sci 2017:2172–6.
- [8] Sharma JN, Vasishtha AR. Functional analysis. 1st ed. Krishna Prakashan Media; 2013.
- [9] Veluchamy T, Manikandan KM. n-Power quasi normal operators on the hilbert space. IOSR J Math 2016;12:6–9.
- [10] Salim DM, Ahmed MK. On the class of (KN) quasi normal operator on Hilbert Space. Math Theor Model 2015;5(10).