

Ticks infesting some demestic animals in Thi-Qar Province, Southern Iraq

Sadek J. Tuama

College of Education, University of Thi-Qar

Khalid J.K. Al-Zihiry

Hayder K. Al-Maliky

College of Medicine, University of Thi-Qar

Abstract:

Four species of Ticks namely, *Hyalomma rufipes*, *H.trancatum*, *Boophilus annulatus* and *Ornithodoros savignyi* were isolated in present study from four species of domestic animals in Thi-Qar Province during the period from January 2002 until December 2004. Infestation percentage were 48.2%, 96.8%, 23% and 30% for Cattle, sheep, Goats and Chicken respectively. Seasonal variations were observed in the Sheep infestation with tick *H.rufipes*, the highest peak was cleared in July (87.8%), while the lowest was (10.2%) during February. The medical and economic importances of recorded ticks were discussed.

Introduction:

Ticks are belong to the subclass Acari, order Parasitiformis, and suborder Ixodida, which includes two major families, Ixodidae (hard ticks) with 13 genera and approximately 645 species and Argasidae (soft ticks) with 5 genera and approximately 170 species distributed worldwide (22), All are parasitic during some parts of their lives. The majority of them infest mammals, though many species attack bird and some found on cold-blooded animals (5). These arthropods are important to human through the direct effects of their feeding and as vectors of various agents of diseases in both man and livestock (6). Diseases transmitted by ticks to livestock constitute a major factor which limits animals production in many tropical and subtropical areas of the world (16), and responsible for high cattle mortalities compared to other diseases (3). Besides their role as disease control vectors, ticks cause physical damage such as injury to hides and loss of blood through their feeding (20).

In human, ticks can cause severe toxic condition such as paralysis and toxicosis, irritation and allergy, forty three species of ticks have been found to cause ticks paralysis with engorging adult female (15), moreover their ability to transmit a great variety of infectious diseases is a major public health concern, notably those of viral origin, characterized by encephalitis and hemorrhagic fever cause the highest morbidity and mortality(8).

Tick bites may also cause a severe allergic response, mediated by IgE specific for tick allergens (10). The insertion of the mouthpart (gnathosoma) into the skin produce an inflammatory reaction, and in many cases the mouthparts are torn off and left in the wound-such an injury

often becomes infected, producing an inflamed sore or ulcer (19).

The aim of present study is to gain informations of ticks in Iraq and to understand its seasonal dynamics.

Materials and Methods:

During the period from January 2002 to December 2004 a large numbers of ticks were collected from domestic animals (Cattle, Sheep, Goats, Chicken) from several localities in Thi-Qar province. Ticks were isolated by forceps, store in clean glass vials tightly lidded containing small pieces of cotton moistered with 1% mycostatin solution, to prevent desiccation and mould growth and labeled with date, location and name of host. The collected ticks were fixed and preserved into 70% ethylalcohol, as cited in Daniel (1993).

Ticks identification made based on six major features which are shape, size, mouthparts (capitulum), color, dorsal shield (scutum), and festoons (posterior abdominal markings). To determine the six major identifying features, the ticks compared with (1).

Results and Discussion:

1607 domestic animals were examined from different areas in Thi-Qar province which include: 319 Cattle, 946 Sheep, 162 Goats and 180 Chicken. Out of these animals 928(57.7%) were found be infected with four species of ticks identified as: *Hyalomma rufipes*, *H. truncatum*, *Boophilus annulatus* and *Ornithodoros savignyi* (Table 1).

Table 1: Infestation percentages of isolated ticks in present study.

Ticks	Cattle			SheeP		
	Exa.	Inf.	Per %	Exa.	Inf.	Per. %
Hyalomma	319	154	48.2	946	599	63.3
Rufipes		-	-		277	29.2
Hyalomma		-	-		3	0.3
Tnncaturfl		-	-		37	3.9
Boophilus		-	-			
Annulalus		-	-			
Ornithodor						
Os savignyi						

Exa. : Examined Inf.: Infected Per, : Percentage

Goats			Chicken		
Exa.	Inf.	Per.%	Exa.	Inf.	Per.%
162	-	-	180	-	-
	38	23.4		-	-
	-	-		6	6.6
	-	-		48	26.6

Hyalomma rufipes, H. truncatum:

H. rufipes were isolated from 753(46.8%) of examined animals distributed as 154(48.2%) from cattle and 599(63.3 %) from sheep, while H. truncatum isolated from 315(19.6 %) of examined animals, 277(29.2%) from sheep and 38(23.4%) from goats.

Ticks of the genus Hyalomma are well-known vectors of viruses and avid parasites of man. Although many species are not involved in disease transmission, the considerable length of Hyalomma mouthparts provokes a painful bite. One of the most important disease transmitted by this ticks is Crimean-Congo-Hemorrhagic Fever (CCHF) which occur sporadically throughout vast area of Africa, Asia and Europe, but can cause mortality(8), moreover the genus Theileria, is the most important being the causative agent of East Coast Fever in cattle (5), the economic impact of Theileriosis can be expressed in term of mortality, loss of production(live-wieght gain, milk production and draught potential)(4).(17) were divide this genus to three subgenus: Hyalomma, Hyalommina and Hyalommosta and givin the complete characteristics and classification keys.

Boophilus annulatus

This ticks were collected from 9(0.5%) of total number of animals, it is collected from 3(0.3%) sheeps and 6(3.3%) chicken.

Boophilus among the genera of ticks using large mammals as host and feed primarily on cattle, less frequently on other large herbivores (6) and constitute amajor problem for the cattle industry in tropical and subtropical areas of the world (9), but rarely attack man (23). B.annulatus still survives in the west Indies, Mexico and in the Mediterranean region, these arthropods transmit various viruses responsible for encephalitis which have a zoonotic origin, Babesia infection

also in *Boophilus* are transmitted from one generation to next transovarially(5).

Ornithodoros savignyi:

This ticks were collected from 85(5.2%) of total number of animals distributed as 37(3.91%) sheeps and 48(26.6%) chicken.

The widely distributed argasid genus *Ornithodoros* has several representative involved in the parasitism of human. A total of 22 species of this genus have been reported on human, and 12 species are found frequently, *O. savignyi* is found in human habitation in India, Africa and some parts of Asia (18) and cause intense local irritation (13), moreover in dry areas of these countries, the ticks is commonly attacks human resting under shady trees and around well where animals gather, etc. The bite of this species can cause long-lasting intense pruritus (8). Members of this genus has a characteristic full oval body shape(1), its larva is retained in the egg shell and become nymph a few hours after hatching, before partaking of their first meal(5). Experimental transmission by West Nile Virus(WNV) has been observed in *O.savignyi* (14) and virus antibodies have detected in human sera from Iraq(1). Specie of *Ornithodoros* can also harbor and transmit leptospiras(5).

Seasonal variations in infestation percentages of sheep with *Hyalomma rufipes*

The collection results of 732 sheep during one year (2004) as a monthly samples that the infestation were found with ticks *Hyalomma rufipes* in all months of this year in different percentages. These differents were calculated to determine the seasonal variations in this infestations (Table 2).

Table (2): Infestation percentages of sheep with ticks *Hyalomma rufipes* depending on monthly samples


Month	Examined	Infected	Percentages %
January	33	10	30.3
February	29	4	10.25
March	25	8	32
April	57	32	56.14
Mav	49	26	53
June	75	63	84
Juh	82	72	87.8
August	91	78	85.71
September	117	99	84.61
October	52	26	50
November	60	17	28.33
December	52	12	23
Total	732	447	61

The result of present study showed that the increase of infestation will observed at the ended of spring and started of summer until it is reached to high percentage in July

(87.8%), and then gradually decreased at the ended of summer and started of autumn reaching to the low level of infestation percentages in February (10.2%) (Figure 1).

This result were similar to these obtained by Stafford (2004) in infestation with ticks *Ixodes scapularis* in USA, which is attributed to that the nymph precede larvae and infect anew generation of animal host. Larvae active later in the summer then become infected when feeding on reservoir host animals.

Figure (1): Seasonal variations in infestations percentages of sheep with *Hyalomma rufipes*



References:

- 1-Adams, D.R.; Anderson, B.E.; Ammirati, C.T. and Helm, K.F.(2003).
Identification and diseases of common U.S. ticks. Internet J. Dermatol., 2 (1):
- 2-Anderson, J.F., Main, A.J., Andreadis, T.G., Wikel, S.K. and Vossbrinck, C.R.
(2003). Transstadial transfer of West Nile Virus by three species of Ixodid ticks
(Acari:
Ixodidae). J. Med. Entomol., 40(4): 528-533.
- 3-Anon, A.G.(1984). A practical field manual, Vol.2, Tick-borne disease control,
in: Ticks and tick-borne disease control, Rome, 457 PP.
- 4-de Castro, J.J., James, A.D., Minjauw, B., Di Giulio, G.U., Permin, A., Pegram,
R.G., Chizyuka, H.G. and Sinyangwe, P. (1997). Long term studies on the
economic impact of ticks on sanga cattle in Zambia. Exp.Appl.Acarol., 21: 3-19.
- 5-Cox, F.E.G.(1999). Modern parasitology.2nd edn., Blackwell science. 276 PP.
- 6-Cumming, G.S.(1998). Host preference in African ticks (Acari: Ixodida): a
quantitative data set. Bull. Entomol. Res., 88: 379-406.
- 7-Daniel, M. (1993). Influence of the microclimate on the vertical distribution of
the tick *Ixodes ricinus* (L.) in central Europe. Acarologia: 105-113.
- 8-Estrada-Pena, A. and Jongejan, F. (1999). Ticks feeding on humans: areview of
records on human-biting Ixodoidea with special references of pathogen
transmission. Exp. Appl. Acarol.23:685-715.
- 9-Fragoso, H.; Rad, P.H.; Ortiz, M. Rodriguez, M.; Redondo, M.; Herrera, L. 10
goloos has and de la Fuente, J. (1998). Protection against *Boophilus*.

annulatus infestations in cattle vaccinated with the B.microplus Bm86-containing vaccine Gavac.off.

Vaccine 16(20): 1990-1992. 815

10-Gauci, M.; Loh, R.K.; Stone, B.F. and Thong, Y.H.(1989). Allergic reaction to the Australian paralysis ticks, Ixodes holocyclus : diagnostic evaluation by skin test and radioimmunoassay. Clin.Exp.Allergy 19: 279-283.

11-Hassanain, M.A., El Garhy, M.F., Abdel-Ghaffar, F.A., El-Sharaby, A. and Abdel Megeed, K.N.(1997). Biological control studies of soft and hard ticks in Egypt I. The effect of Bacillus thuringiensis varieties on soft and hard ticks (Ixodidae). Parasitol. Res. 83: 209-213.

12-Hoogstraal, H. (1956). African Ixodoidea. I.Ticks of the Sudan (with special references to Equatoria province and with preliminary reviews of the genera Boophilus, Margaropus, Hyalomma. Dept. of the Navy, Bureau of medicine and surgery, Washington, D.C.

13-Hoogstraal, H. (1985). Argasid and Nuttallielid ticks as parasites and vectors. Adv. Parasitol., 24: 136-220.

14-Hubalek, Z. and Halouzka, J. (1999). West Nile Fever- a reemerging mosquito-borne viral diseases in Europe. Emer. Infec. Dis. 5 (5): 643-650.000)

15-John, D.T.; Petri, W.A. (2006). Markell and Voge's Medical Parasitology. 9th edn., Elsevier Inc., 463 pp.

16-Jongejan, F. and Uilenberg, G.(1994). Ticks and control methods. Rev. Sci. Tech. Off. Int. Epiz., 13(4): 1201-1226.

17-Kaiser, M.N. and Hoogstraal, H. (1967). Noteworthy bo recent tick records from Egypt. 2. Hosts, distribution and ecology of Rhipicephalus simus. J. Egyptian Public Health Assoc., 42: 231-242).

- 18-Keirans, J.E. (1984). George Henry Falkiner Nuttall ticks catalogue.U.S. Department of Agriculture Miscellaneous puplication, number 1438, pp.1-1785.
- 19-Ruy, J.S.; Lee, J.V.; Ahn, M.H.; Min, D.Y. and Ree, H.I. (1998). A human case of tick bite by Ixodes nipponensis. Korean J. Parasitol., 36: 59-61.
- 20-Smith, R., Pegram, R.G., Burt, S., Killorn, K.J., Oosterwijk, G., Paterson, A. and Wilsmore, A.J.(1991). Effect of dry season supplementation of sanga cattle in Zambia. Trop. Anim. Health prod., 23: 103-105.
- 21-Stafford, K.C. (2004). Tick management handbook. A integrated guide for homeowners, pest control operators, and public health officials for the prevention of tick-associated disease. CDC. 66 pp.
- 22-Strickland, G.T. (1991). Hunter's tropical medicine. 17end. Philadelphia. W.B. Saunders Company.
- 23-Strickland, R.K.; Gerrish, R.R.; Hourrigan, J.L. and Schubert, G.O. (1976). Ticks of veterinary importance. Animal and Plant Health Inspection. Service of the U.S. Dept. of Agric., Handbook. 485: 1-122.

مسح للقراد المتطفل على بعض الحيوانات المنزلية في "

محافظة ذي قار

صادق جعفر طعيمة

كلية التربية - جامعة ذي قار

خالد الزهيري

حيدر المالكي

كلية الطب , جامعة ذي قار

الخلاصة:

تم في الدراسة الحالية عزل اربعة انواع من القراد هي

Hyalomma rufipes, *H.tracatum*, *Boophilus annulatus* and *savignyi Ornithodoros* والتي اجريت على أربعة أنواع من الحيوانات المنزلية لعدة مناطق في محافظة ذي قار، وذلك اثناء المدة المحصورة من بداية شهر كانون الثاني عام 2002 ولغاية نهاية شهر كانون الأول عام 2004، وكانت نسبة الإصابة الكلية بالقراد هي 48.2 ، 96.8 ، 23 % و 30% في الأبقار ، الأغنام، الماعز والدجاج على التوالي ظهرت هناك تغيرات موسمية في نسبة إصابة الأغنام بالنوع الأول من القراد كانت قمته في شهر تموز (87.8) ، في حين كانت اقل نسبة اصابة في شهر شباط حيث بلغت (10.2%) ، فيما لم تظهر هناك دورات موسمية واضحة في بقية الأنواع . نوقشت في البحث الأهمية الصحية والاقتصادية لأنواع القراد المسجلة في الدراسة الحالية.